

THE QUANTUM GROUP OF A PREREGULAR MULTILINEAR FORM

Julien BICHON¹ and Michel DUBOIS-VIOLETTE²

Abstract

We describe the universal quantum group preserving a prerregular multilinear form, by means of an explicit finite presentation of the corresponding Hopf algebra.

Mathematics Subject Classification (2010). 16T05, 81R50, 15A69.

Keywords. Quantum group, multilinear form.

1 Introduction

The quantum group of a nondegenerate bilinear form (i.e. the largest quantum group preserving the bilinear form) was introduced and constructed in [14]. The aim of the paper is to generalize this construction to an appropriate class of m -linear forms ($m \geq 3$), the prerregular multilinear forms introduced in [11, 12].

Let w be a prerregular m -linear form on a vector space V (see Definition 1). We show the existence of a universal Hopf algebra $\mathcal{H}(w)$ preserving w and we provide an explicit *finite* presentation by generators and relations for $\mathcal{H}(w)$. It should be emphasized that the existence part is something that can be considered as well-known (see the end of Section 4) and that our main result is that we have obtained a finite presentation for the universal object $\mathcal{H}(w)$. The condition of prerregularity, introduced in [11, 12] in connection with the analysis of AS-regular algebras, turns out to be particularly relevant in the Hopf algebra context.

In view of the $m = 2$ case [4] and of the construction of the quantum group $SU(m)$ as the universal compact quantum group preserving a well-chosen m -linear form [19], it is natural to expect that the Hopf algebras $\mathcal{H}(w)$ should

¹Laboratoire de Mathématiques, Université Blaise Pascal, Campus des cézeaux BP 80026, F-63171 Aubière Cedex

Julien.Bichon@math.univ-bpclermont.fr

²Laboratoire de Physique Théorique UMR 8627, Université Paris XI, Bâtiment 210 F-91 405 Orsay Cedex

Michel.Dubois-Violette@u-psud.fr

produce quantum analogues of $SL(m)$. We examine two natural examples that show that this is far from being true if $m \geq 3$. The first one is when w is the “signature” m -form, for which we get a non commutative and non cosemisimple Hopf algebra, having the algebra of polynomial functions on $SL(m)$ as a quotient. This example also shows that the Hopf algebra $\mathcal{H}(w)$ differs in general from those Hopf algebras $\mathcal{H}(w, \tilde{w})$ constructed in [12]. The second example is when w is the “totally orthogonal” form, and we get some of the quantum reflection groups studied in [1], whose fusion rules are non-commutative.

The paper is organized as follows. Section 2 is devoted to preliminaries: we fix some notation and conventions and we recall the construction of the quantum group of a nondegenerate bilinear form. In Section 3 we discuss preregular multilinear forms. In Section 4 we describe the universal bialgebra preserving a multilinear form. In Section 5 we provide the explicit finite presentation for the universal Hopf algebra $\mathcal{H}(w)$ preserving a preregular multilinear form. In Section 6 we discuss the relation between $\mathcal{H}(w)$ and another class of Hopf algebras $\mathcal{H}(w, \tilde{w})$ introduced in [12]. Section 7 discusses some examples and the final Section 8 is our conclusion.

2 Preliminaries

2.1 Notations and conventions

Throughout this article \mathbb{K} is a (commutative) field and all vector spaces, algebras, etc. are over \mathbb{K} , the symbol \otimes denotes the tensor product over \mathbb{K} and we use the Einstein convention of summation over repeated up-down indices in the formulas. In the following V is a finite-dimensional vector space with $\dim(V) = n \geq 2$ equipped with a basis $(e_\lambda)_{\lambda \in \{1, \dots, n\}}$ and we endow the dual vector space V^* of V with the dual basis $(\theta^\lambda)_{\lambda \in \{1, \dots, n\}}$ of the basis (e_λ) . Using the finite-dimensionality of V we identify $(V^{\otimes m})^*$ with $(V^*)^{\otimes m}$ and thus, a m -linear form on V is the same thing as an element of $(V^*)^{\otimes m}$. Given an endomorphism Q of V , we denote by Q^t its transposed that is the endomorphism of V^* defined by $\langle \omega, QX \rangle = \langle Q^t \omega, X \rangle$ for any $\omega \in V^*$ and $X \in V$.

2.2 The quantum group of a non-degenerate bilinear form

Let $b = b_{\mu\nu}\theta^\mu \otimes \theta^\nu \in V^* \otimes V^* = (V \otimes V)^*$ be a nondegenerate bilinear form on V , ($b(e_\mu, e_\nu) = b_{\mu\nu}$). The quantum group of the nondegenerate bilinear form b was defined in [14] in the following manner. Denote by $b^{\mu\nu}$ the component of the inverse matrix of $(b_{\mu\nu}) \in M_n(\mathbb{K})$, i.e. one has

$$b^{\mu\lambda}b_{\lambda\nu} = b_{\nu\rho}b^{\rho\mu} = \delta_\nu^\mu \quad (2.1)$$

for $\mu, \nu \in \{1, \dots, n\}$.

Note the obvious fact that $\tilde{b} = b^{\mu\nu}e_\mu \otimes e_\nu \in V \otimes V = (V^* \otimes V^*)^*$ does not depend on the basis (e_λ) but only depends on $b \in V^* \otimes V^* = (V \otimes V)^*$.

Let $\mathcal{H}(b)$ be the unital associative algebra generated by the n^2 elements u_ρ^λ ($\lambda, \rho \in \{1, \dots, n\}$) with relations

$$b_{\mu\nu}u_\lambda^\mu u_\rho^\nu = b_{\lambda\rho}\mathbb{1} \quad (2.2)$$

$$b^{\mu\nu}u_\mu^\lambda u_\nu^\rho = b^{\lambda\rho}\mathbb{1} \quad (2.3)$$

for $\lambda, \rho \in \{1, \dots, n\}$. Then $\mathcal{H}(b)$ has a unique Hopf algebra structure with coproduct Δ , counit ε and antipode S such that

$$\Delta(u_\nu^\mu) = u_\lambda^\mu \otimes u_\nu^\lambda \quad (2.4)$$

$$\varepsilon(u_\nu^\mu) = \delta_\nu^\mu \quad (2.5)$$

$$S(u_\nu^\mu) = b^{\mu\lambda}u_\lambda^\rho b_{\rho\nu} \quad (2.6)$$

for $\mu, \nu \in \{1, \dots, n\}$. The dual object of $\mathcal{H}(b)$ is *the quantum group of the nondegenerate bilinear form b* . The analysis of the category of representations of this quantum group, that is of the category of corepresentations of $\mathcal{H}(b)$, has been done in [4].

Although not completely obvious with the above definition, the Hopf algebra $\mathcal{H}(b)$ is characterized by the following universal property.

Theorem 1. *Let H be a Hopf algebra which coacts on V as*

$$e_\lambda \mapsto e_\mu \otimes v_\lambda^\mu \quad (2.7)$$

$v_\lambda^\mu \in H$ and is such that

$$b_{\mu\nu} v_\lambda^\mu v_\rho^\nu = b_{\lambda\rho} \mathbb{1} \quad (2.8)$$

for $\lambda, \rho \in \{1, \dots, n\}$, (where $\mathbb{1}$ denotes the unit of H). Then there is a unique homomorphism of Hopf algebras $\varphi : \mathcal{H}(b) \rightarrow H$ such that $\varphi(u_\mu^\lambda) = v_\mu^\lambda$ for $\lambda, \mu \in \{1, \dots, n\}$.

In particular the smallest Hopf subalgebra of H which contains the v_μ^λ 's is a quotient of $\mathcal{H}(b)$. For the sake of completeness we give a proof (which will appear as a particular case of a more general result explained later, see Section 5).

Proof. Relations (2.8) imply

$$b^{\sigma\lambda} v_\lambda^\mu b_{\mu\nu} v_\rho^\nu = \delta_\rho^\sigma \mathbb{1}$$

which implies that the antipode S of H satisfies

$$S(v_\nu^\sigma) = b^{\sigma\lambda} v_\lambda^\mu b_{\mu\nu}$$

and therefore one has

$$v_\nu^\sigma S(v_\rho^\nu) = v_\nu^\sigma b^{\nu\lambda} v_\lambda^\mu b_{\mu\rho} = \delta_\rho^\sigma \mathbb{1}$$

from which one obtains by contraction with $b^{\sigma\tau}$

$$b^{\nu\lambda} v_\nu^\sigma v_\lambda^\tau = b^{\sigma\tau} \mathbb{1} \quad (2.9)$$

for any $\sigma, \tau \in \{1, \dots, n\}$. This implies together with (2.8) that $u_\nu^\mu \mapsto v_\nu^\mu$ defines a homomorphism of Hopf algebras of $\mathcal{H}(b)$ onto the Hopf subalgebra of H generated by the v_ν^μ . \square

3 Preregular multilinear forms

As in the previous section V is a finite-dimensional vector space with $\dim(V) = n \geq 2$, (e_λ) is a basis of V with dual basis (θ^λ) of the dual V^* of V , etc.. We shall always make the identifications $(V^{\otimes n})^* = V^{*\otimes n}$ which is allowed in view of the finite dimensionality of V . The following definition is taken from [11] (the main motivation was Theorem 4.3 there).

Definition 1. Let m be an integer with $m \geq 2$. A m -linear form w on V is said to be *preregular* iff it satisfies the following conditions (i) and (ii).

(i) If $X \in V$ is such that $w(X_1, \dots, X_{m-1}, X) = 0$ for any $X_1, \dots, X_{m-1} \in V$, then $X = 0$.

(ii) There is an element $Q_w \in GL(V)$ such that one has

$$w(X_1, \dots, X_{m-1}, X_m) = w(Q_w X_m, X_1, \dots, X_{m-1})$$

for any $X_1, \dots, X_m \in V$.

It follows from (i) that Q_w as in (ii) is unique. Condition (i) when combined with (ii) implies the stronger condition (i').

(i') For any $0 \leq k \leq m - 1$, if $X \in V$ is such that

$$w(X_1, \dots, X_k, X, X_{k+1}, \dots, X_{m-1}) = 0$$

for any $X_1, \dots, X_{m-1} \in V$, then $X = 0$.

Condition (i') will be referred to as *1-site nondegeneracy* while (ii) will be referred to as *twisted cyclicity* with *twisting* element Q_w . Thus a prerregular multilinear form is a multilinear form which is 1-site nondegenerate and twisted cyclic.

Basic examples of prerregular multilinear forms are: the signature form (see Section 7.1), its q -analogue (defined in [19]), the totally orthogonal form (see Section 7.2), and the multilinear forms appearing in §3.3 and §5.3 of [13] which correspond to representative examples of (regular) homogeneous algebras. For the latter class, it is worth noticing here that in the case of quadratic algebras, these multilinear forms give the noncommutative version of the volume element (see Proposition 10 of [12] and the comment following its proof).

By applying n times the relations of (ii) of Definition 1 one obtains the invariance of w by Q_w , that is

$$w(X_1, \dots, X_m) = w(Q_w X_1, \dots, Q_w X_m)$$

for any $X_1, \dots, X_m \in V$. Thus a twisted cyclic w is invariant by its twisting element.

Conversely, let w be an arbitrary Q -invariant m -linear form on V (with $Q \in GL(V)$) then the m -linear form $\pi_Q(w)$ defined by

$$\pi_Q(w)(X_1, \dots, X_m) = \sum_{k=1}^m w(QX_k, \dots, QX_m, X_1, \dots, X_{k-1})$$

for any $X_1, \dots, X_m \in V$ is twisted cyclic with twisting element Q , (in short is Q -cyclic). Notice that if \mathbb{K} is of characteristic 0, then $\frac{1}{m}\pi_Q$ is a projection of the Q -invariant m -linear forms onto the Q -twisted cyclic ones.

Condition (i) is equivalent to the existence of a m -linear form \tilde{w} on V^* such that one has

$$\tilde{w}^{\mu\lambda_1\dots\lambda_{m-1}} w_{\lambda_1\dots\lambda_{m-1}\nu} = \delta_\nu^\mu \quad (3.1)$$

for $\mu, \nu \in \{1, \dots, m\}$, where the components of \tilde{w} and w are defined by $\tilde{w}^{\lambda_1\dots\lambda_m} = \tilde{w}(\theta^{\lambda_1}, \dots, \theta^{\lambda_m})$ and $w_{\lambda_1\dots\lambda_m} = w(e_{\lambda_1}, \dots, e_{\lambda_m})$ for $\lambda_k \in \{1, \dots, m\}$. Notice that in contrast with the case $m = 2$, \tilde{w} is non unique for $m > 2$. The set of all \tilde{w} satisfying (3.1) as above is an affine subspace of $V^{\otimes m}$ which will be denoted by $\text{Aff}(w)$ and referred to as *the polar of w* .

In components, condition (ii) reads

$$w_{\lambda_1\dots\lambda_m} = Q_{\lambda_m}^\lambda w_{\lambda\lambda_1\dots\lambda_{m-1}} \quad (3.2)$$

for $\lambda_k \in \{1, \dots, m\}$, where the Q_ν^μ are the components of Q_w defined by

$$Q_w(e_\nu) = Q_\nu^\mu e_\mu \quad (3.3)$$

for $\nu \in \{1, \dots, n\}$ in the basis (e_λ) .

If one combines (3.1) with (3.2), one gets

$$Q_\nu^\rho \tilde{w}^{\mu\lambda_1\dots\lambda_{m-1}} w_{\rho\lambda_1\dots\lambda_{m-1}} = \delta_\nu^\mu$$

that is

$$\tilde{w}^{\mu\lambda_1\dots\lambda_{m-1}} w_{\nu\lambda_1\dots\lambda_{m-1}} = (Q^{-1})_\nu^\mu \quad (3.4)$$

for $\mu, \nu \in \{1, \dots, n\}$.

Let us say a few words on the case $m = 2$. It is clear that a bilinear form $w = b$ satisfying Condition (i) is just a nondegenerate bilinear form and

that then \tilde{w} as in 3.1 is unique and given by $\tilde{w}^{\mu\nu} = b^{\mu\nu}$ with the notations of Section 2. Hence a preregular bilinear form b is nondegenerate.

Conversely let b be a nondegenerate bilinear form with components $b_{\mu\nu}$ in the basis (e_λ) . Then b is twisted cyclic with

$$Q_\rho^\lambda = b^{\mu\lambda} b_{\mu\rho} \quad (3.5)$$

and hence a nondegenerate bilinear form is preregular.

The classification of nondegenerate bilinear forms over algebraically closed fields is known: the isomorphism class of a nondegenerate bilinear form only depends on the conjugacy class of its twisting element, see [16]. On the other hand, to the best of our knowledge, no general classification result is known for m -linear forms if $m \geq 3$ (isomorphism of multilinear forms is called congruence in [2]). The only basic observation we have is that isomorphic preregular m -linear forms have conjugate twisting elements.

4 The universal bialgebra $\mathcal{B}(w)$

Throughout this section w is a m -linear form on V satisfying Condition (i) of Definition 1. The components of w in the basis (e_λ) are denoted by $w_{\lambda_1 \dots \lambda_m} = w(e_{\lambda_1}, \dots, e_{\lambda_m})$ as before.

Let $\mathcal{B}(w)$ be the unital associative algebra generated by the n^2 elements a_ν^μ ($\mu, \nu \in \{1, \dots, n\}$) with relations

$$w_{\lambda_1 \dots \lambda_m} a_{\mu_1}^{\lambda_1} \dots a_{\mu_m}^{\lambda_m} = w_{\mu_1 \dots \mu_m} \mathbb{1} \quad (4.1)$$

for $\mu_k \in \{1, \dots, n\}$. Then $\mathcal{B}(w)$ has a unique bialgebra structure with co-product Δ and counit ε such that

$$\Delta(a_\nu^\mu) = a_\lambda^\mu \otimes a_\nu^\lambda \quad (4.2)$$

$$\varepsilon(a_\nu^\mu) = \delta_\nu^\mu \quad (4.3)$$

for $\mu, \nu \in \{1, \dots, n\}$. Let $\tilde{w} (\in V^{\otimes m})$ be, as in Section 2, a solution of (3.1). One has in view of (4.1)

$$\tilde{w}^{\mu\lambda_1 \dots \lambda_{m-1}} a_{\lambda_1}^{\rho_1} \dots a_{\lambda_{m-1}}^{\rho_{m-1}} w_{\rho_1 \dots \rho_{m-1} \sigma} a_\nu^\sigma = \delta_\nu^\mu \quad (4.4)$$

for $\mu, \nu \in \{1, \dots, n\}$, which means that in $M_n(\mathcal{B}(w))$ the matrix (a_ν^μ) has $(\tilde{w}^{\mu\lambda_1 \dots \lambda_{m-1}} a_{\lambda_1}^{\rho_1} \dots a_{\lambda_{m-1}}^{\rho_{m-1}} w_{\rho_1 \dots \rho_{m-1} \nu})$ as left inverse.

Theorem 2. *Let B be a bialgebra which coacts on V as*

$$e_\lambda \mapsto e_\mu \otimes v_\lambda^\mu$$

$v_\nu^\mu \in B$ and is such that

$$w_{\rho_1 \dots \rho_m} v_{\lambda_1}^{\rho_1} \dots v_{\lambda_m}^{\rho_m} = w_{\lambda_1 \dots \lambda_m} \mathbb{1}$$

for $\lambda_k \in \{1, \dots, n\}$, (where $\mathbb{1}$ is the unit of B). Then there is a unique homomorphism of bialgebras $\varphi : \mathcal{B}(w) \rightarrow B$ such that

$$\varphi(a_\nu^\mu) = v_\nu^\mu \quad \forall \mu, \nu \in \{1, \dots, n\}.$$

If furthermore B is a Hopf algebra with antipode S , then one has

$$S(v_\nu^\mu) = \tilde{w}^{\mu\lambda_1 \dots \lambda_{m-1}} v_{\lambda_1}^{\rho_1} \dots v_{\lambda_{m-1}}^{\rho_{m-1}} w_{\rho_1 \dots \rho_{m-1} \nu}$$

for $\mu, \nu \in \{1, \dots, n\}$, with \tilde{w} as above.

Proof. The first part is straightforward and classical in a more general context while the second part follows from

$$\begin{aligned} & \tilde{w}^{\mu\lambda_1 \dots \lambda_{m-1}} v_{\lambda_1}^{\rho_1} \dots v_{\lambda_{m-1}}^{\rho_{m-1}} w_{\rho_1 \dots \rho_{m-1} \sigma} v_\tau^\sigma S(v_\nu^\tau) \\ &= S(v_\nu^\mu) = \tilde{w}^{\mu\lambda_1 \dots \lambda_{m-1}} v_{\lambda_1}^{\rho_1} \dots v_{\lambda_{m-1}}^{\rho_{m-1}} w_{\rho_1 \dots \rho_{m-1} \nu} \end{aligned}$$

for $\mu, \nu \in \{1, \dots, n\}$ which follows from (4.4) and from the definition of the antipode. \square

Hence by Manin's construction of the Hopf envelope of a bialgebra [15], there exists a universal Hopf algebra coacting on a multilinear form w , which is finitely generated as an algebra since w satisfies condition (i) in Definition 1 (by the last identities of the previous theorem). We have not been able to provide a finite presentation of this algebra without using the full definition of preregularity.

5 The universal Hopf algebra $\mathcal{H}(w)$

Throughout this section w is a given preregular multilinear form on V with twisting element Q_w simply denoted by $Q \in GL(V)$.

The components of w, Q and Q^{-1} in the basis (e_λ) of V (with dual basis (θ^λ) of V^*) are denoted by $w_{\lambda_1 \dots \lambda_m}, Q_\nu^\mu$ and $(Q^{-1})_\nu^\mu$ with

$$w_{\lambda_1 \dots \lambda_m} = w(e_{\lambda_1}, \dots, e_{\lambda_m})$$

$$Q(e_\nu) = Q_\nu^\mu e_\mu, \quad \text{or} \quad Q_\nu^\mu = \langle \theta^\mu, Q(e_\nu) \rangle$$

$$Q^{-1}(e_\nu) = (Q^{-1})_\nu^\mu e_\mu, \quad \text{or} \quad (Q^{-1})_\nu^\mu = \langle \theta^\mu, Q^{-1}(e_\nu) \rangle$$

for $\lambda_k, \mu, \nu \in \{1, \dots, n\}$.

Let $\mathcal{H}(w)$ be the unital associative algebra generated by the $2n^2$ generators u_ν^μ, s_ν^μ with relations

$$u_\lambda^\mu s_\nu^\lambda = \delta_\nu^\mu \mathbf{1} \tag{5.1}$$

$$Q_\nu^\lambda u_\lambda^\rho (Q^{-1})_\rho^\sigma s_\sigma^\mu = \delta_\nu^\mu \mathbf{1} \tag{5.2}$$

$$w_{\lambda_1 \dots \lambda_m} u_{\mu_1}^{\lambda_1} \dots u_{\mu_m}^{\lambda_m} = w_{\mu_1 \dots \mu_m} \mathbf{1} \tag{5.3}$$

for $\mu, \nu, \mu_k \in \{1, \dots, n\}$.

Proposition 3. *One has in $\mathcal{H}(w)$*

$$w_{\mu_1 \dots \mu_m} s_{\nu_m}^{\mu_m} \dots s_{\nu_1}^{\mu_1} = w_{\nu_1 \dots \nu_m} \mathbf{1} \tag{5.4}$$

$$s_\rho^\mu u_\nu^\rho = \delta_\nu^\mu \mathbf{1} \tag{5.5}$$

$$s_\nu^\lambda Q_\lambda^\tau u_\tau^\rho (Q^{-1})_\rho^\mu = \delta_\nu^\mu \mathbf{1} \tag{5.6}$$

for $\nu_k, \mu, \nu \in \{1, \dots, n\}$. Furthermore, for any solution \tilde{w} of (3.1) i.e. $\forall \tilde{w} \in \text{Aff}(w)$ one has

$$s_\nu^\mu = \tilde{w}^{\mu \lambda_1 \dots \lambda_{m-1}} u_{\lambda_1}^{\rho_1} \dots u_{\lambda_{m-1}}^{\rho_{m-1}} w_{\rho_1 \dots \rho_{m-1} \nu} \tag{5.7}$$

$$Q_\nu^\tau u_\tau^\rho (Q^{-1})_\rho^\mu = \tilde{w}^{\mu \lambda_1 \dots \lambda_{m-1}} s_{\lambda_{m-1}}^{\rho_{m-1}} \dots s_{\lambda_1}^{\rho_1} w_{\rho_1 \dots \rho_{m-1} \nu} \tag{5.8}$$

for $\mu, \nu \in \{1, \dots, n\}$.

Proof. Relations (5.4) are obtained from (5.3) by contraction on the right by $s_{\nu_m}^{\mu_m} \dots s_{\nu_1}^{\mu_1}$ and by using (5.1). From (3.1) and (5.3) it follows that

$$(\tilde{w}^{\mu\lambda_1 \dots \lambda_{m-1}} u_{\lambda_1}^{\rho_1} \dots u_{\lambda_{m-1}}^{\rho_{m-1}} w_{\rho_1 \dots \rho_{m-1} \sigma}) u_{\nu}^{\sigma} = \delta_{\nu}^{\mu} \mathbf{1}$$

which implies (5.7) in view of (5.1) and then (5.5) follows (see also Theorem 2). Similarly, from (3.1) and (5.4) it follows that

$$s_{\nu}^{\sigma} (\tilde{w}^{\mu\lambda_1 \dots \lambda_{m-1}} s_{\lambda_{m-1}}^{\rho_{m-1}} s_{\lambda_1}^{\rho_1} w_{\rho_1 \dots \rho_{m-1} \sigma}) = \delta_{\nu}^{\mu} \mathbf{1}$$

which implies (5.8) in view of (5.2) and then (5.6) follows. \square

Proposition 4. *The algebra $\mathcal{H}(w)$ has a unique structure of Hopf algebra with coproduct Δ , counit ε and antipode S such that*

$$\Delta(u_{\nu}^{\mu}) = u_{\lambda}^{\mu} \otimes u_{\nu}^{\lambda} \quad (5.9)$$

$$\Delta(s_{\nu}^{\mu}) = s_{\nu}^{\lambda} \otimes s_{\lambda}^{\mu} \quad (5.10)$$

$$\varepsilon(u_{\nu}^{\mu}) = \varepsilon(s_{\nu}^{\mu}) = \delta_{\nu}^{\mu} \quad (5.11)$$

$$S(u_{\nu}^{\mu}) = s_{\nu}^{\mu} \quad (5.12)$$

$$S(s_{\nu}^{\mu}) = (Q^{-1})_{\rho}^{\mu} u_{\lambda}^{\rho} Q_{\nu}^{\lambda} \quad (5.13)$$

for $\mu, \nu \in \{1, \dots, n\}$.

Proof. This is a direct verification using the relations from the previous proposition. \square

We did not use Condition (ii) of Definition 1 (twisted cyclicity) to construct the Hopf algebra $\mathcal{H}(w)$ (Q could be any invertible matrix). The twisted cyclicity condition is fully used to prove the following universal property of $\mathcal{H}(w)$.

Theorem 5. *Let w be prerregular m -linear form and let H be a Hopf algebra which coacts on V as*

$$e_{\lambda} \mapsto e_{\mu} \otimes v_{\lambda}^{\mu}$$

$v_{\lambda}^{\mu} \in H$ and is such that

$$w_{\mu_1 \dots \mu_m} v_{\lambda_1}^{\mu_1} \dots v_{\lambda_m}^{\mu_m} = w_{\lambda_1 \dots \lambda_m} \mathbf{1}$$

for $\lambda_k \in \{1, \dots, n\}$, (where $\mathbf{1}$ denotes the unit of H). Then there is a unique homomorphism of Hopf algebras $\varphi : \mathcal{H}(w) \rightarrow H$ such that $\varphi(u_{\mu}^{\lambda}) = v_{\mu}^{\lambda}$ for $\lambda, \mu \in \{1, \dots, n\}$.

This theorem is the announced generalization of Theorem 1. It follows that if $w = b$ is bilinear, then $\mathcal{H}(b)$ as in Section 2 is isomorphic with $\mathcal{H}(w)$ as above.

Proof. All what one has to verify is that the relations (5.1), (5.2) and (5.3) are satisfied with u_ν^μ replaced by v_ν^μ and $s_\nu^\mu = S(u_\nu^\mu)$ replaced by $S(v_\nu^\mu)$, (S being the antipode). Relation (5.3) is satisfied by definition and relation (5.1) is a consequence of the axioms for Hopf algebras. Thus it remains to show that

$$Q_\nu^\lambda v_\lambda^\rho (Q^{-1})_\rho^\sigma S(v_\sigma^\mu) = \delta_\nu^\mu \mathbb{1} \quad (5.14)$$

is satisfied in H . On the other hand, Theorem 2 implies that

$$S(v_\nu^\mu) = \tilde{w}^{\mu\lambda_1 \dots \lambda_{m-1}} v_{\lambda_1}^{\rho_1} \dots v_{\lambda_{m-1}}^{\rho_{m-1}} w_{\rho_1 \dots \rho_{m-1} \nu}$$

with \tilde{w} solution of (3.1). Therefore one has

$$\begin{aligned} Q_\nu^\lambda v_\lambda^\rho (Q^{-1})_\rho^\sigma S(v_\sigma^\mu) &= Q_\nu^\lambda v_\lambda^\rho (Q^{-1})_\rho^\sigma \tilde{w}^{\mu\lambda_1 \dots \lambda_{m-1}} v_{\lambda_1}^{\rho_1} \dots v_{\lambda_{m-1}}^{\rho_{m-1}} w_{\rho_1 \dots \rho_{m-1} \sigma} \\ &= \tilde{w}^{\mu\lambda_1 \dots \lambda_{m-1}} Q_\nu^\lambda v_\lambda^\rho v_{\lambda_1}^{\rho_1} \dots v_{\lambda_{m-1}}^{\rho_{m-1}} w_{\rho_1 \dots \rho_{m-1} \nu} \\ &= \tilde{w}^{\mu\lambda_1 \dots \lambda_{m-1}} Q_\nu^\lambda w_{\lambda \lambda_1 \dots \lambda_{m-1}} \mathbb{1} \\ &= \tilde{w}^{\mu\lambda_1 \dots \lambda_{m-1}} w_{\lambda_1 \dots \lambda_{m-1} \nu} \mathbb{1} = \delta_\nu^\mu \mathbb{1} \end{aligned}$$

where the twisted cyclicity condition is used in the second equality. This proves (5.14). \square

It follows in particular from Theorem 5 that the algebra of polynomial functions on $\text{Aut}(w)$ is a quotient of $\mathcal{H}(w)$. An element of $\text{Aut}(w)$ has to commute with the twisting element Q , so if Q is diagonalizable with distinct eigenvalues, $\text{Aut}(w)$ is very small, being a subgroup of the torus $(K^*)^m$. This still allows $\mathcal{H}(w)$ to be a large algebra (and hence w has a large quantum symmetry group), as the example of quantum $SL(2)$ shows.

6 The Hopf algebras $\mathcal{H}(w, \tilde{w})$, $\tilde{w} \in \text{Aff}(w)$

In this section, where w is a preregular m -linear form on V with twisting element $Q \in GL(V)$ and \tilde{w} is an element of $\text{Aff}(w)$ (that is an element of $V^{\otimes m}$ satisfying (3.1)), we recall the definition of the Hopf algebra $\mathcal{H}(w, \tilde{w})$, constructed in [12], and relate it to $\mathcal{H}(w)$.

Let $\mathcal{H}(w, \tilde{w})$ be the unital associative algebra generated by the n^2 elements v_ν^μ ($\mu, \nu \in \{1, \dots, n\}$) with relations

$$w_{\lambda_1 \dots \lambda_m} v_{\mu_1}^{\lambda_1} \dots v_{\mu_m}^{\lambda_m} = w_{\mu_1 \dots \mu_m} \mathbb{1} \quad (6.1)$$

and

$$\tilde{w}^{\lambda_1 \dots \lambda_m} v_{\lambda_1}^{\mu_1} \dots v_{\lambda_m}^{\mu_m} = \tilde{w}^{\mu_1 \dots \mu_m} \mathbb{1} \quad (6.2)$$

for $\mu_k \in \{1, \dots, n\}$. There is a unique structure of Hopf algebra on $\mathcal{H}(w, \tilde{w})$ with coproduct Δ , counit ε and antipode S such that

$$\Delta v_\nu^\mu = v_\lambda^\mu \otimes v_\nu^\lambda \quad (6.3)$$

$$\varepsilon(v_\nu^\mu) = \delta_\nu^\mu \quad (6.4)$$

and

$$S(v_\nu^\mu) = \tilde{w}^{\mu \lambda_1 \dots \lambda_{m-1}} v_{\lambda_1}^{\rho_1} \dots v_{\lambda_{m-1}}^{\rho_{m-1}} w_{\rho_1 \dots \rho_{m-1} \nu} \quad (6.5)$$

for $\mu, \nu \in \{1, \dots, n\}$.

By Theorem 5 there exists a unique (surjective) homomorphism of Hopf algebras

$$\varphi : \mathcal{H}(w) \rightarrow \mathcal{H}(w, \tilde{w})$$

such that $\varphi(u_\nu^\mu) = v_\nu^\mu$ for $\mu, \nu \in \{1, \dots, n\}$. We will see in the next section that this is not an isomorphism in general.

7 Examples

In this section, in order to illustrate our construction, we examine two examples. We begin with a general useful lemma, whose proof is straightforward combining Relations (5.4) and (5.5).

Lemma 6. *Assume that $m \geq 3$. We have, in $\mathcal{H}(w)$,*

$$w_{\lambda \rho \mu_3 \dots \mu_m} s_{\nu_m}^{\mu_m} \dots s_{\nu_3}^{\mu_3} = w_{\nu_1 \dots \nu_m} u_\lambda^{\nu_1} u_\rho^{\nu_2}$$

for any $\lambda, \rho, \nu_3, \dots, \nu_m \in \{1, \dots, n\}$.

7.1 Example 1: the signature form

Let $w = \varepsilon$ be the signature form on $V = \mathbb{K}^m$ (the volume form), i.e. $\varepsilon_{\lambda_1 \dots \lambda_m} = 0$ if two indices are equal, and $\varepsilon_{\lambda_1 \dots \lambda_m}$ is the signature of the corresponding permutation otherwise. This is a preregular m -linear form, with $Q = I_m$.

Proposition 7. *The Hopf algebra $\mathcal{H}(\varepsilon)$ is not commutative if $m \geq 3$.*

Proof. It is easy to see that it is enough to prove the result at $m = 3$. Let \mathcal{A} be the free algebra on two generators x and y . It is straightforward to check the existence of an algebra map $\mathcal{H}(w) \rightarrow \mathcal{A}$

$$\begin{pmatrix} u_1^1 & u_2^1 & u_3^1 \\ u_1^2 & u_2^2 & u_3^2 \\ u_1^3 & u_2^3 & u_3^3 \end{pmatrix} \mapsto \begin{pmatrix} 1 & x & y \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} s_1^1 & s_2^1 & s_3^1 \\ s_1^2 & s_2^2 & s_3^2 \\ s_1^3 & s_2^3 & s_3^3 \end{pmatrix} \mapsto \begin{pmatrix} 1 & -x & -y \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

Thus $\mathcal{H}(\varepsilon)$ is not commutative. \square

Using Lemma 6 we see that the following relations hold in $\mathcal{H}(\varepsilon)$:

$$u_\nu^\lambda u_\nu^\mu = u_\nu^\mu u_\nu^\lambda, \quad [u_\nu^\lambda, u_\rho^\mu] = [u_\nu^\mu, u_\rho^\lambda]$$

for any λ, μ, ν, ρ . These are the relations defining the Manin matrices studied in [6].

Let $\tilde{\varepsilon}$ be given by $\tilde{\varepsilon}_{\lambda_1 \dots \lambda_m} = \frac{1}{m} \varepsilon_{\lambda_1 \dots \lambda_m}$ (assuming here that $m \neq 0$ in \mathbb{K}). We have $\tilde{\varepsilon} \in \text{Aff}(\varepsilon)$.

Proposition 8. *Assume that $m \geq 3$. Then the canonical Hopf algebra map $\mathcal{H}(\varepsilon) \rightarrow \mathcal{H}(\varepsilon, \tilde{\varepsilon})$ is not injective, and $\mathcal{H}(\varepsilon)$ is not cosemisimple.*

Proof. One can show that $\mathcal{H}(\varepsilon, \tilde{\varepsilon})$ is commutative, as in [19, 17]. It thus follows that the canonical Hopf algebra map $\mathcal{H}(\varepsilon) \rightarrow \mathcal{H}(\varepsilon, \tilde{\varepsilon})$ is not injective.

Assume that $\mathcal{H}(\varepsilon)$ is cosemisimple. Then there exists $w' \in \text{Aff}(\varepsilon)$ satisfying (6.2). But $\tilde{\varepsilon}$ is, up to scalar, the only element in $\text{Aff}(\varepsilon)$ for which (6.2) can hold in $\mathcal{H}(\varepsilon, \tilde{\varepsilon})$ (by some well-known facts on the representation theory of $SL(m)$). Thus, up to a scalar, $w' = \tilde{\varepsilon}$, and relations (6.2) hold for $\tilde{\varepsilon}$ in $\mathcal{H}(\varepsilon)$. Thus the canonical map $\mathcal{H}(\varepsilon) \rightarrow \mathcal{H}(\varepsilon, \tilde{\varepsilon})$ is an isomorphism: contradiction. \square

7.2 Example 2: the totally orthogonal form

We now consider the totally orthogonal m -linear form θ on $V = \mathbb{K}^n$: $\theta_{\lambda_1 \dots \lambda_m} = 1$ if all indices are equal and 0 otherwise. This is a preregular m -linear form with $Q = I_n$.

Proposition 9. *For $m \geq 3$, the Hopf algebra $\mathcal{H}(\theta)$ is isomorphic to the algebra presented by generators a_λ^μ , $\mu, \lambda \in \{1, \dots, n\}$, and relations*

$$a_\lambda^\mu a_\nu^\mu = 0 = a_\mu^\lambda a_\mu^\nu, \quad \forall \mu, \forall \lambda \neq \nu$$

$$\sum_{\lambda=1}^n (a_\lambda^\mu)^m = 1 = \sum_{\lambda=1}^n (a_\mu^\lambda)^m, \quad \forall \mu$$

Thus $\mathcal{H}(\theta)$ is isomorphic with the Hopf algebra denoted $A_h^m(n)$ in [1].

Proof. Let \mathcal{A} denote the algebra having the presentation given in the statement of the proposition. It is a direct verification to check that there exists a unique Hopf algebra structure on \mathcal{A} such that

$$\Delta(a_\nu^\mu) = a_\lambda^\mu \otimes a_\nu^\lambda, \quad \varepsilon(a_\nu^\mu) = \delta_\nu^\mu, \quad S(a_\nu^\mu) = (a_\mu^\nu)^{m-1}$$

and that \mathcal{A} satisfies the assumption in Theorem 5. Hence there exists a surjective Hopf algebra map $\mathcal{H}(\theta) \rightarrow \mathcal{A}$, $u_\lambda^\mu \mapsto a_\lambda^\mu$.

To construct the inverse isomorphism, we have to check that the defining relations in \mathcal{A} are satisfied in $\mathcal{H}(\theta)$. The second family of relations holds in $\mathcal{H}(\theta)$ by Relations (5.3). We have, by Lemma 6, $u_\lambda^\mu u_\nu^\mu = 0$ for $\lambda \neq \nu$. Using the antipode we get $s_\lambda^\mu s_\nu^\mu = 0$ for $\lambda \neq \nu$. Moreover, using Relations (5.3), we see that $s_\mu^\lambda = (u_\lambda^\mu)^{m-1}$, hence $s_\mu^\lambda s_\mu^\nu = 0$ for $\lambda \neq \nu$, and using the antipode, we see that the first family of relations holds in $\mathcal{H}(\theta)$. We get a surjective algebra map $\mathcal{A} \rightarrow \mathcal{H}(\theta)$, $a_\lambda^\mu \mapsto u_\lambda^\mu$, and we are done. \square

The Hopf algebra $\mathcal{H}(\theta) \simeq A_h^m(n)$ (which also can be described by a free wreath product operation [5]) is, when $\mathbb{K} = \mathbb{C}$, cosemimple and its fusion rules are non-abelian, see [1]. It would be interesting to characterize the preregular multilinear forms for which the diagrammatic techniques in [1] can be extended.

8 Conclusion : The quantum group of a pre-regular multilinear form

The Hopf algebra $\mathcal{H}(w, \tilde{w})$ which was considered in [12] in connection with its coaction on the algebras $\mathcal{A}(w, N)$ defined by the “twisted potential” w (for $m \geq N \geq 2$) has the defect to be dependent of the auxiliary variable $\tilde{w} \in \text{Aff}(w)$, (excepted for $m = 2$ for which $\text{card}(\text{Aff}(w)) = 1$).

The Hopf algebra $\mathcal{H}(w)$ coacts as well on the algebras $\mathcal{A}(w, N)$ defined in Section 5 of [12]. This latter class of algebras includes all AS-regular algebras which are N -Koszul as shown in [12], Theorem 11 (see also in [11]).

It is clear from Theorem 5 that the right Hopf algebra which only depends on w is the universal Hopf algebra $\mathcal{H}(w)$ defined in Section 5. Therefore we define *the quantum group of the preregular multilinear form w* to be the dual object of $\mathcal{H}(w)$. The *representations* of this quantum group are the corepresentations of the Hopf algebra $\mathcal{H}(w)$ which plays the role of the Hopf algebra of “representative functions” on it.

In fact the Hopf algebras $\mathcal{H}(w, \tilde{w})$ for $\tilde{w} \in \text{Aff}(w)$ all are quotients of $\mathcal{H}(w)$, which means that the corresponding quantum groups are quantum subgroups of the above quantum group of the preregular multilinear form w . In the case $m = 2$, that is when w is a nondegenerate bilinear form, then \tilde{w} is unique and $\mathcal{H}(w, \tilde{w})$ coincides with $\mathcal{H}(w)$ as explained in Section 2 (Theorem 1). Thus for $m = 2$, the canonical projection

$$\pi : \mathcal{H}(w) \rightarrow \mathcal{H}(w, \tilde{w})$$

is an isomorphism. However for $m \geq 3$, Example 1 of the previous section shows that this not always true, but Example 2 shows it can be true. Those examples also show that the uniform description of the corepresentation category of $\mathcal{H}(w)$ for the case $m = 2$ in [4] will no longer hold when $m \geq 3$.

The Hopf algebras $\mathcal{H}(w)$ and $\mathcal{H}(w, \tilde{w})$ above belong to a particular class of Hopf algebras named cosovereign Hopf algebras in [3] where they were considered and studied in a categorical perspective. Examples of Hopf algebras of this type are those arising from compact quantum groups [18]. The class of cosovereign Hopf algebras is also exactly the class of Hopf algebras

considered in [7] (see also in [10]) and the analysis of [7] was later extended to a larger class of Hopf algebras, which mixed the notion of cosovereign Hopf algebra with the dual notion of sovereign Hopf algebra [3], in [8] and [9]. These classes of Hopf algebras generalize the class of Hopf algebras with antipode satisfying $S^2 = I$ for which cyclic cohomology and the generalization of the Weil homomorphism have tractable descriptions.

References

- [1] T. Banica, R. Vergnioux, Fusion rules for quantum reflection groups. *J. Noncommut. Geom.* 3 (2009), 327-359.
- [2] G. Belitskii, V. Sergeichuk, Congruence of multilinear forms, *Linear Algebra Appl.* 418 (2006), 751-762.
- [3] J. Bichon, Cosovereign Hopf algebras, *J. Pure Appl. Algebra* **157**, No.2-3 (2001), 121-133.
- [4] J. Bichon, The representation category of the quantum group of a non-degenerate bilinear form, *Comm. Algebra* **31**, No. 10 (2003), 4831-4851.
- [5] J. Bichon, Free wreath product by the quantum permutation group, *Algebr. Represent. Theory* 7 (2004), 343-362.
- [6] A. Chervov, G. Falqui, V. Rubtsov, Algebraic properties of Manin matrices. I, *Adv. in Appl. Math.* 43 (2009), no. 3, 239-315.
- [7] A. Connes and H. Moscovici, Hopf algebras, cyclic cohomology and the transverse index theorem, *Comm. Math. Phys.* 198 (1998), 199-246.
- [8] A. Connes and H. Moscovici, Cyclic cohomology and Hopf algebras, *Lett. Math. Phys.* 48 (1999), 97-108.
- [9] A. Connes and H. Moscovici, Cyclic cohomology and Hopf algebra symmetry, *Lett. Math. Phys.* 53 (2000), 1-28.
- [10] M. Crainic, Cyclic cohomology of Hopf algebras. *J. Pure Appl. Algebra* 166 (2002), 29-66.
- [11] M. Dubois-Violette, Graded algebras and multilinear forms, *C. R. Acad. Sci. Paris, Ser. I* **341** (2005), 719-724.

- [12] M. Dubois-Violette, Multilinear forms and graded algebras, *J. Algebra* **317** (2007), 198-225.
- [13] M. Dubois-Violette. Noncommutative coordinate algebras. In E. Blanchard, editor, *Quanta of Maths, dédié à A. Connes*, volume 14 of *Clay Mathematics Proceedings*, pages 171–199. Clay Mathematics Institute, 2010.
- [14] M. Dubois-Violette, G. Launer, The quantum group of a non-degenerate bilinear form, *Phys. Lett. B* **245**, No.2 (1990), 175-177.
- [15] Yu. I. Manin, Quantum groups and noncommutative geometry *Université de Montréal, Centre de Recherches Mathématiques, Montréal, QC*, 1988.
- [16] C. Riehm, The equivalence of bilinear forms, *J. Algebra* 31 (1974), 45-66.
- [17] M. Rosso, Algèbres enveloppantes quantifiées, groupes quantiques compacts de matrices et calcul différentiel non commutatif, *Duke Math. J.* 61 (1990), 11-40.
- [18] S.L. Woronowicz, Compact matrix pseudogroups, *Comm. Math. Phys.* 111 (1987), 613-665.
- [19] S.L. Woronowicz, Tannaka-Krein duality for compact matrix pseudogroups. Twisted SU(N) groups, *Invent. Math.* 93 (1988), 35-76.