

Approximation of control and inverse problems for PDEs using variational methods: A review

ARNAUD DIEGO MÜNCH

Université Clermont Auvergne - Clermont-Ferrand - FRANCE

Workshop on "Recent advances in PDEs: Analysis, Numerics and Control"

In honor of Enrique Fernández-Cara for his 60th birthday

Sevilla - January 25th-27th 2017



General Context

We discuss **hyperbolic** and **parabolic** equations and try to emphasize the interest of **space-time variational methods** with respect to time marching methods.

Boundary controllability of wave like equation with initial data in $L^2 \times H^{-1}$

$\Omega \subset \mathbb{R}^N$ bounded domain with C^2 -boundary; $T > 0$; $Q_T := \Omega \times (0, T)$; $d \in L^\infty(Q_T)$;
 $\Gamma_0 \subset \partial\Omega$

$$\begin{cases} Ly := y_{tt} - \Delta y + dy = 0, & Q_T := \Omega \times (0, T), \\ y = \nu 1_{\Gamma_0}(x), & \Sigma_T := \partial\Omega \times (0, T), \\ (y(\cdot, 0), y_t(\cdot, 0)) = (y_0, y_1) \in \mathbf{H} := L^2(\Omega) \times H^{-1}(\Omega), & \Omega. \end{cases} \quad (1)$$

EXISTENCE - UNIQUENESS (Lions'69)

$\forall \nu \in L^2(\Sigma_T)$, $\exists! y = y(\nu) \in C^0([0, T]; L^2(\Omega)) \cap C^1([0, T]; H^{-1}(\Omega))$ and

$$\|y\|_{L^\infty(0, T; L^2(\Omega))} \leq C_{\Omega, T} \left(\|y_0, y_1\|_{\mathbf{H}} + \|\nu\|_{L^2(\Sigma_T)} \right)$$

NULL CONTROLLABILITY (Lions'88, Bardos-Lebeau-Rauch'92, Lasiecka'94, ...) If (T, Γ_0, Ω) satisfies a geometric condition, system (1) is **null controllable** at time T uniformly with respect to the initial condition (y_0, y_1) : $\exists \nu \in L^2(\Sigma_T)$ such that

$$(y_\nu(\cdot, T), (y_\nu)_t(\cdot, T)) = (0, 0), \quad \text{in } \Omega. \quad (2)$$

Boundary controllability of wave like equation with initial data in $L^2 \times H^{-1}$

$\Omega \subset \mathbb{R}^N$ bounded domain with C^2 -boundary; $T > 0$; $Q_T := \Omega \times (0, T)$; $d \in L^\infty(Q_T)$;
 $\Gamma_0 \subset \partial\Omega$

$$\begin{cases} Ly := y_{tt} - \Delta y + dy = 0, & Q_T := \Omega \times (0, T), \\ y = \mathbf{v} 1_{\Gamma_0}(x), & \Sigma_T := \partial\Omega \times (0, T), \\ (y(\cdot, 0), y_t(\cdot, 0)) = (y_0, y_1) \in \mathbf{H} := L^2(\Omega) \times H^{-1}(\Omega), & \Omega. \end{cases} \quad (1)$$

EXISTENCE - UNIQUENESS (Lions'69)

$\forall \mathbf{v} \in L^2(\Sigma_T)$, $\exists! y = y(\mathbf{v}) \in C^0([0, T]; L^2(\Omega)) \cap C^1([0, T]; H^{-1}(\Omega))$ and

$$\|y\|_{L^\infty(0, T; L^2(\Omega))} \leq C_{\Omega, T} \left(\|y_0, y_1\|_{\mathbf{H}} + \|\mathbf{v}\|_{L^2(\Sigma_T)} \right)$$

NULL CONTROLLABILITY (Lions'88, Bardos-Lebeau-Rauch'92, Lasiecka'94, ...) If (T, Γ_0, Ω) satisfies a geometric condition, system (1) is **null controllable** at time T uniformly with respect to the initial condition (y_0, y_1) : $\exists \mathbf{v} \in L^2(\Sigma_T)$ such that

$$(y_{\mathbf{v}}(\cdot, T), (y_{\mathbf{v}})_t(\cdot, T)) = (0, 0), \quad \text{in } \Omega. \quad (2)$$

Observability for the adjoint system

The controllability property of the hyperbolic equation is equivalent to the observability for the corresponding adjoint problem :

$$\begin{cases} L^* \varphi := \varphi_{tt} - \Delta \varphi + d\varphi = 0 & \text{in } Q_T, \\ \varphi = 0 & \text{on } \Sigma_T, \\ (\varphi(\cdot, T), \varphi_t(\cdot, T)) = (\varphi_0, \varphi_1) \in \mathbf{V} & \text{in } \Omega \end{cases} \quad (3)$$

$$\mathbf{V} := H_0^1(\Omega) \times L^2(\Omega) = \mathbf{H}'.$$

OBSERVABILITY INEQUALITY- System (3) is **observable in time T** if there exists a positive constant $C_{obs} > 0$ such that

$$\|(\varphi_0, \varphi_1)\|_{\mathbf{V}}^2 \leq C_{obs} \int_0^T \int_{\Gamma_0} \left| \frac{\partial \varphi}{\partial \nu} \right|^2 d\sigma dt \quad \forall (\varphi_0, \varphi_1) \in \mathbf{V}. \quad (4)$$

$C_{obs} = C_{obs}(T, \Gamma_0, \Omega, \|d\|_{L^\infty(Q_T)})$ - Observability constant

Minimal L^2 -norm control

$$\begin{cases} \text{Minimize } J(y, v) = \frac{1}{2} \int_0^T \int_{\Gamma_0} |v|^2 d\sigma dt \\ \text{Subject to } (y, v) \in \mathcal{C}(y_0, y_1; T) \end{cases} \quad (5)$$

where $\mathcal{C}(y_0, y_1; T)$ denotes the non-empty linear manifold

$$\mathcal{C}(y_0, y_1; T) = \{ (y, v) : v \in L^2(\Sigma_T), y \text{ solves (1) and satisfies (2)} \}.$$

Using the [Fenchel-Rockafellar theorem \[Ekeland-Temam 74\], \[Brezis 84\]](#) we get that

$$\inf_{(y, v) \in \mathcal{C}(y_0, y_1; T)} J(y, v) = - \min_{(\varphi_0, \varphi_1) \in V} J^*(\varphi_0, \varphi_1)$$

$$\begin{cases} \text{Minimize } J^*(\varphi_0, \varphi_1) = \frac{1}{2} \int_0^T \int_{\Gamma_0} \left| \frac{\partial \varphi}{\partial \nu} \right|^2 d\sigma dt + \langle (\varphi_0, \varphi_1), (-y_1, y_0) \rangle_{V, H} \\ \text{Subject to } (\varphi_0, \varphi_1) \in V \text{ where } L^* \varphi = 0 \quad + IC + BC \end{cases} \quad (6)$$

Optimal control: $v = \frac{\partial \varphi}{\partial \nu} 1_{\Gamma_0}$

Minimal L^2 -norm control

$$\begin{cases} \text{Minimize } J(y, v) = \frac{1}{2} \int_0^T \int_{\Gamma_0} |v|^2 d\sigma dt \\ \text{Subject to } (y, v) \in \mathcal{C}(y_0, y_1; T) \end{cases} \quad (5)$$

where $\mathcal{C}(y_0, y_1; T)$ denotes the non-empty linear manifold

$$\mathcal{C}(y_0, y_1; T) = \{ (y, v) : v \in L^2(\Sigma_T), y \text{ solves (1) and satisfies (2)} \}.$$

Using the **Fenchel-Rockafellar theorem** [Ekeland-Temam 74], [Brezis 84] we get that

$$\inf_{(y, v) \in \mathcal{C}(y_0, y_1; T)} J(y, v) = - \min_{(\varphi_0, \varphi_1) \in \mathbf{V}} J^*(\varphi_0, \varphi_1)$$

$$\begin{cases} \text{Minimize } J^*(\varphi_0, \varphi_1) = \frac{1}{2} \int_0^T \int_{\Gamma_0} \left| \frac{\partial \varphi}{\partial \nu} \right|^2 d\sigma dt + \langle (\varphi_0, \varphi_1), (-y_1, y_0) \rangle_{\mathbf{V}, \mathbf{H}} \\ \text{Subject to } (\varphi_0, \varphi_1) \in \mathbf{V} \text{ where } L^* \varphi = 0 \quad + IC + BC \end{cases} \quad (6)$$


Optimal control: $v = \frac{\partial \varphi}{\partial \nu} \mathbf{1}_{\Gamma_0}$

Approximation and minimization of J^* over $\mathbf{V} := H_0^1(\Omega) \times L^2(\Omega)$

$$\begin{cases} \text{Min } J^*(\varphi_0, \varphi_1) = \frac{1}{2} \int_0^T \int_{\Gamma_0} \left| \frac{\partial \varphi}{\partial \nu} \right|^2 d\sigma dt + \langle (\varphi_0, \varphi_1), (-y_1, y_0) \rangle_{\mathbf{V}, \mathbf{H}} \\ \text{Subject to } (\varphi_0, \varphi_1) \in \mathbf{V} = H_0^1(\Omega) \times L^2(\Omega) \text{ where } L^* \varphi = 0 \quad + IC + BC \end{cases}$$

At least **two methods** to solve this extremal problem:

- ▶ The "bad" one : minimize J^* w.r.t. (φ_0, φ_1) by a gradient method. This requires to solve $L^* \varphi = 0$ by a time marching method . However, it is not possible to achieve at the finite dimensional level the constraint $L^* \varphi = 0$!!!! The "trick", developed initially by Glowinski ¹ is first to discretize the equation and then to exactly control the corresponding finite dimensional system.




¹R. Glowinski, C.H. Li, J.-L. Lions, A numerical approach to the exact boundary controllability of the wave equation, (1990)

Approximation and minimization of J^* over $\mathbf{V} := H_0^1(\Omega) \times L^2(\Omega)$

$$\begin{cases} \text{Min } J^*(\varphi_0, \varphi_1) = \frac{1}{2} \int_0^T \int_{\Gamma_0} \left| \frac{\partial \varphi}{\partial \nu} \right|^2 d\sigma dt + \langle (\varphi_0, \varphi_1), (-y_1, y_0) \rangle_{\mathbf{V}, \mathbf{H}} \\ \text{Subject to } (\varphi_0, \varphi_1) \in \mathbf{V} = H_0^1(\Omega) \times L^2(\Omega) \text{ where } L^* \varphi = 0 + IC + BC \end{cases}$$

At least **two methods** to solve this extremal problem:

- The "bad" one : minimize J^* w.r.t. (φ_0, φ_1) by a gradient method. This requires to solve $L^* \varphi = 0$ by a time marching method .
However, it is not possible to achieve at the finite dimensional level the constraint $L^* \varphi = 0$!!!! The "trick", developed initially by Glowinski ¹ is **first** to discretize the equation and **then to exactly control** the corresponding finite dimensional system.




¹R. Glowinski, C.H. Li, J.-L. Lions, A numerical approach to the exact boundary controllability of the wave equation, (1990)

Approximation and minimization of J^* over $\mathbf{V} := H_0^1(\Omega) \times L^2(\Omega)$

$$\begin{cases} \text{Min } J^*(\varphi_0, \varphi_1) = \frac{1}{2} \int_0^T \int_{\Gamma_0} \left| \frac{\partial \varphi}{\partial \nu} \right|^2 d\sigma dt + \langle (\varphi_0, \varphi_1), (-y_1, y_0) \rangle_{\mathbf{V}, \mathbf{H}} \\ \text{Subject to } (\varphi_0, \varphi_1) \in \mathbf{V} = H_0^1(\Omega) \times L^2(\Omega) \text{ where } L^* \varphi = 0 + IC + BC \end{cases}$$

At least **two methods** to solve this extremal problem:

- ▶ The "bad" one : minimize J^* w.r.t. (φ_0, φ_1) by a gradient method. This requires to solve $L^* \varphi = 0$ by a time marching method .
However, it is not possible to achieve at the finite dimensional level the constraint $L^* \varphi = 0$!!!! The "trick", developed initially by Glowinski ¹ is **first** to discretize the equation and **then to exactly control** the corresponding finite dimensional system.




¹R. Glowinski, C.H. Li, J.-L. Lions, A numerical approach to the exact boundary controllability of the wave equation, (1990)

Approximation and minimization of J^* over $\mathbf{V} := H_0^1(\Omega) \times L^2(\Omega)$

$$\begin{cases} \text{Min } J^*(\varphi_0, \varphi_1) = \frac{1}{2} \int_0^T \int_{\Gamma_0} \left| \frac{\partial \varphi}{\partial \nu} \right|^2 d\sigma dt + \langle (\varphi_0, \varphi_1), (-y_1, y_0) \rangle_{\mathbf{V}, \mathbf{H}} \\ \text{Subject to } (\varphi_0, \varphi_1) \in \mathbf{V} = H_0^1(\Omega) \times L^2(\Omega) \text{ where } L^* \varphi = 0 + IC + BC \end{cases}$$

At least **two methods** to solve this extremal problem:

- The "bad" one : minimize J^* w.r.t. (φ_0, φ_1) by a gradient method. This requires to solve $L^* \varphi = 0$ by a time marching method .
However, it is not possible to achieve at the finite dimensional level the constraint $L^* \varphi = 0$!!!! The "trick", developed initially by Glowinski ¹ is **first** to discretize the equation and **then to exactly control** the corresponding finite dimensional system.



¹R. Glowinski, C.H. Li, J.-L. Lions, A numerical approach to the exact boundary controllability of the wave equation, (1990)


1D - Negative Commutation diagram

CENTERED FINITE DIFFERENCE IN SPACE AND TIME - UNIFORM DISCRETIZATION -

Constant coefficients $c := 1$, $d := 0$

$$(\bar{S}_{h,\Delta t}) \begin{cases} \Delta_{\Delta t} y_{h,\Delta t} - \Delta_h y_{h,\Delta t} = 0, \\ + \text{Initial and boundary terms} \end{cases} \quad (7)$$

produces **a non discrete uniformly bounded and convergent control** under the (CFL) condition $\Delta t < h$.



For high frequency components of the discrete solution, the discrete observability constant $C_{obs,h}$ blows up as $h \rightarrow 0$

[Glowinski-Lions'90] then [Zuazua team later].

Numerical example

$$\Omega = (0, 1) - \Gamma_0 = \{1\} - T = 2.4$$

$$y_0(x) = \begin{cases} 16x & x \in [0, 1/2], \\ 0 & x \in]1/2, 1]. \end{cases} ; \quad y_1(x) = 0. \quad (8)$$

The control v with minimal L^2 -norm is discontinuous :

$$v(t) = \begin{cases} 0 & t \in [0, 0.9] \cup [1.9, T], \\ 8(t - 1.4) & t \in]0.9, 1.9[, \end{cases} \quad (9)$$

leading to $\|v\|_{L^2(0,T)} = 4/\sqrt{3} \approx 2.3094$.

Usual centered finite difference scheme - Discrete control





Figure: Control $P(\mathbf{v}_h)(t)$ vs. $t \in [0, T]$, $\Delta t/h = 0.98$, $T = 2.4$ and $h = 1/10, 1/20, 1/30$ and $h = 1/40$.

1D - Positive Commutation diagram with a modified scheme

2

$$(\bar{S}_{h,\Delta t}) \left\{ \begin{array}{l} \Delta_{\Delta t} y_{h,\Delta t} + \frac{1}{4}(h^2 - \Delta t^2) \underbrace{\Delta_h \Delta_{\Delta t} y_{h,\Delta t}}_{(\Delta y)_{tt}} - \Delta_h y_{h,\Delta t} = 0, \\ + \text{Initial conditions and Boundary terms} \end{array} \right. \quad (10)$$

produces a discrete uniformly bounded and converging control under the condition $\Delta t < h\sqrt{T/2}$.




Within this approach (discretize then control), remedies in the general case (general domain, non constant coefficients) are unknown.

1D - Positive Commutation diagram with a modified scheme

2

$$(\bar{S}_{h,\Delta t}) \left\{ \begin{array}{l} \Delta_{\Delta t} y_{h,\Delta t} + \frac{1}{4}(h^2 - \Delta t^2) \underbrace{\Delta_h \Delta_{\Delta t} y_{h,\Delta t}}_{(\Delta y)_{tt}} - \Delta_h y_{h,\Delta t} = 0, \\ + \text{Initial conditions and Boundary terms} \end{array} \right. \quad (10)$$

produces a discrete uniformly bounded and converging control under the condition $\Delta t < h\sqrt{T/2}$.



Within this approach (discretize then control), remedies in the general case (general domain, non constant coefficients) are unknown.

Modified scheme - control




Figure: Modified scheme - Control $P(\mathbf{v}_h)(t)$ vs. $t \in [0, T]$ - $\Delta t = 1.095445h$, $T = 2.4$ and $h = 1/20, 1/40, 1/80, 1/160$.

Second method to bypass the fact that $L^* \varphi_h \neq 0$: the "good" one

- ▶ Minimize the conjugate functional J^* w.r.t. φ directly ! Mainly use by the German optimal control community. **This allows to relax the constraint $L^* \varphi_h = 0$!**

We replace the observability inequality

$$\begin{cases} \|\varphi_0, \varphi_1\|_{\mathbf{V}}^2 \leq C_{obs} \left\| \frac{\partial \varphi}{\partial \nu} \right\|_{L^2(\Gamma_T)}^2, & \forall (\varphi_0, \varphi_1), \\ L^* \varphi = 0, & \varphi|_{\Sigma_T} = 0 \end{cases} \quad (11)$$

by a "generalized observability inequality" :

$$\|\varphi(\cdot, 0), \varphi_t(\cdot, 0)\|_{\mathbf{V}}^2 \leq C_{\Omega, T}(1 + C_{obs}) \left(\left\| \frac{\partial \varphi}{\partial \nu} \right\|_{L^2(\Gamma_T)}^2 + \|L^* \varphi\|_{L^2(Q_T)}^2 \right), \quad \forall \varphi \in \Phi \quad (12)$$

Advantages ? If $\varphi_h \in \Phi_h$ a finite dimensional subspace of Φ , then

$$\|\varphi_h(\cdot, 0), \varphi_{h,t}(\cdot, 0)\|_{\mathbf{V}}^2 \leq C_{\Omega, T}(1 + C_{obs}) \left(\left\| \frac{\partial \varphi_h}{\partial \nu} \right\|_{L^2(\Gamma_T)}^2 + \|L^* \varphi_h\|_{L^2(Q_T)}^2 \right), \quad \forall \varphi_h \in \Phi_h \quad (13)$$

with a constant $C_{\Omega, T}(1 + C_{obs})$ independent of h !!!

Second method to bypass the fact that $L^* \varphi_h \neq 0$: the "good" one

- Minimize the conjugate functional J^* w.r.t. φ directly ! Mainly use by the German optimal control community. **This allows to relax the constraint $L^* \varphi_h = 0$!**

We replace the observability inequality

$$\begin{cases} \|\varphi_0, \varphi_1\|_{\mathbf{V}}^2 \leq C_{obs} \left\| \frac{\partial \varphi}{\partial \nu} \right\|_{L^2(\Gamma_T)}^2, & \forall (\varphi_0, \varphi_1), \\ L^* \varphi = 0, \quad \varphi|_{\Sigma_T} = 0 \end{cases} \quad (11)$$

by a "generalized observability inequality" :

$$\|\varphi(\cdot, 0), \varphi_t(\cdot, 0)\|_{\mathbf{V}}^2 \leq C_{\Omega, T}(1 + C_{obs}) \left(\left\| \frac{\partial \varphi}{\partial \nu} \right\|_{L^2(\Gamma_T)}^2 + \|L^* \varphi\|_{L^2(Q_T)}^2 \right), \quad \forall \varphi \in \Phi \quad (12)$$

Advantages ? If $\varphi_h \in \Phi_h$ a finite dimensional subspace of Φ , then

$$\|\varphi_h(\cdot, 0), \varphi_{h,t}(\cdot, 0)\|_{\mathbf{V}}^2 \leq C_{\Omega, T}(1 + C_{obs}) \left(\left\| \frac{\partial \varphi_h}{\partial \nu} \right\|_{L^2(\Gamma_T)}^2 + \|L^* \varphi_h\|_{L^2(Q_T)}^2 \right), \quad \forall \varphi_h \in \Phi_h \quad (13)$$

with a constant $C_{\Omega, T}(1 + C_{obs})$ independent of h !!!

Second method to bypass the fact that $L^* \varphi_h \neq 0$: the "good" one

- ▶ Minimize the conjugate functional J^* w.r.t. φ directly ! Mainly use by the German optimal control community. **This allows to relax the constraint $L^* \varphi_h = 0$!**

We replace the observability inequality

$$\begin{cases} \|\varphi_0, \varphi_1\|_{\mathbf{V}}^2 \leq C_{obs} \left\| \frac{\partial \varphi}{\partial \nu} \right\|_{L^2(\Gamma_T)}^2, & \forall (\varphi_0, \varphi_1), \\ L^* \varphi = 0, & \varphi|_{\Sigma_T} = 0 \end{cases} \quad (11)$$

by a "generalized observability inequality" :

$$\|\varphi(\cdot, 0), \varphi_t(\cdot, 0)\|_{\mathbf{V}}^2 \leq C_{\Omega, T}(1 + C_{obs}) \left(\left\| \frac{\partial \varphi}{\partial \nu} \right\|_{L^2(\Gamma_T)}^2 + \|L^* \varphi\|_{L^2(Q_T)}^2 \right), \quad \forall \varphi \in \Phi \quad (12)$$

Advantages ? If $\varphi_h \in \Phi_h$ a finite dimensional subspace of Φ , then

$$\|\varphi_h(\cdot, 0), \varphi_{h,t}(\cdot, 0)\|_{\mathbf{V}}^2 \leq C_{\Omega, T}(1 + C_{obs}) \left(\left\| \frac{\partial \varphi_h}{\partial \nu} \right\|_{L^2(\Gamma_T)}^2 + \|L^* \varphi_h\|_{L^2(Q_T)}^2 \right), \quad \forall \varphi_h \in \Phi_h \quad (13)$$

with a constant $C_{\Omega, T}(1 + C_{obs})$ independent of h !!!

Minimization of J^* w.r.t. φ

We replace the problem

$$\begin{cases} \text{Min } J^*(\varphi_0, \varphi_1) = \frac{1}{2} \int_0^T \int_{\Gamma_0} \left| \frac{\partial \varphi}{\partial \nu} \right|^2 d\sigma dt + \langle y_0, \varphi_t(\cdot, 0) \rangle_{L^2} - \langle y_1, \varphi(\cdot, 0) \rangle_{H^{-1}, H_0^1} \\ \text{Subject to } (\varphi_0, \varphi_1) \in \mathbf{V} = H_0^1(\Omega) \times L^2(\Omega) \quad \text{where } L^* \varphi = 0 \end{cases} \quad (14)$$

by the equivalent problem

$$\begin{cases} \min J^*(\varphi) = \frac{1}{2} \int_0^T \int_{\Gamma_0} \left| \frac{\partial \varphi}{\partial \nu} \right|^2 d\sigma dt + \langle y_0, \varphi_t(\cdot, 0) \rangle_{L^2} - \langle y_1, \varphi(\cdot, 0) \rangle_{H^{-1}, H_0^1} \\ \text{Subject to } \varphi \in \mathbf{W} := \left\{ \varphi : \varphi \in C^0(0, T; H_0^1(\Omega)) \cap C^1(0, T; L^2(\Omega)), L^* \varphi = 0 \in L^2(Q_T) \right\} \end{cases} \quad (15)$$

Remark- \mathbf{W} endowed with the norm $\|\varphi\|_{\mathbf{W}} := \left\| \frac{\partial \varphi}{\partial \nu} \right\|_{L^2(\Gamma_T)}$ is an Hilbert space.

Minimization of J^*

We now replace the problem

$$\begin{cases} \text{Min } J^*(\varphi_0, \varphi_1) = \frac{1}{2} \int_0^T \int_{\Gamma_0} \left| \frac{\partial \varphi}{\partial \nu} \right|^2 d\sigma dt + \langle y_0, \varphi_t(\cdot, 0) \rangle_{L^2} - \langle y_1, \varphi(\cdot, 0) \rangle_{H^{-1}, H_0^1} \\ \text{Subject to } (\varphi_0, \varphi_1) \in \mathbf{V} = H_0^1(\Omega) \times L^2(\Omega) \quad \text{where } L^* \varphi = 0 \end{cases} \quad (16)$$

by the equivalent problem

$$\begin{cases} \min J_r^*(\varphi) = \frac{1}{2} \int_0^T \int_{\Gamma_0} \left| \frac{\partial \varphi}{\partial \nu} \right|^2 d\sigma dt + \frac{r}{2} \|L^* \varphi\|_{L^2(Q_T)}^2 + \langle y_0, \varphi_t(\cdot, 0) \rangle_{L^2} - \langle y_1, \varphi(\cdot, 0) \rangle_{H^{-1}, H_0^1} \\ \text{Subject to } \varphi \in \mathbf{W} := \left\{ \varphi : \varphi \in C^0(0, T; H_0^1(\Omega)) \cap C^1(0, T; L^2(\Omega)), L^* \varphi = 0 \in L^2(Q_T) \right\} \end{cases} \quad (17)$$

for all $r \geq 0$.

Remark- \mathbf{W} endowed with the norm $\|\varphi\|_{\mathbf{W}} := \left\| \frac{\partial \varphi}{\partial \nu} \right\|_{L^2(\Gamma_T)}$ is an Hilbert space.

Relaxation of $L^*\varphi = 0$ via a Lagrange Multiplier

In order to address the $L^2(Q_T)$ constraint $L^*\varphi = 0$, we introduce a Lagrange multiplier $\lambda \in L^2(Q_T)$; we consider the saddle point problem³:

$$\left\{ \begin{array}{l} \sup_{\lambda \in L^2(Q_T)} \inf_{\varphi \in \Phi} \mathcal{L}_r(\varphi, \lambda), \\ \mathcal{L}_r(\varphi, \lambda) := J_r(\varphi) + \langle L^*\varphi, \lambda \rangle_{L^2(Q_T)} \\ \Phi := \left\{ \varphi : \varphi \in C^0(0, T; H_0^1(\Omega)) \cap C^1(0, T; L^2(\Omega)), L^*\varphi \in L^2(Q_T) \right\} \supset \mathbf{W} \end{array} \right. \quad (18)$$

Remark- For all $\eta > 0$, Φ is endowed with the scalar product,

$$\langle \varphi, \bar{\varphi} \rangle_{\Phi} := \left\langle \frac{\partial \varphi}{\partial \nu}, \frac{\partial \bar{\varphi}}{\partial \nu} \right\rangle_{L^2(\Gamma_T)} + \eta \langle L^*\varphi, L^*\bar{\varphi} \rangle_{L^2(Q_T)}, \quad \forall \varphi, \bar{\varphi} \in \Phi.$$

$\|\varphi\|_{\Phi} := \sqrt{\langle \varphi, \varphi \rangle_{\Phi}}$ is a norm and $(\Phi, \|\cdot\|_{\Phi})$ is an Hilbert space.

³N. Cindea, AM, A mixed formulation for the direct approximation of the control of minimal L^2 -norm for linear type wave equations, (2015)

Relaxation of $L^*\varphi = 0$ via a Lagrange Multiplier

In order to address the $L^2(Q_T)$ constraint $L^*\varphi = 0$, we introduce a Lagrange multiplier $\lambda \in L^2(Q_T)$; we consider the saddle point problem³:

$$\left\{ \begin{array}{l} \sup_{\lambda \in L^2(Q_T)} \inf_{\varphi \in \Phi} \mathcal{L}_r(\varphi, \lambda), \\ \mathcal{L}_r(\varphi, \lambda) := J_r(\varphi) + \langle L^*\varphi, \lambda \rangle_{L^2(Q_T)} \\ \Phi := \left\{ \varphi : \varphi \in C^0(0, T; H_0^1(\Omega)) \cap C^1(0, T; L^2(\Omega)), L^*\varphi \in L^2(Q_T) \right\} \supset \mathbf{W} \end{array} \right. \quad (18)$$

Remark- For all $\eta > 0$, Φ is endowed with the scalar product,

$$\langle \varphi, \bar{\varphi} \rangle_{\Phi} := \left\langle \frac{\partial \varphi}{\partial \nu}, \frac{\partial \bar{\varphi}}{\partial \nu} \right\rangle_{L^2(\Gamma_T)} + \eta \langle L^*\varphi, L^*\bar{\varphi} \rangle_{L^2(Q_T)}, \quad \forall \varphi, \bar{\varphi} \in \Phi.$$

$\|\varphi\|_{\Phi} := \sqrt{\langle \varphi, \varphi \rangle_{\Phi}}$ is a norm and $(\Phi, \|\cdot\|_{\Phi})$ is an Hilbert space.

³N. Cindea, AM, A mixed formulation for the direct approximation of the control of minimal L^2 -norm for linear type wave equations, (2015)

Mixed formulation

Find $(\varphi, \lambda) \in \Phi \times L^2(Q_T)$ solution of

$$\begin{cases} a_r(\varphi, \bar{\varphi}) + b(\bar{\varphi}, \lambda) = l(\bar{\varphi}), & \forall \bar{\varphi} \in \Phi \\ b(\varphi, \bar{\lambda}) = 0, & \forall \bar{\lambda} \in L^2(Q_T), \end{cases} \quad (19)$$

where

$$\begin{aligned} a_r : \Phi \times \Phi &\rightarrow \mathbb{R}, & a_r(\varphi, \bar{\varphi}) &= \left\langle \frac{\partial \varphi}{\partial \nu}, \frac{\partial \bar{\varphi}}{\partial \nu} \right\rangle_{L^2(\Gamma_T)} + r \langle L^* \varphi, L^* \bar{\varphi} \rangle_{L^2(Q_T)} \\ b : \Phi \times L^2(Q_T) &\rightarrow \mathbb{R}, & b(\varphi, \lambda) &= \langle L^* \varphi, \lambda \rangle_{L^2(Q_T)} \\ l : \Phi &\rightarrow \mathbb{R}, & l(\varphi) &= - \langle y_0, \varphi_t(\cdot, 0) \rangle_{L^2} + \langle y_1, \varphi(\cdot, 0) \rangle_{H^{-1}, H_0^1} \end{aligned}$$

Theorem

For all $r \geq 0$,

1. The mixed formulation is well-posed.
2. The unique solution $(\varphi, \lambda) \in \Phi \times L^2(Q_T)$ is the unique saddle-point of the Lagrangian $\mathcal{L}_r : \Phi \times L^2(Q_T) \rightarrow \mathbb{R}$ defined by

$$\mathcal{L}_r(\varphi, \lambda) = \frac{1}{2} a_r(\varphi, \varphi) + b(\varphi, \lambda) - l(\varphi). \quad (20)$$

3. The optimal function φ given by 2. satisfies $\varphi \in \mathbf{W}$ and is the minimizer of J_r^* over \mathbf{W} while the optimal function $\lambda \in L^2(Q_T)$ is the state of the controlled wave equation in the weak sense.
4. We have the following estimates

$$\|\varphi\|_{\Phi} \leq \|y_0, y_1\|_{\mathbf{H}},$$

$$\|\lambda\|_{L^2} \leq \frac{1}{\delta} \left(1 + \max\left(1, \frac{r}{\eta}\right) \right) \|y_0, y_1\|_{\mathbf{H}}, \quad \delta = (C_{\Omega, T} + \eta)^{-1/2}$$

Well-posedness 2

The kernel $\mathcal{N}(b) = \{\varphi \in \Phi; b(\varphi, \lambda) = 0 \quad \forall \lambda \in L^2(Q_T)\}$ coincides with \mathbf{W} : we get

$$a_r(\varphi, \varphi) = \|\varphi\|_{\Phi}^2, \quad \forall \varphi \in \mathcal{N}(b) = \mathbf{W}.$$

It remains to check the **inf-sup constant property** : $\exists \delta > 0$ such that

$$\inf_{\lambda \in L^2} \sup_{\varphi \in \Phi} \frac{b(\varphi, \lambda)}{\|\varphi\|_{\Phi} \|\lambda\|_{L^2}} \geq \delta. \quad (21)$$

For any fixed $\lambda \in L^2(Q_T)$, we define $\varphi^0 \in \Phi$ as the unique solution of

$$L^* \varphi^0 = \lambda \text{ in } Q_T, \quad (\varphi^0(\cdot, 0), \varphi_t^0(\cdot, 0)) = (0, 0) \text{ on } \Omega, \quad \varphi^0 = 0 \text{ on } \Sigma_T.$$

We get $b(\varphi^0, \lambda) = \|\lambda\|_{L^2}^2$ and $\|\varphi^0\|_{\Phi}^2 = \left\| \frac{\partial \varphi^0}{\partial \nu} \right\|_{L^2(\Gamma_T)}^2 + \eta \|\lambda\|_{L^2}^2$.

The estimate $\left\| \frac{\partial \varphi^0}{\partial \nu} \right\|_{L^2(\Gamma_T)} \leq \sqrt{C_{\Omega, T}} \|\lambda\|_{L^2(Q_T)}$ implies that

$$\sup_{\varphi \in \Phi} \frac{b(\varphi, \lambda)}{\|\varphi\|_{\Phi} \|\lambda\|_{L^2}} \geq \frac{b(\varphi^0, \lambda)}{\|\varphi^0\|_{\Phi} \|\lambda\|_{L^2}} \geq \frac{1}{\sqrt{C_{\Omega, T} + \eta}} > 0$$

leading to the inf-sup property with $\delta = (C_{\Omega, T} + \eta)^{-1/2}$.

Well-posedness 2

The kernel $\mathcal{N}(b) = \{\varphi \in \Phi; b(\varphi, \lambda) = 0 \quad \forall \lambda \in L^2(Q_T)\}$ coincides with \mathbf{W} : we get

$$a_r(\varphi, \varphi) = \|\varphi\|_{\Phi}^2, \quad \forall \varphi \in \mathcal{N}(b) = \mathbf{W}.$$

It remains to check the **inf-sup constant property** : $\exists \delta > 0$ such that

$$\inf_{\lambda \in L^2} \sup_{\varphi \in \Phi} \frac{b(\varphi, \lambda)}{\|\varphi\|_{\Phi} \|\lambda\|_{L^2}} \geq \delta. \quad (21)$$

For any fixed $\lambda \in L^2(Q_T)$, we define $\varphi^0 \in \Phi$ as the unique solution of

$$L^* \varphi^0 = \lambda \text{ in } Q_T, \quad (\varphi^0(\cdot, 0), \varphi_t^0(\cdot, 0)) = (0, 0) \text{ on } \Omega, \quad \varphi^0 = 0 \text{ on } \Sigma_T.$$

We get $b(\varphi^0, \lambda) = \|\lambda\|_{L^2}^2$ and $\|\varphi^0\|_{\Phi}^2 = \left\| \frac{\partial \varphi^0}{\partial \nu} \right\|_{L^2(\Gamma_T)}^2 + \eta \|\lambda\|_{L^2}^2$.

The estimate $\left\| \frac{\partial \varphi^0}{\partial \nu} \right\|_{L^2(\Gamma_T)} \leq \sqrt{C_{\Omega, T}} \|\lambda\|_{L^2(Q_T)}$ implies that

$$\sup_{\varphi \in \Phi} \frac{b(\varphi, \lambda)}{\|\varphi\|_{\Phi} \|\lambda\|_{L^2}} \geq \frac{b(\varphi^0, \lambda)}{\|\varphi^0\|_{\Phi} \|\lambda\|_{L^2}} \geq \frac{1}{\sqrt{C_{\Omega, T} + \eta}} > 0$$

leading to the inf-sup property with $\delta = (C_{\Omega, T} + \eta)^{-1/2}$.

Well-posedness 2

The kernel $\mathcal{N}(b) = \{\varphi \in \Phi; b(\varphi, \lambda) = 0 \quad \forall \lambda \in L^2(Q_T)\}$ coincides with \mathbf{W} : we get

$$a_r(\varphi, \varphi) = \|\varphi\|_{\Phi}^2, \quad \forall \varphi \in \mathcal{N}(b) = \mathbf{W}.$$

It remains to check the **inf-sup constant property** : $\exists \delta > 0$ such that

$$\inf_{\lambda \in L^2} \sup_{\varphi \in \Phi} \frac{b(\varphi, \lambda)}{\|\varphi\|_{\Phi} \|\lambda\|_{L^2}} \geq \delta. \quad (21)$$

For any fixed $\lambda \in L^2(Q_T)$, we define $\varphi^0 \in \Phi$ as the unique solution of

$$L^* \varphi^0 = \lambda \text{ in } Q_T, \quad (\varphi^0(\cdot, 0), \varphi_t^0(\cdot, 0)) = (0, 0) \text{ on } \Omega, \quad \varphi^0 = 0 \text{ on } \Sigma_T.$$

We get $b(\varphi^0, \lambda) = \|\lambda\|_{L^2}^2$ and $\|\varphi^0\|_{\Phi}^2 = \left\| \frac{\partial \varphi^0}{\partial \nu} \right\|_{L^2(\Gamma_T)}^2 + \eta \|\lambda\|_{L^2}^2$.

The estimate $\left\| \frac{\partial \varphi^0}{\partial \nu} \right\|_{L^2(\Gamma_T)} \leq \sqrt{C_{\Omega, T}} \|\lambda\|_{L^2(Q_T)}$ implies that

$$\sup_{\varphi \in \Phi} \frac{b(\varphi, \lambda)}{\|\varphi\|_{\Phi} \|\lambda\|_{L^2}} \geq \frac{b(\varphi^0, \lambda)}{\|\varphi^0\|_{\Phi} \|\lambda\|_{L^2}} \geq \frac{1}{\sqrt{C_{\Omega, T} + \eta}} > 0$$

leading to the inf-sup property with $\delta = (C_{\Omega, T} + \eta)^{-1/2}$.

Well-posedness 2

The kernel $\mathcal{N}(b) = \{\varphi \in \Phi; b(\varphi, \lambda) = 0 \quad \forall \lambda \in L^2(Q_T)\}$ coincides with \mathbf{W} : we get

$$a_r(\varphi, \varphi) = \|\varphi\|_{\Phi}^2, \quad \forall \varphi \in \mathcal{N}(b) = \mathbf{W}.$$

It remains to check the **inf-sup constant property** : $\exists \delta > 0$ such that

$$\inf_{\lambda \in L^2} \sup_{\varphi \in \Phi} \frac{b(\varphi, \lambda)}{\|\varphi\|_{\Phi} \|\lambda\|_{L^2}} \geq \delta. \quad (21)$$

For any fixed $\lambda \in L^2(Q_T)$, we define $\varphi^0 \in \Phi$ as the unique solution of

$$L^* \varphi^0 = \lambda \text{ in } Q_T, \quad (\varphi^0(\cdot, 0), \varphi_t^0(\cdot, 0)) = (0, 0) \text{ on } \Omega, \quad \varphi^0 = 0 \text{ on } \Sigma_T.$$

We get $b(\varphi^0, \lambda) = \|\lambda\|_{L^2}^2$ and $\|\varphi^0\|_{\Phi}^2 = \left\| \frac{\partial \varphi^0}{\partial \nu} \right\|_{L^2(\Gamma_T)}^2 + \eta \|\lambda\|_{L^2}^2$.

The estimate $\left\| \frac{\partial \varphi^0}{\partial \nu} \right\|_{L^2(\Gamma_T)} \leq \sqrt{C_{\Omega, T}} \|\lambda\|_{L^2(Q_T)}$ implies that

$$\sup_{\varphi \in \Phi} \frac{b(\varphi, \lambda)}{\|\varphi\|_{\Phi} \|\lambda\|_{L^2}} \geq \frac{b(\varphi^0, \lambda)}{\|\varphi^0\|_{\Phi} \|\lambda\|_{L^2}} \geq \frac{1}{\sqrt{C_{\Omega, T} + \eta}} > 0$$

leading to the inf-sup property with $\delta = (C_{\Omega, T} + \eta)^{-1/2}$.

The multiplier λ

Taking $r = 0$, the first equation reads

$$a_{r=0}(\varphi, \bar{\varphi}) + b(\bar{\varphi}, \lambda) = l(\bar{\varphi}), \quad \forall \bar{\varphi} \in \Phi \quad (22)$$

i.e.

$$\iint_{\Gamma_T} \frac{\partial \varphi}{\partial \nu} \frac{\partial \bar{\varphi}}{\partial \nu} + \iint_{Q_T} \lambda L^* \bar{\varphi} = - \langle y_0, \bar{\varphi}_t(\cdot, 0) \rangle_{L^2} + \langle y_1, \bar{\varphi}(\cdot, 0) \rangle_{H^{-1}, H_0^1}, \quad \forall \bar{\varphi} \in \Phi \quad (23)$$

which means $\lambda \in L^2(Q_T)$ is solution in the sense of transposition of

$$\begin{cases} L\lambda = 0, & \text{in } Q_T \\ (\lambda(\cdot, 0), \lambda_t(\cdot, 0)) = (y_0, y_1) \in L^2(\Omega) \times H^{-1}(\Omega) \\ (\lambda(\cdot, T), \lambda_t(\cdot, T)) = (0, 0), \\ \lambda = \frac{\partial \varphi}{\partial \nu} & \text{on } \Gamma_T \end{cases} \quad (24)$$

Therefore, λ coincides with the weak solution of the wave equation controlled by v .

$$\lambda \in C^0([0, T], L^2(\Omega)) \cap C^1([0, T], H^{-1}(\Omega))$$

The multiplier λ

Taking $r = 0$, the first equation reads

$$a_{r=0}(\varphi, \bar{\varphi}) + b(\bar{\varphi}, \lambda) = l(\bar{\varphi}), \quad \forall \bar{\varphi} \in \Phi \quad (22)$$

i.e.

$$\iint_{\Gamma_T} \frac{\partial \varphi}{\partial \nu} \frac{\partial \bar{\varphi}}{\partial \nu} + \iint_{Q_T} \lambda L^* \bar{\varphi} = - \langle y_0, \bar{\varphi}_t(\cdot, 0) \rangle_{L^2} + \langle y_1, \bar{\varphi}(\cdot, 0) \rangle_{H^{-1}, H_0^1}, \quad \forall \bar{\varphi} \in \Phi \quad (23)$$

which means $\lambda \in L^2(Q_T)$ is solution in the sense of transposition of

$$\begin{cases} L\lambda = 0, & \text{in } Q_T \\ (\lambda(\cdot, 0), \lambda_t(\cdot, 0)) = (y_0, y_1) \in L^2(\Omega) \times H^{-1}(\Omega) \\ (\lambda(\cdot, T), \lambda_t(\cdot, T)) = (0, 0), \\ \lambda = \frac{\partial \varphi}{\partial \nu} & \text{on } \Gamma_T \end{cases} \quad (24)$$

Therefore, λ coincides with the weak solution of the wave equation controlled by v .

$$\lambda \in C^0([0, T], L^2(\Omega)) \cap C^1([0, T], H^{-1}(\Omega))$$

"Dual of the dual" - Equivalent minimization w.r.t. λ

Lemma

Let \mathcal{P}_r be the linear operator from L^2 into L^2 defined by

$$\mathcal{P}_r \lambda := L^* \varphi, \quad \forall \lambda \in L^2 \quad \text{where } \varphi \in \Phi \text{ solves } a_r(\varphi, \bar{\varphi}) = b(\bar{\varphi}, \lambda), \quad \forall \bar{\varphi} \in \Phi.$$

For any $r > 0$, the operator \mathcal{P}_r is a **strongly elliptic, symmetric** isomorphism from L^2 into L^2 .

Theorem

$$\sup_{\lambda \in L^2} \inf_{\varphi \in \Phi} \mathcal{L}_r(\varphi, \lambda) = - \inf_{\lambda \in L^2} J_r^*(\lambda) + \mathcal{L}_r(\varphi_0, 0)$$

where $\varphi_0 \in \Phi$ solves $a_r(\varphi_0, \bar{\varphi}) = l(\bar{\varphi}), \forall \bar{\varphi} \in \Phi$ and $J_r^* : L^2 \rightarrow \mathbb{R}$ defined by

$$J_r^*(\lambda) := \frac{1}{2} \langle \mathcal{P}_r \lambda, \lambda \rangle_{L^2(\Omega_T)} - b(\varphi_0, \lambda)$$

"Dual of the dual" - Equivalent minimization w.r.t. λ

Lemma

Let \mathcal{P}_r be the linear operator from L^2 into L^2 defined by

$$\mathcal{P}_r \lambda := L^* \varphi, \quad \forall \lambda \in L^2 \quad \text{where } \varphi \in \Phi \text{ solves } a_r(\varphi, \bar{\varphi}) = b(\bar{\varphi}, \lambda), \quad \forall \bar{\varphi} \in \Phi.$$

For any $r > 0$, the operator \mathcal{P}_r is a **strongly elliptic, symmetric** isomorphism from L^2 into L^2 .

Theorem

$$\sup_{\lambda \in L^2} \inf_{\varphi \in \Phi} \mathcal{L}_r(\varphi, \lambda) = - \inf_{\lambda \in L^2} J_r^*(\lambda) + \mathcal{L}_r(\varphi_0, 0)$$

where $\varphi_0 \in \Phi$ solves $a_r(\varphi_0, \bar{\varphi}) = l(\bar{\varphi}), \forall \bar{\varphi} \in \Phi$ and $J_r^* : L^2 \rightarrow \mathbb{R}$ defined by

$$J_r^*(\lambda) := \frac{1}{2} \langle \mathcal{P}_r \lambda, \lambda \rangle_{L^2(\Omega_r)} - b(\varphi_0, \lambda)$$

"Dual of the dual" - Equivalent minimization w.r.t. λ

Lemma

Let \mathcal{P}_r be the linear operator from L^2 into L^2 defined by

$$\mathcal{P}_r \lambda := L^* \varphi, \quad \forall \lambda \in L^2 \quad \text{where } \varphi \in \Phi \text{ solves } a_r(\varphi, \bar{\varphi}) = b(\bar{\varphi}, \lambda), \quad \forall \bar{\varphi} \in \Phi.$$

For any $r > 0$, the operator \mathcal{P}_r is a **strongly elliptic, symmetric** isomorphism from L^2 into L^2 .

Theorem

$$\sup_{\lambda \in L^2} \inf_{\varphi \in \Phi} \mathcal{L}_r(\varphi, \lambda) = - \inf_{\lambda \in L^2} J_r^{**}(\lambda) + \mathcal{L}_r(\varphi_0, 0)$$

where $\varphi_0 \in \Phi$ solves $a_r(\varphi_0, \bar{\varphi}) = l(\bar{\varphi}), \forall \bar{\varphi} \in \Phi$ and $J_r^{**} : L^2 \rightarrow \mathbb{R}$ defined by

$$J_r^{**}(\lambda) := \frac{1}{2} \langle \mathcal{P}_r \lambda, \lambda \rangle_{L^2(Q_T)} - b(\varphi_0, \lambda)$$

Conformal Approximation of the mixed formulation

Let then Φ_h and Λ_h be two finite dimensional spaces parametrized by the variable h such that

$$\Phi_h \subset \Phi, \quad \Lambda_h \subset L^2(Q_T), \quad \forall h > 0.$$

Then, we can introduce the following approximated problems : find $(\varphi_h, \lambda_h) \in \Phi_h \times \Lambda_h$ solution of

$$\begin{cases} a_r(\varphi_h, \bar{\varphi}_h) + b(\bar{\varphi}_h, \lambda_h) & = I(\bar{\varphi}_h), & \forall \bar{\varphi}_h \in \Phi_h \\ b(\varphi_h, \bar{\lambda}_h) & = 0, & \forall \bar{\lambda}_h \in \Lambda_h. \end{cases} \quad (25)$$

For any $h > 0$, the well-posedness is again a consequence of two properties

- ▶ the coercivity of the bilinear form a_r on the subset $\mathcal{N}_h(b) = \{\varphi_h \in \Phi_h; b(\varphi_h, \lambda_h) = 0 \quad \forall \lambda_h \in \Lambda_h\}$. From

$$a_r(\varphi, \varphi) \geq \frac{r}{\eta} \|\varphi\|_{\Phi}^2, \quad \forall \varphi \in \Phi \quad (r > 0!!!)$$

the form a_r is coercive on the full space Φ , and so *a fortiori* on $\mathcal{N}_h(b) \subset \Phi_h \subset \Phi$.

- ▶ The second property is a discrete inf-sup condition : there exists $\delta > 0$ such that

$$\delta_h := \inf_{\lambda_h \in \Lambda_h} \sup_{\varphi_h \in \Phi_h} \frac{b(\varphi_h, \lambda_h)}{\|\varphi_h\|_{\Phi_h} \|\lambda_h\|_{\Lambda_h}} \geq \delta. \quad (26)$$

Necessary condition: $\dim(\Phi_h) > \dim(\Lambda_h)$

Conformal Approximation of the mixed formulation

Let then Φ_h and Λ_h be two finite dimensional spaces parametrized by the variable h such that

$$\Phi_h \subset \Phi, \quad \Lambda_h \subset L^2(Q_T), \quad \forall h > 0.$$

Then, we can introduce the following approximated problems : find $(\varphi_h, \lambda_h) \in \Phi_h \times \Lambda_h$ solution of

$$\begin{cases} a_r(\varphi_h, \bar{\varphi}_h) + b(\bar{\varphi}_h, \lambda_h) &= I(\bar{\varphi}_h), & \forall \bar{\varphi}_h \in \Phi_h \\ b(\varphi_h, \bar{\lambda}_h) &= 0, & \forall \bar{\lambda}_h \in \Lambda_h. \end{cases} \quad (25)$$

For any $h > 0$, the well-posedness is again a consequence of two properties

- ▶ the coercivity of the bilinear form a_r on the subset $\mathcal{N}_h(b) = \{\varphi_h \in \Phi_h; b(\varphi_h, \lambda_h) = 0 \quad \forall \lambda_h \in \Lambda_h\}$. From

$$a_r(\varphi, \varphi) \geq \frac{r}{\eta} \|\varphi\|_{\Phi}^2, \quad \forall \varphi \in \Phi \quad (r > 0!!)$$

the form a_r is coercive on the full space Φ , and so *a fortiori* on $\mathcal{N}_h(b) \subset \Phi_h \subset \Phi$.

- ▶ The second property is a discrete inf-sup condition : there exists $\delta > 0$ such that

$$\delta_h := \inf_{\lambda_h \in \Lambda_h} \sup_{\varphi_h \in \Phi_h} \frac{b(\varphi_h, \lambda_h)}{\|\varphi_h\|_{\Phi_h} \|\lambda_h\|_{\Lambda_h}} \geq \delta. \quad (26)$$

Necessary condition: $\dim(\Phi_h) > \dim(\Lambda_h)$

Conformal Approximation of the mixed formulation

Let then Φ_h and Λ_h be two finite dimensional spaces parametrized by the variable h such that

$$\Phi_h \subset \Phi, \quad \Lambda_h \subset L^2(Q_T), \quad \forall h > 0.$$

Then, we can introduce the following approximated problems : find $(\varphi_h, \lambda_h) \in \Phi_h \times \Lambda_h$ solution of

$$\begin{cases} a_r(\varphi_h, \bar{\varphi}_h) + b(\bar{\varphi}_h, \lambda_h) &= I(\bar{\varphi}_h), & \forall \bar{\varphi}_h \in \Phi_h \\ b(\varphi_h, \bar{\lambda}_h) &= 0, & \forall \bar{\lambda}_h \in \Lambda_h. \end{cases} \quad (25)$$

For any $h > 0$, the well-posedness is again a consequence of two properties

- ▶ the coercivity of the bilinear form a_r on the subset $\mathcal{N}_h(b) = \{\varphi_h \in \Phi_h; b(\varphi_h, \lambda_h) = 0 \quad \forall \lambda_h \in \Lambda_h\}$. From

$$a_r(\varphi, \varphi) \geq \frac{r}{\eta} \|\varphi\|_{\Phi}^2, \quad \forall \varphi \in \Phi \quad (r > 0!!)$$

the form a_r is coercive on the full space Φ , and so *a fortiori* on $\mathcal{N}_h(b) \subset \Phi_h \subset \Phi$.

- ▶ The second property is a discrete inf-sup condition : there exists $\delta > 0$ such that

$$\delta_h := \inf_{\lambda_h \in \Lambda_h} \sup_{\varphi_h \in \Phi_h} \frac{b(\varphi_h, \lambda_h)}{\|\varphi_h\|_{\Phi_h} \|\lambda_h\|_{\Lambda_h}} \geq \delta. \quad (26)$$

Necessary condition: $\dim(\Phi_h) > \dim(\Lambda_h)$

Finite dimensional linear system

Let $n_h = \dim \Phi_h$, $m_h = \dim \Lambda_h$ and let the real matrices $A_{r,h} \in \mathbb{R}^{n_h, n_h}$, $B_h \in \mathbb{R}^{m_h, n_h}$, $J_h \in \mathbb{R}^{m_h, m_h}$ and $L_h \in \mathbb{R}^{n_h}$ be defined by

$$\begin{cases} a_r(\varphi_h, \overline{\varphi_h}) = \langle A_{r,h}\{\varphi_h\}, \{\overline{\varphi_h}\} \rangle_{\mathbb{R}^{n_h}, \mathbb{R}^{n_h}}, & \forall \varphi_h, \overline{\varphi_h} \in \Phi_h, \\ b(\varphi_h, \lambda_h) = \langle B_h\{\varphi_h\}, \{\lambda_h\} \rangle_{\mathbb{R}^{m_h}, \mathbb{R}^{m_h}}, & \forall \varphi_h \in \Phi_h, \forall \lambda_h \in \Lambda_h, \\ l(\varphi_h) = \langle L_h, \{\varphi_h\} \rangle, & \forall \varphi_h \in \Phi_h \end{cases}$$

where $\{\varphi_h\} \in \mathbb{R}^{n_h}$ denotes the vector associated to φ_h and $\langle \cdot, \cdot \rangle_{\mathbb{R}^{n_h}, \mathbb{R}^{n_h}}$ the usual scalar product over \mathbb{R}^{n_h} . Problem (25) reads as follows :

find $\{\varphi_h\} \in \mathbb{R}^{n_h}$ and $\{\lambda_h\} \in \mathbb{R}^{m_h}$ such that

$$\begin{pmatrix} A_{r,h} & B_h^T \\ B_h & 0 \end{pmatrix}_{\mathbb{R}^{n_h+m_h}, \mathbb{R}^{n_h+m_h}} \begin{pmatrix} \{\varphi_h\} \\ \{\lambda_h\} \end{pmatrix}_{\mathbb{R}^{n_h+m_h}} = \begin{pmatrix} L_h \\ 0 \end{pmatrix}_{\mathbb{R}^{n_h+m_h}}. \quad (27)$$

$A_{r,h}$ is symmetric and positive definite for any $h > 0$ and any $r > 0$.

The full matrix of order $m_h + n_h$ in (27) is symmetric but not positive definite.

Choice of the conformal spaces Φ_h and Λ_h

We introduce a triangulation \mathcal{T}_h such that $\overline{Q_T} = \cup_{K \in \mathcal{T}_h} K$ and we assume that $\{\mathcal{T}_h\}_{h>0}$ is a regular family. We note $h := \max\{\text{diam}(K), K \in \mathcal{T}_h\}$.

- We define the finite dimensional space

$$\Lambda_h = \{\lambda_h \in C^0(\overline{Q_T}), \lambda_h|_K \in \mathbb{P}_1(K) \quad \forall K \in \mathcal{T}_h, \lambda_h = 0 \text{ on } \Sigma_T\} \subset L^2(Q_T)$$

- The space Φ_h must be chosen such that $L^* \varphi_h \in L^2(Q_T)$ for any $\varphi_h \in \Phi_h$. We introduce the space Φ_h as follows:

$$\Phi_h = \{\varphi_h \in \Phi_h \in C^1(\overline{Q_T}) : \varphi_h|_K \in \mathbb{P}(K) \quad \forall K \in \mathcal{T}_h, \varphi_h = 0 \text{ on } \Sigma_T\} \subset \Phi$$

where $\mathbb{P}(K)$ denotes an appropriate space of polynomial functions in x and t .

Choice of the conformal spaces Φ_h and Λ_h

We introduce a triangulation \mathcal{T}_h such that $\overline{Q_T} = \cup_{K \in \mathcal{T}_h} K$ and we assume that $\{\mathcal{T}_h\}_{h>0}$ is a regular family. We note $h := \max\{\text{diam}(K), K \in \mathcal{T}_h\}$.

- We define the finite dimensional space

$$\Lambda_h = \{\lambda_h \in C^0(\overline{Q_T}), \lambda_h|_K \in \mathbb{P}_1(K) \quad \forall K \in \mathcal{T}_h, \lambda_h = 0 \text{ on } \Sigma_T\} \subset L^2(Q_T)$$

- The space Φ_h must be chosen such that $L^* \varphi_h \in L^2(Q_T)$ for any $\varphi_h \in \Phi_h$. We introduce the space Φ_h as follows:

$$\Phi_h = \{\varphi_h \in \Phi_h \in C^1(\overline{Q_T}) : \varphi_h|_K \in \mathbb{P}(K) \quad \forall K \in \mathcal{T}_h, \varphi_h = 0 \text{ on } \Sigma_T\} \subset \Phi$$

where $\mathbb{P}(K)$ denotes an appropriate space of polynomial functions in x and t .

Choice of the conformal spaces Φ_h and Λ_h

We introduce a triangulation \mathcal{T}_h such that $\overline{Q_T} = \cup_{K \in \mathcal{T}_h} K$ and we assume that $\{\mathcal{T}_h\}_{h>0}$ is a regular family. We note $h := \max\{\text{diam}(K), K \in \mathcal{T}_h\}$.

- We define the finite dimensional space

$$\Lambda_h = \{\lambda_h \in C^0(\overline{Q_T}), \lambda_h|_K \in \mathbb{P}_1(K) \quad \forall K \in \mathcal{T}_h, \lambda_h = 0 \text{ on } \Sigma_T\} \subset L^2(Q_T)$$

- The space Φ_h must be chosen such that $L^* \varphi_h \in L^2(Q_T)$ for any $\varphi_h \in \Phi_h$. We introduce the space Φ_h as follows:

$$\Phi_h = \{\varphi_h \in \Phi_h \in C^1(\overline{Q_T}) : \varphi_h|_K \in \mathbb{P}(K) \quad \forall K \in \mathcal{T}_h, \varphi_h = 0 \text{ on } \Sigma_T\} \subset \Phi$$

where $\mathbb{P}(K)$ denotes an appropriate space of polynomial functions in x and t .

C^1 finite element over Q_T

4

$$\Phi_h = \{\varphi_h \in \Phi_h \in C^1(\overline{Q_T}) : \varphi_h|_K \in \mathbb{P}(K) \quad \forall K \in \mathcal{T}_h, \varphi_h = 0 \text{ on } \Sigma_T\}$$

where $\mathbb{P}(K)$ denotes an appropriate space of polynomial functions in x and t .

We may consider the following choices for $\mathbb{P}(K)$:

1. The *Bogner-Fox-Schmit* (BFS for short) C^1 element defined for rectangles. It involves the values of $\varphi_h, \varphi_{h,x}, \varphi_{h,t}, \varphi_{h,xt}$ on the four vertices of each rectangle K .
2. The *reduced Hsieh-Clough-Tocher* (HCT for short) C^1 element defined for triangles. This is a so-called composite finite element and the values of $\varphi_h, \varphi_{h,x}, \varphi_{h,t}$ on the three vertices of each triangle K .

C^1 finite element over Q_T

4

$$\Phi_h = \{\varphi_h \in \Phi_h \in C^1(\overline{Q_T}) : \varphi_h|_K \in \mathbb{P}(K) \quad \forall K \in \mathcal{T}_h, \varphi_h = 0 \text{ on } \Sigma_T\}$$

where $\mathbb{P}(K)$ denotes an appropriate space of polynomial functions in x and t .

We may consider the following choices for $\mathbb{P}(K)$:

1. The *Bogner-Fox-Schmit* (BFS for short) C^1 element defined for **rectangles**. It involves the values of $\varphi_h, \varphi_{h,x}, \varphi_{h,t}, \varphi_{h,xt}$ on the four vertices of each rectangle K .
2. The *reduced Hsieh-Clough-Tocher* (HCT for short) C^1 element defined for **triangles**. This is a so-called composite finite element and the values of $\varphi_h, \varphi_{h,x}, \varphi_{h,t}$ on the three vertices of each triangle K .

C^1 finite element over Q_T

4

$$\Phi_h = \{\varphi_h \in \Phi_h \in C^1(\overline{Q_T}) : \varphi_h|_K \in \mathbb{P}(K) \quad \forall K \in \mathcal{T}_h, \varphi_h = 0 \text{ on } \Sigma_T\}$$

where $\mathbb{P}(K)$ denotes an appropriate space of polynomial functions in x and t .

We may consider the following choices for $\mathbb{P}(K)$:

1. The *Bogner-Fox-Schmit* (BFS for short) C^1 element defined for **rectangles**. It involves the values of $\varphi_h, \varphi_{h,x}, \varphi_{h,t}, \varphi_{h,xt}$ on the four vertices of each rectangle K .
2. The *reduced Hsieh-Clough-Tocher* (HCT for short) C^1 element defined for **triangles**. This is a so-called composite finite element and the values of $\varphi_h, \varphi_{h,x}, \varphi_{h,t}$ on the three vertices of each triangle K .

Convergence rate in Φ

From [D. Boffi, F. Brezzi, M. Fortin, *Mixed finite element methods and applications*. 2013],

Proposition (BFS element for $N = 1$ - Convergence in $\Phi \times L^2$)

Let $h > 0$, let $k \leq 2$. If $(\varphi, \lambda) \in H^{k+2}(Q_T) \times H^k(Q_T)$, $\exists K > 0$

$$\|\varphi - \varphi_h\|_{\Phi} \leq K \left(1 + \frac{1}{\sqrt{\eta}\delta_h} + \frac{1}{\sqrt{\eta}} \right) h^k,$$

$$\|\lambda - \lambda_h\|_{L^2(Q_T)} \leq K \left(\left(1 + \frac{1}{\sqrt{\eta}\delta_h} \right) \frac{1}{\delta_h} + \frac{1}{\sqrt{\eta}\delta_h} \right) h^k.$$

Corollary (Estimates on the approximation of the control)

Under the previous assumptions, the approximation $v_h := \nabla \varphi_h \cdot \nu 1_{\Gamma_0}$ satisfies

$$\|v - v_h\|_{L^2(\Gamma_0 \times (0, T))} \leq K \left(1 + \frac{1}{\sqrt{\eta}\delta_h} + \frac{1}{\sqrt{\eta}} \right) h^k. \quad (28)$$

Convergence rate in Φ

From [D. Boffi, F. Brezzi, M. Fortin, Mixed finite element methods and applications. 2013],

Proposition (BFS element for $N = 1$ - Convergence in $\Phi \times L^2$)

Let $h > 0$, let $k \leq 2$. If $(\varphi, \lambda) \in H^{k+2}(Q_T) \times H^k(Q_T)$, $\exists K > 0$

$$\|\varphi - \varphi_h\|_{\Phi} \leq K \left(1 + \frac{1}{\sqrt{\eta}\delta_h} + \frac{1}{\sqrt{\eta}} \right) h^k,$$

$$\|\lambda - \lambda_h\|_{L^2(Q_T)} \leq K \left(\left(1 + \frac{1}{\sqrt{\eta}\delta_h} \right) \frac{1}{\delta_h} + \frac{1}{\sqrt{\eta}\delta_h} \right) h^k.$$

Corollary (Estimates on the approximation of the control)

Under the previous assumptions, the approximation $\mathbf{v}_h := \nabla\varphi_h \cdot \boldsymbol{\nu} \mathbf{1}_{\Gamma_0}$ satisfies

$$\|\mathbf{v} - \mathbf{v}_h\|_{L^2(\Gamma_0 \times (0, T))} \leq K \left(1 + \frac{1}{\sqrt{\eta}\delta_h} + \frac{1}{\sqrt{\eta}} \right) h^k. \quad (28)$$

$N = 1$ - Numerical experiments

$$\Omega = (0, 1) - \Gamma_0 = \{1\} - T = 2.4$$

$$\text{(EX)} \quad y_0(x) = 4x \mathbf{1}_{(0,1/2)}(x), \quad y_1(x) = 0, \quad x \in \Omega$$

$$v(t) = 2(1-t) \mathbf{1}_{(1/2,3/2)}(t), \quad t \in (0, T), \quad \|v\|_{L^2(0,T)} = 1/\sqrt{3} \approx 0.5773. \quad (29)$$

$N = 1$ - Numerical experiments




Figure: Control of minimal L^2 -norm v and its approximation v_h on $(0, T)$; $r = 10^{-2}$; $h = 2.46 \times 10^{-2}$

Example 1 - $N = 1$ - Numerical experiments

h	1.41×10^{-1}	7.01×10^{-2}	3.53×10^{-2}	1.76×10^{-2}	8.83×10^{-3}
$\ v_h\ _{L^2(0,T)}$	0.6003	0.5850	0.5776	0.5752	0.5747
$\ v - v_h\ _{L^2(0,T)}$	2.87×10^{-1}	2.05×10^{-1}	1.47×10^{-1}	1.08×10^{-1}	8.18×10^{-2}
$\ \lambda_h\ _{L^2(Q_T)}$	0.62	0.598	0.586	0.581	0.578
$\ L^* \varphi_h\ _{L^2(Q_T)}$	1.02×10^{-1}	7.53×10^{-2}	5.8×10^{-2}	4.55×10^{-2}	3.6×10^{-2}
$\ L^* \varphi_h\ _{H^{-1}(Q_T)}$	1.92×10^{-16}	3.83×10^{-16}	7.46×10^{-16}	1.51×10^{-15}	2.81×10^{-15}

Table: BFS element - $r = 1$.

$$r = 1 : \quad \|v - v_h\|_{L^2(0,T)} \approx 1.12 \cdot h^{0.52}, \quad \|L^* \varphi_h\|_{L^2(Q_T)} \approx 15.67 \cdot h^{0.72},$$

$$r = 10^{-2} : \quad \|v - v_h\|_{L^2(0,T)} \approx 0.83 \cdot h^{0.45}, \quad \|L^* \varphi_h\|_{L^2(Q_T)} \approx 0.24 \cdot h^{0.37}.$$

A curiosity : $\|v_h\|_{L^2(0,T)}$ is close to $\|y_h\|_{L^2(Q_T)}$!?!!

Example 1 - $N = 1$ - Numerical experiments




Figure: The dual variable φ_h in Q_T ; $h = 2.46 \times 10^{-2}$; $r = 10^{-2}$.

Example 1 - $N = 1$ - Numerical experiments




Figure: The primal variable λ_h in Q_T ; $h = 2.46 \times 10^{-2}$; $r = 10^{-2}$.

Mesh adaptation




Figure: Iterative refinement of the triangular mesh over Q_T with respect to the variable λ_h : 142, 412, 1 154, 2 556 ; $r = 2 \times 10^{-3}$.

Example 1 - $N = 1$ - Numerical experiments




Figure: The dual variable φ_h in Q_T corresponding to the finer mesh; $r = 2 \times 10^{-3}$.

Example 1 - $N = 1$ - Numerical experiments




Figure: The primal variable λ_h in Q_T corresponding to the finer mesh.

Minimization of J_r^{**} with respect to λ

$$J_r^{**}(\lambda) := \frac{1}{2} \langle \mathcal{P}_r \lambda, \lambda \rangle_{L^2(Q_T)} - b(\varphi_0, \lambda)$$




Figure: Relative residus $\|g^n\|_{L^2(Q_T)} / \|g^0\|_{L^2(Q_T)}$ w.r.t. the iterate n for $r = 10^2$ (\star), $r = 1$ (\square), $r = 10^{-2}$ (\circ) and $r = h^2$ (\triangleleft); $h = 9.99 \times 10^{-3}$.

Minimization of J_r^{**} with respect to λ

$$J_r^{**}(\lambda) := \frac{1}{2} \langle \mathcal{P}_r \lambda, \lambda \rangle_{L^2(Q_T)} - b(\varphi_0, \lambda)$$

h	1.56×10^{-1}	7.92×10^{-2}	3.99×10^{-2}	1.99×10^{-2}	9.99×10^{-3}
# iterates	20	26	31	44	61
$m_h = \text{card}(\{\lambda_h\})$	231	840	3198	12555	49749
$\ \lambda_h(1, \cdot)\ _{L^2(0, T)}$	0.6089	0.5867	0.5775	0.5746	0.5742
$\ v - \lambda_h(1, \cdot)\ _{L^2(0, T)}$	2.40×10^{-1}	1.68×10^{-1}	1.28×10^{-1}	9.69×10^{-2}	7.62×10^{-2}
$\ \lambda_h\ _{L^2(Q_T)}$	0.6178	0.5963	0.5857	0.5806	0.5784

Table: BFS element - Conjugate gradient algorithm - $r = 1$.

Remind: $\|v\|_{L^2(0, T)} \approx 0.5773$

Comparison with the bi-harmonic regularization [Glowinski'92]

$$\begin{cases} \min_{(\varphi_0, \varphi_1) \in \tilde{V}} J_\epsilon^*(\varphi_0, \varphi_1) := J^*(\varphi_0, \varphi_1) + \frac{\epsilon}{2} \|\varphi_0, \varphi_1\|_{\tilde{V}}^2, & \epsilon > 0, \\ \tilde{V} := H^2(\Omega) \cap H_0^1(\Omega) \times H_0^1(\Omega) \end{cases} \quad (30)$$

Time Marching method here ! : $h = \Delta x$; $\Delta t = 0.8\Delta x$

h	1.56×10^{-1}	7.92×10^{-2}	3.99×10^{-2}	1.99×10^{-2}	9.99×10^{-3}
# iterates	62	> 5000	78	58	39
$\text{card}(\{\varphi_{0h}, \varphi_{1h}\})$	44	84	164	324	644
$\ v_h\ _{L^2(0,T)}$	0.5484	0.5603	0.5671	0.5712	0.5736
$\ v - v_h\ _{L^2(0,T)}$	2.72×10^{-1}	2.23×10^{-1}	1.81×10^{-1}	1.47×10^{-1}	1.24×10^{-1}
$\ y_h\ _{L^2(Q_T)}$	0.5386	0.5557	0.5649	0.5701	0.5731

Table: Bi-harmonic Tychonoff regularization; $\epsilon = h^{1.8}$.

Remark : If ϵ is too small (e.g. $\epsilon = h^2$), the gradient algorithm diverges.

Comparison with the bi-harmonic regularization [Glowinski'92]

$$\begin{cases} \min_{(\varphi_0, \varphi_1) \in \tilde{V}} J_\epsilon^*(\varphi_0, \varphi_1) := J^*(\varphi_0, \varphi_1) + \frac{\epsilon}{2} \|\varphi_0, \varphi_1\|_{\tilde{V}}^2, & \epsilon > 0, \\ \tilde{V} := H^2(\Omega) \cap H_0^1(\Omega) \times H_0^1(\Omega) \end{cases} \quad (30)$$

Time Marching method here ! : $h = \Delta x$; $\Delta t = 0.8\Delta x$

h	1.56×10^{-1}	7.92×10^{-2}	3.99×10^{-2}	1.99×10^{-2}	9.99×10^{-3}
# iterates	62	> 5000	78	58	39
$\text{card}(\{\varphi_{0h}, \varphi_{1h}\})$	44	84	164	324	644
$\ \mathbf{v}_h\ _{L^2(0, T)}$	0.5484	0.5603	0.5671	0.5712	0.5736
$\ \mathbf{v} - \mathbf{v}_h\ _{L^2(0, T)}$	2.72×10^{-1}	2.23×10^{-1}	1.81×10^{-1}	1.47×10^{-1}	1.24×10^{-1}
$\ \mathbf{y}_h\ _{L^2(Q_T)}$	0.5386	0.5557	0.5649	0.5701	0.5731

Table: Bi-harmonic Tychonoff regularization; $\epsilon = h^{1.8}$.


Remark : If ϵ is too small (e.g. $\epsilon = h^2$), the gradient algorithm diverges.

The discrete inf-sup test - A posteriori evaluation of δ_h

$$\delta_h := \inf_{\lambda_h \in \Lambda_h} \sup_{\varphi_h \in \Phi_h} \frac{b(\varphi_h, \lambda_h)}{\|\varphi_h\|_{\Phi_h} \|\lambda_h\|_{\Lambda_h}} \geq \delta. \quad (31)$$

Taking $\eta = r > 0$ so that $a_r(\varphi, \bar{\varphi}) = (\varphi, \bar{\varphi})_{\Phi}$, we have ⁵

$$\delta_h = \inf \left\{ \sqrt{\delta} : B_h A_{r,h}^{-1} B_h^T \{\lambda_h\} = \delta J_h \{\lambda_h\}, \quad \forall \{\lambda_h\} \in \mathbb{R}^{m_h} \setminus \{0\} \right\}. \quad (32)$$



$$\delta_h \approx C_r \frac{h}{\sqrt{r}} \quad \text{as } h \rightarrow 0^+$$

If $r = h^2$, (Φ_h, Λ_h) passes the discrete inf-sup test !

BFS finite element - $h \rightarrow \sqrt{r} \delta_{h,r}$ for $r = 1$ (□),
 $r = 10^{-2}$ (○), $r = h$ (☆) and $r = h^2$ (<)


⁵K. Bathe, D. Chapelle, The discrete inf-sup test, (2003)

The discrete inf-sup test - A posteriori evaluation of δ_h

$$\delta_h := \inf_{\lambda_h \in \Lambda_h} \sup_{\varphi_h \in \Phi_h} \frac{b(\varphi_h, \lambda_h)}{\|\varphi_h\|_{\Phi_h} \|\lambda_h\|_{\Lambda_h}} \geq \delta. \quad (31)$$

Taking $\eta = r > 0$ so that $a_r(\varphi, \bar{\varphi}) = (\varphi, \bar{\varphi})_{\Phi}$, we have ⁵

$$\delta_h = \inf \left\{ \sqrt{\delta} : B_h A_{r,h}^{-1} B_h^T \{\lambda_h\} = \delta J_h \{\lambda_h\}, \quad \forall \{\lambda_h\} \in \mathbb{R}^{m_h} \setminus \{0\} \right\}. \quad (32)$$



$$\delta_h \approx C_r \frac{h}{\sqrt{r}} \quad \text{as } h \rightarrow 0^+$$

If $r = h^2$, (Φ_h, Λ_h) passes the discrete inf-sup test !

BFS finite element - $h \rightarrow \sqrt{r}\delta_{h,r}$ for $r = 1$ (□),
 $r = 10^{-2}$ (○), $r = h$ (★) and $r = h^2$ (<)

⁵K. Bathe, D. Chapelle, The discrete inf-sup test, (2003)

Stabilized mixed formulation "à la Barbosa-Hughes"

6

$\alpha > 0$

$$\begin{cases} \sup_{\lambda \in \Lambda} \inf_{\varphi \in \Phi} \mathcal{L}_{r,\alpha}(\varphi, \lambda), \\ \mathcal{L}_{r,\alpha}(\varphi, \lambda) := \mathcal{L}_r(\varphi, \lambda) - \frac{\alpha}{2} \|L\lambda\|_{L^2(H^{-1}(\Omega))}^2 - \frac{\alpha}{2} \|\lambda - \partial_\nu \varphi\|_{L^2(\Gamma_T)}^2. \end{cases} \quad (33)$$

$$\Lambda := \left\{ \lambda : \lambda \in C([0, T]; L^2(\Omega)) \cap C^1([0, T]; H^{-1}(\Omega)), \right. \\ \left. L\lambda \in L^2([0, T]; H^{-1}(\Omega)), \lambda(\cdot, 0) = \lambda_t(\cdot, 0) = 0, \lambda|_{\Gamma_T} \in L^2(\Gamma_T) \right\}.$$

Λ is a Hilbert space endowed with the following inner product

$$\langle \lambda, \bar{\lambda} \rangle_\Lambda := \int_0^T \langle L\lambda(t), L\bar{\lambda}(t) \rangle_{H^{-1}(\Omega)} dt + \iint_{\Gamma_T} \lambda \bar{\lambda} d\sigma dt, \quad \forall \lambda, \bar{\lambda} \in \Lambda$$

using notably that

$$\|\lambda\|_{L^2(Q_T)} \leq C_{\Omega, T} \sqrt{\langle \lambda, \lambda \rangle_\Lambda}, \quad \forall \lambda \in \Lambda \quad (34)$$

for some positive constant $C_{\Omega, T}$. We denote $\|\lambda\|_\Lambda := \sqrt{\langle \lambda, \lambda \rangle_\Lambda}$.

Stabilized mixed formulation "à la Barbosa-Hughes"

6

$\alpha > 0$

$$\left\{ \begin{array}{l} \sup_{\lambda \in \Lambda} \inf_{\varphi \in \Phi} \mathcal{L}_{r,\alpha}(\varphi, \lambda), \\ \mathcal{L}_{r,\alpha}(\varphi, \lambda) := \mathcal{L}_r(\varphi, \lambda) - \frac{\alpha}{2} \|L\lambda\|_{L^2(H^{-1}(\Omega))}^2 - \frac{\alpha}{2} \|\lambda - \partial_\nu \varphi\|_{L^2(\Gamma_T)}^2. \end{array} \right. \quad (33)$$

$$\Lambda := \left\{ \lambda : \lambda \in C([0, T]; L^2(\Omega)) \cap C^1([0, T]; H^{-1}(\Omega)), \right. \\ \left. L\lambda \in L^2([0, T]; H^{-1}(\Omega)), \lambda(\cdot, 0) = \lambda_t(\cdot, 0) = 0, \lambda|_{\Gamma_T} \in L^2(\Gamma_T) \right\}.$$

Λ is a Hilbert space endowed with the following inner product

$$\langle \lambda, \bar{\lambda} \rangle_\Lambda := \int_0^T \langle L\lambda(t), L\bar{\lambda}(t) \rangle_{H^{-1}(\Omega)} dt + \iint_{\Gamma_T} \lambda \bar{\lambda} d\sigma dt, \quad \forall \lambda, \bar{\lambda} \in \Lambda$$

using notably that

$$\|\lambda\|_{L^2(Q_T)} \leq C_{\Omega, T} \sqrt{\langle \lambda, \lambda \rangle_\Lambda}, \quad \forall \lambda \in \Lambda \quad (34)$$

for some positive constant $C_{\Omega, T}$. We denote $\|\lambda\|_\Lambda := \sqrt{\langle \lambda, \lambda \rangle_\Lambda}$.

Stabilized mixed formulation "à la Barbosa-Hughes" - 2

Then, $\forall \alpha \in (0, 1)$, we consider the following mixed formulation: find $(\varphi, \lambda) \in (\Phi, \Lambda)$

$$\begin{cases} a_{r,\alpha}(\varphi, \bar{\varphi}) + b_{\alpha}(\bar{\varphi}, \lambda) & = l_1(\bar{\varphi}), & \forall \bar{\varphi} \in \Phi \\ b_{\alpha}(\varphi, \bar{\lambda}) - c_{\alpha}(\lambda, \bar{\lambda}) & = 0, & \forall \bar{\lambda} \in \Lambda, \end{cases} \quad (35)$$

where

$$a_{r,\alpha} : \Phi \times \Phi \rightarrow \mathbb{R}, \quad a_{r,\alpha}(\varphi, \bar{\varphi}) = (1 - \alpha) \iint_{\Gamma_T} \partial_{\nu} \varphi \partial_{\nu} \bar{\varphi} d\sigma dt + r \iint_{Q_T} L^* \varphi L^* \bar{\varphi} dx dt \quad (36)$$

$$b_{\alpha} : \Phi \times \Lambda \rightarrow \mathbb{R}, \quad b_{\alpha}(\varphi, \lambda) = \iint_{Q_T} L^* \varphi \lambda dx dt - \alpha \iint_{\Gamma_T} \partial_{\nu} \varphi \lambda d\sigma dt \quad (37)$$

$$c_{\alpha} : \Lambda \times \Lambda \rightarrow \mathbb{R}, \quad c_{\alpha}(\lambda, \bar{\lambda}) = \alpha \int_0^T \langle L\lambda(t), L\bar{\lambda}(t) \rangle_{H^{-1}(\Omega)} dt + \alpha \iint_{\Gamma_T} \lambda \bar{\lambda} d\sigma dt \quad (38)$$

Proposition

$\forall \alpha \in (0, 1)$, the stabilized mixed formulation is well-posed. Moreover, the unique pair $(\varphi, \lambda) \in \Phi \times \Lambda$ satisfies

$$\theta \|\varphi\|_{\Phi}^2 + \alpha \|\lambda\|_{\Lambda}^2 \leq \frac{(1 - \alpha)^2 + \alpha\theta}{\theta} \|y_0, y_1\|_{L^2 \times H^{-1}}^2 \quad (39)$$

with $\theta := \min(1 - \alpha, r/\eta)$.

Proposition

If $\alpha \in (0, 1)$, the solution $(\varphi, \lambda) \in \Phi \times L^2(\Omega)$ coincides with the stabilized solution $(\varphi_\alpha, \lambda_\alpha) \in \Phi \times \Lambda$

Proposition

$\forall \alpha \in (0, 1)$, the stabilized mixed formulation is well-posed. Moreover, the unique pair $(\varphi, \lambda) \in \Phi \times \Lambda$ satisfies

$$\theta \|\varphi\|_{\Phi}^2 + \alpha \|\lambda\|_{\Lambda}^2 \leq \frac{(1 - \alpha)^2 + \alpha\theta}{\theta} \|y_0, y_1\|_{L^2 \times H^{-1}}^2 \quad (39)$$

with $\theta := \min(1 - \alpha, r/\eta)$.

Proposition

If $\alpha \in (0, 1)$, the solution $(\varphi, \lambda) \in \Phi \times L^2(\Omega)$ coincides with the stabilized solution $(\varphi_\alpha, \lambda_\alpha) \in \Phi \times \Lambda$

Stabilized mixed formulation "à la Barbosa-Hughes" - Numerical approximation

$$\alpha \in (0, 1), r > 0.$$

$$\Phi_h \subset \Phi, \quad \tilde{\Lambda}_h \subset \Lambda, \quad \forall h > 0.$$

Find $(\varphi_h, \lambda_h) \in \Phi_h \times \tilde{\Lambda}_h$ solution of

$$\begin{cases} a_{r,\alpha}(\varphi_h, \bar{\varphi}_h) + b_\alpha(\lambda_h, \bar{\varphi}_h) = l_1(\bar{\varphi}_h), & \forall \bar{\varphi}_h \in \Phi_h \\ b_\alpha(\bar{\lambda}_h, \varphi_h) - c_\alpha(\lambda_h, \bar{\lambda}_h) = 0, & \forall \bar{\lambda}_h \in \tilde{\Lambda}_h. \end{cases} \quad (40)$$

In view of the properties of $a_{r,\alpha}$, c_α , l_1 , this formulation is well-posed.

$$\tilde{\Lambda}_h = \{\lambda \in \Phi_h; \lambda(\cdot, 0) = \lambda_t(\cdot, 0) = 0\}. \quad (41)$$

Proposition (BFS element for $N = 1$ - Rate of convergence in $\Phi \times \Lambda$)

Let $h > 0$, let $k \leq 2$ be a positive integer and $\alpha \in (0, 1)$. Let (y, λ) and (y_h, λ_h) be the solution of (35) and (40) respectively. If (y, λ) belongs to $H^{k+2}(Q_T) \times H^{k+2}(Q_T)$, then $\exists K = K(\|\varphi\|_{H^{k+2}(Q_T)}, \alpha, r, \eta)$ independent of h , such that

$$\|\varphi - \varphi_h\|_\Phi + \|\lambda - \lambda_h\|_\Lambda \leq Kh^k. \quad (42)$$

Remark - no δ_h here !!!! $r > 0$ is arbitrary

Stabilized mixed formulation "à la Barbosa-Hughes" - Numerical approximation

$$\alpha \in (0, 1), r > 0.$$

$$\Phi_h \subset \Phi, \quad \tilde{\Lambda}_h \subset \Lambda, \quad \forall h > 0.$$

Find $(\varphi_h, \lambda_h) \in \Phi_h \times \tilde{\Lambda}_h$ solution of

$$\begin{cases} a_{r,\alpha}(\varphi_h, \bar{\varphi}_h) + b_\alpha(\lambda_h, \bar{\varphi}_h) = l_1(\bar{\varphi}_h), & \forall \bar{\varphi}_h \in \Phi_h \\ b_\alpha(\bar{\lambda}_h, \varphi_h) - c_\alpha(\lambda_h, \bar{\lambda}_h) = 0, & \forall \bar{\lambda}_h \in \tilde{\Lambda}_h. \end{cases} \quad (40)$$

In view of the properties of $a_{r,\alpha}$, c_α , l_1 , this formulation is well-posed.

$$\tilde{\Lambda}_h = \{\lambda \in \Phi_h; \lambda(\cdot, 0) = \lambda_t(\cdot, 0) = 0\}. \quad (41)$$

Proposition (BFS element for $N = 1$ - Rate of convergence in $\Phi \times \Lambda$)

Let $h > 0$, let $k \leq 2$ be a positive integer and $\alpha \in (0, 1)$. Let (y, λ) and (y_h, λ_h) be the solution of (35) and (40) respectively. If (y, λ) belongs to $H^{k+2}(Q_T) \times H^{k+2}(Q_T)$, then $\exists K = K(\|\varphi\|_{H^{k+2}(Q_T)}, \alpha, r, \eta)$ independent of h , such that

$$\|\varphi - \varphi_h\|_\Phi + \|\lambda - \lambda_h\|_\Lambda \leq Kh^k. \quad (42)$$

Remark - no δ_h here !!!! $r > 0$ is arbitrary

Remark 1: The situation may be simpler with a different cost !?

$$\begin{cases} \text{Minimize } J(y, v) = \frac{1}{2} \iint_{Q_T} |y|^2 dx dt + \frac{1}{2} \int_0^T \int_{\Gamma_0} |v|^2 d\sigma dt \\ \text{Subject to } (y, v) \in \mathcal{C}(y_0, y_1; T) \end{cases} \quad (43)$$

$$v = \frac{\partial \varphi}{\partial \nu} \text{ in } (0, T) \times \Gamma_0 \text{ and } y = \mu \text{ in } Q_T.$$

$$\begin{cases} \text{Minimize } J^*(\mu, \varphi_0, \varphi_1) = \frac{1}{2} \iint_{Q_T} |\mu|^2 dx dt + \frac{1}{2} \int_0^T \int_{\Gamma_0} \left| \frac{\partial \varphi}{\partial \nu} \right|^2 d\sigma dt \\ \quad + \langle (\varphi_0, \varphi_1), (y_0, y_1) \rangle \\ \text{Subject to } (\mu, \varphi_0, \varphi_1) \in L^2(Q_T) \times \mathbf{V}, \end{cases} \quad (44)$$

where φ solves the nonhomogeneous backward problem

$$L^* \varphi = \mu \text{ in } Q_T, \quad \varphi = 0 \text{ on } \Sigma_T, \quad (\varphi(\cdot, 0), \varphi'(\cdot, 0)) = (\varphi_0, \varphi_1) \quad (45)$$

Remark 1: The situation may be much simpler with a different cost !!?!

7

Replacing μ by $L^*\varphi$ and minimiz over φ lead to

$$\left\{ \begin{array}{l} \text{Minimize } J_1^*(\varphi) = \frac{1}{2} \iint_{Q_T} |L^*\varphi|^2 dx dt + \frac{1}{2} \int_0^T \int_{\Gamma_0} \left| \frac{\partial \varphi}{\partial \nu} \right|^2 d\sigma dt \\ \quad \quad \quad + \langle (\varphi(\cdot, 0), \varphi_t(\cdot, 0)), (y_0, y_1) \rangle \\ \text{Subject to } \varphi \in \Phi \end{array} \right. \quad (46)$$

and to the **well-posed variational formulation**: find $\varphi \in \Phi$ such that

$$\underbrace{\iint_{Q_T} L^*\varphi L^*\bar{\varphi} dx dt + \int_0^T \int_{\Gamma_0} \frac{\partial \varphi}{\partial \nu} \frac{\partial \bar{\varphi}}{\partial \nu} d\sigma dt}_{a_r(\varphi, \varphi) \text{ with } r=1} = \langle (\bar{\varphi}(\cdot, 0), \bar{\varphi}_t(\cdot, 0)), (y_0, y_1) \rangle, \quad \forall \bar{\varphi} \in \Phi \quad (47)$$

Non constant coefficient: $Ly := y_{tt} - (c(x)y_x)_x + d(x, t)y$, $c \in C^1(\overline{\Omega})$

$$c(x) = \begin{cases} 1 & x \in [0, 0.45] \\ \in [1., 5.] \quad (c'(x) > 0), & x \in (0.45, 0.55) \\ 5 & x \in [0.55, 1] \end{cases} \quad (48)$$




Figure: Approximation of the control solution \hat{y}_h over $Q_T - h = (1/80, 1/80)$.

Remark 2: The distributed case

$$Ly = v \mathbf{1}_{q_T}, \quad q_T = \omega \times (0, T) \subset \Omega \times (0, T)$$

$$\left\{ \begin{array}{l} \min J^*(\varphi) = \frac{1}{2} \int_0^T \int_\omega |\varphi|^2 dx dt + \langle y_0, \varphi_t(\cdot, 0) \rangle_{H^1, H^{-1}} - \langle y_1, \varphi(\cdot, 0) \rangle_{L^2} \\ \text{Subject to } \varphi \in \mathbf{W} := \left\{ \varphi : \varphi \in L^2(q_T), \varphi|_{\Sigma_T} = 0, L^* \varphi = 0 \in L^2(0, T, H^{-1}(\Omega)) \right\} \end{array} \right\} \quad (49)$$

Optimal control : $v = \varphi \mathbf{1}_{q_T}$

Generalized observability inequality : $\exists C_{obs} > 0$ s.t.


$$\|\varphi_0, \varphi_1\|_{L^2(\Omega) \times H^{-1}(\Omega)}^2 \leq C_{obs} \left(\|\varphi\|_{L^2(q_T)}^2 + \|L^* \varphi\|_{L^2(0, T; H^{-1})}^2 \right), \quad \forall \varphi \in \Phi$$

Lagrange Multiplier :

$$b(\varphi, \lambda) = \int_0^T \langle \lambda(\cdot, t), L^* \varphi(\cdot, t) \rangle_{H_0^1(\Omega), H^{-1}(\Omega)} dt, \quad \lambda \in L^2(0, T; H_0^1(\Omega))$$

The distributed case : Non cylindrical situation in 1D with constant coefficient

The variational approach is well-adapted to the non cylindrical situation. ^{8 9}



Time dependent domains $q_T \subset Q_T = \Omega \times (0, T)$ and corresponding meshes

⁸ C. Castro, N. Cindea, AM, [Controllability of the 1D wave equation with inner moving force](#), SICON (2014)]

⁹ G. Lebeau, J. Le Rousseau, P. Terpolilli, E. Trélat, [Geometric control condition for the wave equation with a time-dependent domain](#), (2016)

Remark 3 : Inverse problems -

Given a **distributed observation** $y_{obs} \in L^2(q_T)$, $f \in X := L^2(H^{-1})$, **reconstruct** y such that

$$Ly = f \quad \text{in } Q_T, \quad y = 0 \quad \text{on } \Sigma_T, \quad y - y_{obs} = 0 \quad \text{on } q_T$$

$$(LS) \quad \begin{cases} \text{minimize} & J(y_0, y_1) := \frac{1}{2} \|y - y_{obs}\|_{L^2(q_T)}^2 \\ \text{subject to} & (y_0, y_1) \in L^2 \times H^{-1} \text{ where } Ly - f = 0 \end{cases}$$

The "**Discretization then Inverse problem**" procedure is discussed in [L. Baudouin, M. De Buhan, S. Ervedoza, 2013]

Keeping y as the main variable ¹⁰....

$$(P) \quad \begin{cases} \inf J(y) := \frac{1}{2} \|y - y_{obs}\|_{L^2(q_T)}^2 + \frac{r}{2} \|Ly - f\|_X^2, \quad r \geq 0 \\ \text{subject to} \quad y \in W := \{y \in Z; Ly - f = 0 \text{ in } X\} \end{cases}$$

The multiplier $\lambda \in X'$ is a "**measure**" of the quality of y_{obs} to reconstruct y .

¹⁰N. Cindea, AM, Inverse problem for linear hyperbolic equations using mixed formulations, Inverse Problems, (2015).

Remark 3 : Inverse problems -

Given a **distributed observation** $y_{obs} \in L^2(q_T)$, $f \in X := L^2(H^{-1})$, **reconstruct** y such that

$$Ly = f \quad \text{in } Q_T, \quad y = 0 \quad \text{on } \Sigma_T, \quad y - y_{obs} = 0 \quad \text{on } q_T$$

$$(LS) \quad \begin{cases} \text{minimize} & J(y_0, y_1) := \frac{1}{2} \|y - y_{obs}\|_{L^2(q_T)}^2 \\ \text{subject to} & (y_0, y_1) \in L^2 \times H^{-1} \text{ where } Ly - f = 0 \end{cases}$$

The "**Discretization then Inverse problem**" procedure is discussed in [L. Baudouin, M. De Buhan, S. Ervedoza, 2013]

Keeping y as the main variable ¹⁰....

$$(P) \quad \begin{cases} \inf J(y) := \frac{1}{2} \|y - y_{obs}\|_{L^2(q_T)}^2 + \frac{r}{2} \|Ly - f\|_X^2, & r \geq 0 \\ \text{subject to} & y \in W := \{y \in Z; Ly - f = 0 \text{ in } X\} \end{cases}$$

The multiplier $\lambda \in X'$ is a "**measure**" of the quality of y_{obs} to reconstruct y .

¹⁰N. Cindea, AM, Inverse problem for linear hyperbolic equations using mixed formulations, Inverse Problems, (2015).

Remark 3 : Inverse problems -

Given a **distributed observation** $y_{obs} \in L^2(q_T)$, $f \in X := L^2(H^{-1})$, **reconstruct** y such that

$$Ly = f \quad \text{in } Q_T, \quad y = 0 \quad \text{on } \Sigma_T, \quad y - y_{obs} = 0 \quad \text{on } q_T$$

$$(LS) \quad \begin{cases} \text{minimize} & J(y_0, y_1) := \frac{1}{2} \|y - y_{obs}\|_{L^2(q_T)}^2 \\ \text{subject to} & (y_0, y_1) \in L^2 \times H^{-1} \text{ where } Ly - f = 0 \end{cases}$$

The "**Discretization then Inverse problem**" procedure is discussed in [L. Baudouin, M. De Buhan, S. Ervedoza, 2013]


Keeping y as the main variable ¹⁰....

$$(P) \quad \begin{cases} \inf J(y) := \frac{1}{2} \|y - y_{obs}\|_{L^2(q_T)}^2 + \frac{r}{2} \|Ly - f\|_X^2, & r \geq 0 \\ \text{subject to} & y \in W := \{y \in Z; Ly - f = 0 \text{ in } X\} \end{cases}$$


The multiplier $\lambda \in X'$ is a "**measure**" of the quality of y_{obs} to reconstruct y .

¹⁰N. Cindea, AM, Inverse problem for linear hyperbolic equations using mixed formulations, Inverse Problems, (2015).

2D example - Observation on q_T



(a)



(b)

Mesh number	0	1	2
Number of elements	5 730	44 900	196 040
Number of nodes	3 432	24 633	103 566

Characteristics of the three meshes associated with Q_T .


2D example - Observation on q_T

$$\begin{cases} -\Delta y_0 = 10, & \text{in } \Omega \\ y_0 = 0, & \text{on } \partial\Omega, \end{cases} \quad y_1 = 0. \quad (50)$$


Mesh number	0	1	2
$\frac{\ \bar{y}_h - y_h\ _{L^2(Q_T)}}{\ \bar{y}_h\ _{L^2(Q_T)}}$	1.88×10^{-1}	8.04×10^{-2}	5.41×10^{-2}
$\ Ly_h\ _{L^2(Q_T)}$	3.21	2.01	1.17
$\ \lambda_h\ _{L^2(Q_T)}$	8.26×10^{-5}	3.62×10^{-5}	2.24×10^{-5}

$$r = h^2 - T = 2$$

2D example - Observation on q_T



(a)



(b)

y and y_h in Q_T

Example 2 - $N = 2$ - The Bunimovich's stadium - Reconstruction from partial boundary observation

$$T = 3$$




Figure: Bunimovich's stadium and the subset Γ of $\partial\Omega$ on which the observations are available. Example of mesh of the domain Q_T .

Example 2 - $N = 2$ - Recovering of the initial data

$T = 3$





Figure: (a) Initial data y_0 given by (50). (b) Reconstructed initial data $y_h(\cdot, 0)$.

Remark 4 : Simultaneous reconstruction of source and solution

Similarly, the "generalized observability inequality" [Yamamoto-Zhang, 2001]¹¹

$$\|\mu\|_{H^{-1}(\Omega)}^2 \leq C_{obs} \left(\|c(x)\partial_\nu y\|_{L^2(\Gamma_T)}^2 + \|Ly - \sigma\mu\|_{L^2(Q_T)}^2 \right), \quad \forall (y, \mu) \in Y. \quad (\mathcal{H}_2)$$

where

$$Y := \left\{ (y, \mu); y \in C(H_0^1) \cap C^1(L^2), \mu \in H^{-1}(\Omega), Ly - \sigma\mu \in L^2(Q_T), y(\cdot, 0) = y_t(\cdot, 0) = 0 \right\}. \quad (51)$$

allows to reconstruct with robustness ¹² and simultaneously the spatial part $\mu(x)$ of the source and the solution y of

$$Ly := \sigma(t)\mu(x), \text{ in } Q_T, \quad y = 0 \text{ on } \Sigma_T, \quad (y(\cdot, 0), y_t(\cdot, 0)) = (0, 0), \text{ in } \Omega. \quad (52)$$

from the observation $\partial_\nu y|_{\Gamma_0}$, assuming $\sigma \in C^1([0, T]), \sigma(0) \neq 0$.

¹¹Yamamoto, Zhang, Global uniqueness and stability for an inverse wave source problem for less regular data. 2001

¹²N. Cindea, AM, Simultaneous reconstruction of the solution and the source of hyperbolic equations from boundary measurements: a robust numerical approach, Inverse Problems, 2016.

Parabolic case

$\Omega \subset \mathbb{R}^N$; $Q_T = \Omega \times (0, T)$; $q_T = \omega \times (0, T)$

$$\begin{cases} y_t - \nabla \cdot (c(x)\nabla y) + d(x, t)y = v 1_\omega, & \text{in } Q_T, \\ y = 0, & \text{in } \Sigma_T, \\ y(x, 0) = y_0(x), & \text{in } \Omega. \end{cases} \quad (53)$$

$c := (c_{i,j}) \in C^1(\bar{\Omega}; \mathcal{M}_N(\mathbb{R}))$; $(c(x)\xi, \xi) \geq c_0|\xi|^2$ in $\bar{\Omega}$ ($c_0 > 0$),

$d \in L^\infty(Q_T)$, $y_0 \in L^2(\Omega)$;

$v = v(x, t)$ is the *control* $y = y(x, t)$ is the associated state.

The linear manifold

$$\mathcal{C}(y_0, T) = \{ (y, v) : v \in L^2(q_T), y \text{ solves (53) and satisfies } y(T, \cdot) = 0 \}.$$

non empty (see FURSIKOV-IMANUVILOV'96, ROBBIANO-LEBEAU'95)).

NOTATIONS -

$Ly := y_t - \nabla \cdot (c(x)\nabla y) + d(x, t)y$; $L^*\varphi := -\varphi_t - \nabla \cdot (c(x)\nabla \varphi) + d(x, t)\varphi$

Parabolic case

$$\Omega \subset \mathbb{R}^N; Q_T = \Omega \times (0, T); q_T = \omega \times (0, T)$$

$$\begin{cases} y_t - \nabla \cdot (c(x)\nabla y) + d(x, t)y = v 1_\omega, & \text{in } Q_T, \\ y = 0, & \text{in } \Sigma_T, \\ y(x, 0) = y_0(x), & \text{in } \Omega. \end{cases} \quad (53)$$

$$c := (c_{i,j}) \in C^1(\bar{\Omega}; \mathcal{M}_N(\mathbb{R})); (c(x)\xi, \xi) \geq c_0|\xi|^2 \text{ in } \bar{\Omega} \quad (c_0 > 0),$$

$$d \in L^\infty(Q_T), y_0 \in L^2(\Omega);$$

$v = v(x, t)$ is the *control* $y = y(x, t)$ is the associated state.

The linear manifold

$$\mathcal{C}(y_0, T) = \{ (y, v) : v \in L^2(q_T), y \text{ solves (53) and satisfies } y(T, \cdot) = 0 \}.$$

non empty (see FURSIKOV-IMANUVILOV'96, ROBBIANO-LEBEAU'95)).

NOTATIONS -

$$Ly := y_t - \nabla \cdot (c(x)\nabla y) + d(x, t)y; \quad L^* \varphi := -\varphi_t - \nabla \cdot (c(x)\nabla \varphi) + d(x, t)\varphi$$

Minimal L^2 norm control

$$\inf_{\phi_T \in H} J^*(\phi_T), \quad J^*(\phi_T) = \frac{1}{2} \int_{Q_T} \phi^2 dx dt + \int_{\Omega} \phi(0, \cdot) y_0 dx$$

where ϕ solves the backward system

$$L^* \phi = 0 \text{ in } Q_T, \quad \phi = 0 \text{ on } \Sigma_T, \quad \phi(T, \cdot) = \phi_T \text{ in } \Omega.$$

The Hilbert space H is defined as the completion of $\mathcal{D}(\Omega)$ with respect to the norm

$$\|\phi_T\|_H = \left(\int_{Q_T} \phi^2(t, x) dx dt \right)^{1/2}.$$

From the observability inequality

$$\|\phi(0, \cdot)\|_{L^2(\Omega)}^2 \leq C_{obs}(\omega, T) \|\phi_T\|_H^2 \quad \forall \phi_T \in L^2(\Omega),$$

J^* is coercive on H . The HUM control is given by $v = \phi \mathcal{X}_\omega$ on Q_T .

The problem is ill-posed: H is "huge": In 1D, from [Micu, Zuazua, 2011] ¹³

the set of initial data y_0 , for which the corresponding ϕ_T , minimizer of J^* , does not belong to any negative Sobolev spaces, is dense in $L^2(0, 1)$!!!

¹³S. Micu, E. Zuazua, *Regularity issues for the null-controllability of the linear 1-d heat equation*, 2011

Minimal L^2 norm control

$$\inf_{\phi_T \in H} J^*(\phi_T), \quad J^*(\phi_T) = \frac{1}{2} \int_{Q_T} \phi^2 dx dt + \int_{\Omega} \phi(0, \cdot) y_0 dx$$

where ϕ solves the backward system

$$L^* \phi = 0 \text{ in } Q_T, \quad \phi = 0 \text{ on } \Sigma_T, \quad \phi(T, \cdot) = \phi_T \text{ in } \Omega.$$

The Hilbert space H is defined as the completion of $\mathcal{D}(\Omega)$ with respect to the norm

$$\|\phi_T\|_H = \left(\int_{Q_T} \phi^2(t, x) dx dt \right)^{1/2}.$$

From the observability inequality

$$\|\phi(0, \cdot)\|_{L^2(\Omega)}^2 \leq C_{obs}(\omega, T) \|\phi_T\|_H^2 \quad \forall \phi_T \in L^2(\Omega),$$

J^* is coercive on H . The HUM control is given by $v = \phi \mathcal{X}_\omega$ on Q_T .

The problem is ill-posed: H is "huge": In 1D, from [Micu, Zuazua, 2011] ¹³

the set of initial data y_0 , for which the corresponding ϕ_T , minimizer of J^* , does not belong to any negative Sobolev spaces, is dense in $L^2(0, 1)$!!!

¹³S. Micu, E. Zuazua, *Regularity issues for the null-controllability of the linear 1-d heat equation*, 2011

Minimal L^2 norm control

$$\inf_{\phi_T \in H} J^*(\phi_T), \quad J^*(\phi_T) = \frac{1}{2} \int_{Q_T} \phi^2 dx dt + \int_{\Omega} \phi(0, \cdot) y_0 dx$$

where ϕ solves the backward system

$$L^* \phi = 0 \text{ in } Q_T, \quad \phi = 0 \text{ on } \Sigma_T, \quad \phi(T, \cdot) = \phi_T \text{ in } \Omega.$$

The Hilbert space H is defined as the completion of $\mathcal{D}(\Omega)$ with respect to the norm

$$\|\phi_T\|_H = \left(\int_{Q_T} \phi^2(t, x) dx dt \right)^{1/2}.$$

From the observability inequality

$$\|\phi(0, \cdot)\|_{L^2(\Omega)}^2 \leq C_{obs}(\omega, T) \|\phi_T\|_H^2 \quad \forall \phi_T \in L^2(\Omega),$$

J^* is coercive on H . The HUM control is given by $v = \phi \chi_{\omega}$ on Q_T .

The problem is ill-posed: H is "huge": In 1D, from [Micu, Zuazua, 2011] ¹³

the set of initial data y_0 , for which the corresponding ϕ_T , minimizer of J^* , does not belong to any negative Sobolev spaces, is dense in $L^2(0, 1)$!!!

¹³S. Micu, E. Zuazua, *Regularity issues for the null-controllability of the linear 1-d heat equation*, 2011

$N = 1 - L^2(q_T)$ -norm of the HUM control with respect to time




Figure: $y_0(x) = \sin(\pi x) - T = 1 - \omega = (0.2, 0.8) - t \rightarrow \|v(\cdot, t)\|_{L^2(0,1)}$ in $[0, T]$

$N = 1 - L^2$ -norm of the HUM control with respect to time: Zoom near T




Figure: $y_0(x) = \sin(\pi x) - T = 1 - \omega = (0.2, 0.8) - t \rightarrow \|v(\cdot, t)\|_{L^2(0,1)}$ in $[0.92T, T]$

Optimal backward solution ϕ on $\partial\omega \times [0, T]$

$$T = 1, \quad y_0(x) = \sin(\pi x), \quad a(x) = a_0 = 1/10, \quad \omega = (0.2, 0.8)$$




Figure: $T = 1 - \omega = (0.2, 0.8) - \phi^N(\cdot, 0.8)$ for $N = 80$ on $[0, T]$ (**Left**) and on $[0.92T, T]$ (**Right**).

[Carthel-Glowinski-Lions'94, JOTA], [AM-Zuazua'11, Inverse Problems]

Remedies : Carleman weights !!

Change of the norm : framework of Fursikov-Imanuvilov'96 ¹⁴


$$\begin{cases} \text{Minimize } J(y, v) = \frac{1}{2} \iint_{Q_T} \rho^2 |y|^2 dx dt + \frac{1}{2} \iint_{q_T} \rho_0^2 |v|^2 dx dt \\ \text{Subject to } (y, v) \in \mathcal{C}(y_0, T). \end{cases} \quad (54)$$

where ρ, ρ_0 are non-negative continuous weights functions such that $\rho, \rho_0 \in L^\infty(Q_{T-\delta}) \quad \forall \delta > 0$.

$$\Rightarrow \min_{p \in P} J^*(p) := \frac{1}{2} \iint_{Q_T} \rho^{-2} |L^* p|^2 dx dt + \frac{1}{2} \iint_{q_T} \rho_0^{-2} |p|^2 dx dt - (y_0, p(\cdot, 0)), \quad (55)$$

\Rightarrow

$$\begin{cases} \text{find } p \in P \text{ s.t.} \\ \iint_{Q_T} \rho^{-2} L^* p L^* q dx dt + \iint_{q_T} \rho_0^{-2} p q dx dt = (y_0, q(\cdot, 0)), \quad \forall q \in P \end{cases} \quad (56)$$

¹⁴A.V. Fursikov and O. Yu. Imanuvilov, *Controllability of Evolution Equations*, Lecture Notes Series, number 34. Seoul National University, Korea, (1996) 1-163. 

Remedies : Carleman weights !!

Change of the norm : framework of Fursikov-Imanuvilov'96 ¹⁴


$$\begin{cases} \text{Minimize } J(y, v) = \frac{1}{2} \iint_{Q_T} \rho^2 |y|^2 dx dt + \frac{1}{2} \iint_{q_T} \rho_0^2 |v|^2 dx dt \\ \text{Subject to } (y, v) \in \mathcal{C}(y_0, T). \end{cases} \quad (54)$$

where ρ, ρ_0 are non-negative continuous weights functions such that $\rho, \rho_0 \in L^\infty(Q_{T-\delta}) \quad \forall \delta > 0$.

$$\Rightarrow \min_{p \in P} J^*(p) := \frac{1}{2} \iint_{Q_T} \rho^{-2} |L^* p|^2 dx dt + \frac{1}{2} \iint_{q_T} \rho_0^{-2} |p|^2 dx dt - (y_0, p(\cdot, 0)), \quad (55)$$

\Rightarrow

$$\begin{cases} \text{find } p \in P \text{ s.t.} \\ \iint_{Q_T} \rho^{-2} L^* p L^* q dx dt + \iint_{q_T} \rho_0^{-2} p q dx dt = (y_0, q(\cdot, 0)), \quad \forall q \in P \end{cases} \quad (56)$$

¹⁴A.V. Fursikov and O. Yu. Imanuvilov, *Controllability of Evolution Equations*, Lecture Notes Series, number 34. Seoul National University, Korea, (1996) 1-163. 

Primal (direct) approach

Following Fursikov-Imanuvilov'96, we assume Carleman type weights :

$$\left\{ \begin{array}{l} \rho(x, t) = \exp\left(\frac{\beta(x)}{T-t}\right), \quad \rho_0(x, t) = (T-t)^{3/2}\rho(x, t), \quad \beta(x) = K_1 \left(e^{K_2} - e^{\beta_0(x)}\right) \\ \text{where the } K_i \text{ are sufficiently large positive constants (depending on } T, c_0 \text{ and } \|c\|_{C^1}) \\ \text{and } \beta_0 \in C^\infty(\bar{\Omega}), \beta_0 > 0 \text{ in } \Omega, (\beta_0)|_{\partial\Omega} = 0, |\nabla\beta_0| > 0 \text{ outside } \omega. \end{array} \right. \quad (57)$$

We introduce

$$P_0 = \{q \in C^2(\bar{Q}_T) : q = 0 \text{ on } \Sigma_T\}.$$

In this linear space, the bilinear form

$$(p, q)_P := \iint_{Q_T} \rho^{-2} L^* p L^* q \, dx \, dt + \iint_{Q_T} \rho_0^{-2} p q \, dx \, dt$$

is a scalar product (unique continuation property).

Let P be the completion of P_0 for this scalar product.

Primal (direct) approach

Following Fursikov-Imanuvilov'96, we assume Carleman type weights :

$$\left\{ \begin{array}{l} \rho(x, t) = \exp\left(\frac{\beta(x)}{T-t}\right), \quad \rho_0(x, t) = (T-t)^{3/2} \rho(x, t), \quad \beta(x) = K_1 \left(e^{K_2} - e^{\beta_0(x)}\right) \\ \text{where the } K_i \text{ are sufficiently large positive constants (depending on } T, c_0 \text{ and } \|c\|_{C^1}) \\ \text{and } \beta_0 \in C^\infty(\bar{\Omega}), \beta_0 > 0 \text{ in } \Omega, (\beta_0)|_{\partial\Omega} = 0, |\nabla\beta_0| > 0 \text{ outside } \omega. \end{array} \right. \quad (57)$$

We introduce

$$P_0 = \{q \in C^2(\bar{Q}_T) : q = 0 \text{ on } \Sigma_T\}.$$

In this linear space, the bilinear form

$$(p, q)_P := \iint_{Q_T} \rho^{-2} L^* p L^* q \, dx \, dt + \iint_{Q_T} \rho_0^{-2} p q \, dx \, dt$$

is a scalar product (unique continuation property).

Let P be the completion of P_0 for this scalar product.

Lemma (Fursikov-Imanuvilov'96, Fernández-Cara-Guerrero'06)

Let ρ and ρ_0 be given by (57). For any $\delta > 0$, one has

$$P \hookrightarrow C^0([0, T - \delta]; H_0^1(\Omega)),$$

where the embedding is continuous. In particular, there exists $C > 0$, only depending on ω , T , c_0 and $\|c\|_{C^1}$, such that, for all $q \in P$,

$$\|q(\cdot, 0)\|_{H_0^1(\Omega)}^2 \leq C \left(\iint_{Q_T} \rho^{-2} |L^* q|^2 dx dt + \iint_{q_T} \rho_0^{-2} |q|^2 dx dt \right). \quad (58)$$

Primal (direct) approach

Proposition

Let ρ and ρ_0 be given by (57). Let (y, v) be the corresponding optimal pair for J . Then there exists $p \in P$ such that

$$y = \rho^{-2} L^* p, \quad v = -\rho_0^{-2} p|_{q_T}. \quad (59)$$

The function p is the unique solution in P of

$$\iint_{Q_T} \rho^{-2} L^* p L^* q \, dx \, dt + \iint_{q_T} \rho_0^{-2} p q \, dx \, dt = \int_0^1 y_0(x) q(x, 0) \, dx, \quad \forall q \in P \quad (60)$$

Remark

p solves, at least in \mathcal{D}' , the following differential problem, that is second order in time and fourth order in space:

$$\begin{cases} L(\rho^{-2} L^* p) + \rho_0^{-2} p 1_\omega = 0, & (x, t) \in (0, 1) \times (0, T) \\ p(x, t) = 0, \quad (-\rho^{-2} L^* p)(x, t) = 0 & (x, t) \in \{0, 1\} \times (0, T) \\ (-\rho^{-2} L^* p)(x, 0) = y_0(x), \quad (-\rho^{-2} L^* p)(x, T) = 0, & x \in (0, 1). \end{cases} \quad (61)$$

Primal (direct) approach

Proposition

Let ρ and ρ_0 be given by (57). Let (y, v) be the corresponding optimal pair for J . Then there exists $p \in P$ such that

$$y = \rho^{-2} L^* p, \quad v = -\rho_0^{-2} p|_{q_T}. \quad (59)$$

The function p is the unique solution in P of

$$\iint_{Q_T} \rho^{-2} L^* p L^* q \, dx \, dt + \iint_{q_T} \rho_0^{-2} p q \, dx \, dt = \int_0^1 y_0(x) q(x, 0) \, dx, \quad \forall q \in P \quad (60)$$

Remark

p solves, at least in \mathcal{D}' , the following differential problem, that is second order in time and fourth order in space:

$$\begin{cases} L(\rho^{-2} L^* p) + \rho_0^{-2} p 1_\omega = 0, & (x, t) \in (0, 1) \times (0, T) \\ p(x, t) = 0, \quad (-\rho^{-2} L^* p)(x, t) = 0 & (x, t) \in \{0, 1\} \times (0, T) \\ (-\rho^{-2} L^* p)(x, 0) = y_0(x), \quad (-\rho^{-2} L^* p)(x, T) = 0, & x \in (0, 1). \end{cases} \quad (61)$$

Conformal approximation

Let \mathcal{T}_h be a uniform triangulation, with $\overline{Q_T} = \bigcup_{K \in \mathcal{T}_h} K$.

The following (conformal) finite element approximations of the space P are introduced:

$$P_h = \{ q_h \in C_{x,t}^{1,0}(\overline{Q_T}) : q_h|_K \in (\mathbb{P}_{3,x} \otimes \mathbb{P}_{1,t})(K) \quad \forall K \in \mathcal{T}_h, \quad q_h|_{\Sigma_T} \equiv 0 \},$$

The variational equality (60) is approximated as follows:

$$\left\{ \begin{array}{l} \iint_{Q_T} \rho^{-2} L^* p_h L^* q_h \, dx \, dt + \iint_{q_T} \rho_0^{-2} p_h q_h \, dx \, dt = \int_0^1 y_0(x) q_h(x, 0) \, dx \\ \forall q_h \in P_h; \quad p_h \in P_h. \end{array} \right. \quad (62)$$

Conformal approximation

Let \mathcal{T}_h be a uniform triangulation, with $\overline{Q_T} = \bigcup_{K \in \mathcal{T}_h} K$.

The following (conformal) finite element approximations of the space P are introduced:

$$P_h = \{ q_h \in C_{x,t}^{1,0}(\overline{Q_T}) : q_h|_K \in (\mathbb{P}_{3,x} \otimes \mathbb{P}_{1,t})(K) \quad \forall K \in \mathcal{T}_h, \quad q_h|_{\Sigma_T} \equiv 0 \},$$

The variational equality (60) is approximated as follows:

$$\left\{ \begin{array}{l} \iint_{Q_T} \rho^{-2} L^* p_h L^* q_h \, dx \, dt + \iint_{q_T} \rho_0^{-2} p_h q_h \, dx \, dt = \int_0^1 y_0(x) q_h(x, 0) \, dx \\ \forall q_h \in P_h; \quad p_h \in P_h. \end{array} \right. \quad (62)$$

Conformal approximation

Theorem (Fernandez-Cara, AM)

Let $p_h \in P_h$ be the unique solution to (63). Let us set

$$y_h := \rho^{-2} L^* p_h, \quad v_h := -\rho_0^{-2} p_h \mathbf{1}_{Q_T}.$$

Then one has

$$\|y - y_h\|_{L^2(Q_T)} \rightarrow 0 \text{ and } \|v - v_h\|_{L^2(Q_T)} \rightarrow 0, \quad \text{as } h \rightarrow 0$$

where (y, v) is the minimizer of J .

In practice, we introduce the variable

$m_h := \rho_0^{-1} p_h \in \rho_0^{-1} P_h \subset \rho_0^{-1} P \subset C([0, T], H_0^1(\Omega))$ and we solve

$$\begin{cases} \iint_{Q_T} \rho^{-2} L^*(\rho_0 m_h) L^*(\rho_0 \overline{m}_h) dx dt + \iint_{Q_T} m_h \overline{m}_h dx dt = \int_0^1 y_0 \rho_0(\cdot, 0) \overline{m}_h(\cdot, 0) dx \\ \forall m_h \in \rho_0^{-1} P_h; \quad \overline{m}_h \in \rho_0^{-1} P_h. \end{cases} \quad (63)$$

Conformal approximation

Theorem (Fernandez-Cara, AM)

Let $p_h \in P_h$ be the unique solution to (63). Let us set

$$y_h := \rho^{-2} L^* p_h, \quad v_h := -\rho_0^{-2} p_h \mathbf{1}_{Q_T}.$$

Then one has

$$\|y - y_h\|_{L^2(Q_T)} \rightarrow 0 \text{ and } \|v - v_h\|_{L^2(Q_T)} \rightarrow 0, \quad \text{as } h \rightarrow 0$$

where (y, v) is the minimizer of J .

In practice, we introduce the variable

$m_h := \rho_0^{-1} p_h \in \rho_0^{-1} P_h \subset \rho_0^{-1} P \subset C([0, T], H_0^1(\Omega))$ and we solve

$$\begin{cases} \iint_{Q_T} \rho^{-2} L^*(\rho_0 m_h) L^*(\rho_0 \overline{m}_h) dx dt + \iint_{Q_T} m_h \overline{m}_h dx dt = \int_0^1 y_0 \rho_0(\cdot, 0) \overline{m}_h(\cdot, 0) dx \\ \forall m_h \in \rho_0^{-1} P_h; \quad \overline{m}_h \in \rho_0^{-1} P_h. \end{cases} \quad (63)$$

Theorem (Fernandez-Cara, AM)

Let $p_h \in P_h$ be the unique solution to (63). Let us set

$$y_h := \rho^{-2} L^* p_h, \quad v_h := -\rho_0^{-2} \rho_h \mathbf{1}_{Q_T}.$$

Then one has

$$\|y - y_h\|_{L^2(Q_T)} \rightarrow 0 \text{ and } \|v - v_h\|_{L^2(Q_T)} \rightarrow 0, \quad \text{as } h \rightarrow 0$$

where (y, v) is the minimizer of J .

In practice, we introduce the variable

$m_h := \rho_0^{-1} p_h \in \rho_0^{-1} P_h \subset \rho_0^{-1} P \subset C([0, T], H_0^1(\Omega))$ and we solve

$$\begin{cases} \iint_{Q_T} \rho^{-2} L^*(\rho_0 m_h) L^*(\rho_0 \overline{m}_h) dx dt + \iint_{Q_T} m_h \overline{m}_h dx dt = \int_0^1 y_0 \rho_0(\cdot, 0) \overline{m}_h(\cdot, 0) dx \\ \forall m_h \in \rho_0^{-1} P_h; \quad \overline{m}_h \in \rho_0^{-1} P_h. \end{cases} \quad (63)$$

Experiments with $\omega = (0.2, 0.8)$

$$T = 1/2, y_0(x) \equiv \sin(\pi x), a(x) \equiv 10^{-1}.$$




Figure: $\omega = (0.2, 0.8)$. The state y_h (Left) and the control v_h (Right).

Experiments with $\omega = (0.3, 0.6)$

$$T = 1/2, y_0(x) \equiv \sin(\pi x), a(x) \equiv 10^{-1}.$$




Figure: $\omega = (0.3, 0.6)$. $y(x)$

Experiments with $\omega = (0.3, 0.4)$





Figure: $\omega = (0.3, 0.4)$. The state y_h (Left) and the control v_h (Right).

Experiments with $\omega = (0.2, 0.4)$




Figure: $\omega = (0.2, 0.4)$. $y(x)$

17

18

¹⁷E. Fernández-Cara and AM, *Numerical null controllability of the 1-d heat equation: primal algorithms*, (2013),

¹⁸E. Fernández-Cara and A. Münch, *Numerical null controllability of the 1-d heat equation: Carleman weights and duality*, JOTA, (2013)

Application: Controllability for semi-linear heat equation

19

$$\begin{cases} y_t - 0.1y_{xx} - 5y \log^{1.4}(1 + |y|) = v1_{(0.2,0.8)}, & (x, t) \in (0, 1) \times (0, 1/2) \\ y(x, t) = 0, & (x, t) \in \{0, 1\} \times (0, 1/2) \\ y(x, 0) = 40 \sin(\pi x), & x \in (0, 1). \end{cases} \quad (64)$$

Without control, blow up at $t \approx 0.318$.




Figure: Fixed point method - $h = (1/60, 1/60)$ - $y_0(x) = 40 \sin(\pi x)$ - Control v_h (**Left**) and corresponding controlled solution y_h (**Right**) in Q_T .

L^2 -weighted norm


No contribution of y in the cost ²⁰

$$\begin{cases} \text{Minimize } J(y, v) = \frac{1}{2} \iint_{q_T} \rho_0^2 |v|^2 dx dt \\ \text{Subject to } (y, v) \in \mathcal{C}(y_0, T). \end{cases} \quad (65)$$

where ρ_0 are non-negative continuous weights functions such that $\rho, \rho_0 \in L^\infty(Q_{T-\delta}) \quad \forall \delta > 0$.

$$\min_{\varphi \in \widetilde{W}_{\rho_0, \rho}} \mathcal{J}^*(\varphi) = \frac{1}{2} \iint_{q_T} \rho_0^{-2} |\varphi(x, t)|^2 dx dt + (y_0, \varphi(\cdot, 0))_{L^2(\Omega)}. \quad (66)$$

$$\widetilde{W}_{\rho_0, \rho} = \{\varphi \in \widetilde{\Phi}_{\rho_0, \rho} : \rho^{-1} L^* \varphi = 0 \text{ in } L^2(Q_T)\}$$

²⁰D. Araujo de Souza, AM, [A mixed formulation for the direct approximation of the control of minimal \$L^2\$ -weighted norm for the linear heat equation.](#) (2016) 


L^2 -weighted norm

No contribution of y in the cost ²⁰

$$\begin{cases} \text{Minimize } J(y, v) = \frac{1}{2} \iint_{q_T} \rho_0^2 |v|^2 dx dt \\ \text{Subject to } (y, v) \in \mathcal{C}(y_0, T). \end{cases} \quad (65)$$

where ρ_0 are non-negative continuous weights functions such that $\rho, \rho_0 \in L^\infty(Q_{T-\delta}) \quad \forall \delta > 0$.

$$\begin{aligned} \min_{\varphi \in \widetilde{W}_{\rho_0, \rho}} \hat{J}^*(\varphi) &= \frac{1}{2} \iint_{q_T} \rho_0^{-2} |\varphi(x, t)|^2 dx dt + (y_0, \varphi(\cdot, 0))_{L^2(\Omega)}. \quad (66) \\ \widetilde{W}_{\rho_0, \rho} &= \{\varphi \in \widetilde{\Phi}_{\rho_0, \rho} : \rho^{-1} L^* \varphi = 0 \text{ in } L^2(Q_T)\} \end{aligned}$$

²⁰D. Araujo de Souza, AM, [A mixed formulation for the direct approximation of the control of minimal \$L^2\$ -weighted norm for the linear heat equation.](#) (2016) 

INVERSE PROBLEM FOR HEAT - RECONSTRUCTION OF y FROM y_{q_T}

$\Omega \subset \mathbb{R}^N$ ($N \geq 1$) - $T > 0$, $c \in C^1(\bar{\Omega}, \mathbb{R})$, $d \in L^\infty(Q_T)$, $y_0 \in \mathbf{H}$

$$\begin{cases} Ly := y_t - \nabla \cdot (c \nabla y) + dy = f, & Q_T := \Omega \times (0, T) \\ y = 0, & \Sigma_T := \partial\Omega \times (0, T) \\ y(\cdot, 0) = y_0, & \Omega. \end{cases} \quad (67)$$

► Inverse Problem : **Distributed observation** on $q_T = \omega \times (0, T)$, $\omega \subset \Omega$

$$\begin{cases} X = L^2(q_T), \\ \text{Given } (y_{obs}, f) \in (L^2(q_T), X), \text{ find } y \text{ s.t. } \{(67) \text{ and } y - y_{obs} = 0 \text{ on } q_T\} \end{cases}$$

WELL-KNOWN DIFFICULTY:

$$\left(Ly \in L^2(Q_T), y \in L^2(q_T), y|_{\Sigma_T} = 0 \right) \implies y \in C([\delta, T], H_0^1(\Omega)), \quad \forall \delta > 0$$

Second order mixed formulation as in the previous part

We then define the following extremal problem :

$$\left\{ \begin{array}{l} \text{Minimize } J(y) := \frac{1}{2} \iint_{Q_T} \rho_0^{-2} |y(x, t) - y_{obs}(x, t)|^2 dx dt + r \iint_{Q_T} (\rho^{-1} Ly)^2 dx dt \\ \text{Subject to } y \in \mathcal{W} := \left\{ y \in \mathcal{Y} : \rho^{-1} Ly = 0 \text{ in } L^2(Q_T) \right\} \end{array} \right. \quad (P)$$

with $\rho_0, \rho \in \mathcal{R}$ where $(\rho_\star \in \mathbb{R}_\star^+)$

$$\mathcal{R} := \{w : w \in C(Q_T); w \geq \rho_\star > 0 \text{ in } Q_T; w \in L^\infty(\Omega \times (\delta, T)) \forall \delta > 0\}$$

H_0^1 – L^2 first order formulation

First order formulation involving y and the flux $\mathbf{p} = c(x)\nabla y$.

$$\begin{cases} \mathcal{I}(y, \mathbf{p}) := y_t - \nabla \cdot \mathbf{p} + d y = f, & \mathcal{J}(y, \mathbf{p}) := c(x)\nabla y - \mathbf{p} = \mathbf{0} & \text{in } Q_T, \\ y = 0 & & \text{on } \Sigma_T, \\ y(x, 0) = y_0(x) & & \text{in } \Omega. \end{cases} \quad (68)$$

$$(y_0, f) \in L^2(\Omega) \times L^2(Q_T) \implies p \in \mathbf{L}^2(Q_T), y \in L^2(0, T, H_0^1(\Omega)), y_t \in L^2(0, T, H^{-1}(\Omega))$$

► Inverse Problem : Distributed observation on $q_T = \omega \times (0, T), \omega \subset \Omega$

$$\begin{cases} X = L^2(q_T), \\ \text{Given } (y_{\text{obs}}, f) \in (L^2(q_T), X), \text{ find } (y, \mathbf{p}) \text{ s.t. } \{(68) \text{ and } y - y_{\text{obs}} = 0 \text{ on } q_T\} \end{cases}$$

Key tool to set up the variational method: A global Carleman estimate obtained from [Immanuvilov, Puel, Yamamoto, 2010]: $\exists C > 0$

$$\|\rho_{p,0}^{-1} y\|_{L^2(Q_T)}^2 \leq C \left(\|\rho_p^{-1} \mathcal{J}(y, \mathbf{p})\|_{L^2(Q_T)}^2 + \|\rho_{p,2}^{-1} \mathcal{I}(y, \mathbf{p})\|_{L^2(Q_T)}^2 + \|\rho_{p,0}^{-1} y\|_{L^2(q_T)}^2 \right),$$

$H_0^1 - L^2$ first order formulation

First order formulation involving y and the flux $\mathbf{p} = c(x)\nabla y$.

$$\begin{cases} \mathcal{I}(y, \mathbf{p}) := y_t - \nabla \cdot \mathbf{p} + dy = f, & \mathcal{J}(y, \mathbf{p}) := c(x)\nabla y - \mathbf{p} = \mathbf{0} & \text{in } Q_T, \\ y = 0 & & \text{on } \Sigma_T, \\ y(x, 0) = y_0(x) & & \text{in } \Omega. \end{cases} \quad (68)$$

$$(y_0, f) \in L^2(\Omega) \times L^2(Q_T) \implies p \in \mathbf{L}^2(Q_T), y \in L^2(0, T, H_0^1(\Omega)), y_t \in L^2(0, T, H^{-1}(\Omega))$$

► Inverse Problem : **Distributed observation** on $q_T = \omega \times (0, T), \omega \subset \Omega$

$$\begin{cases} X = L^2(q_T), \\ \text{Given } (y_{obs}, f) \in (L^2(q_T), X), \text{ find } (y, \mathbf{p}) \text{ s.t. } \{(68) \text{ and } y - y_{obs} = 0 \text{ on } q_T\} \end{cases}$$

Key tool to set up the variational method: A global Carleman estimate obtained from [Immanuvilov, Puel, Yamamoto, 2010]: $\exists C > 0$

$$\|\rho_{p,0}^{-1}y\|_{L^2(Q_T)}^2 \leq C \left(\|\rho_p^{-1}\mathcal{J}(y, \mathbf{p})\|_{L^2(Q_T)}^2 + \|\rho_{p,2}^{-1}\mathcal{I}(y, \mathbf{p})\|_{L^2(Q_T)}^2 + \|\rho_{p,0}^{-1}y\|_{L^2(q_T)}^2 \right),$$

$H_0^1 - L^2$ first order formulation

First order formulation involving y and the flux $\mathbf{p} = c(x)\nabla y$.

$$\begin{cases} \mathcal{I}(y, \mathbf{p}) := y_t - \nabla \cdot \mathbf{p} + d y = f, & \mathcal{J}(y, \mathbf{p}) := c(x)\nabla y - \mathbf{p} = \mathbf{0} & \text{in } Q_T, \\ y = 0 & & \text{on } \Sigma_T, \\ y(x, 0) = y_0(x) & & \text{in } \Omega. \end{cases} \quad (68)$$

$$(y_0, f) \in L^2(\Omega) \times L^2(Q_T) \implies p \in \mathbf{L}^2(Q_T), y \in L^2(0, T, H_0^1(\Omega)), y_t \in L^2(0, T, H^{-1}(\Omega))$$

► Inverse Problem : **Distributed observation** on $q_T = \omega \times (0, T), \omega \subset \Omega$

$$\begin{cases} X = L^2(q_T), \\ \text{Given } (y_{obs}, f) \in (L^2(q_T), X), \text{ find } (y, \mathbf{p}) \text{ s.t. } \{(68) \text{ and } y - y_{obs} = 0 \text{ on } q_T\} \end{cases}$$

Key tool to set up the variational method: A global Carleman estimate obtained from [Immanuvilov, Puel, Yamamoto, 2010]: $\exists C > 0$

$$\|\rho_{p,0}^{-1} y\|_{L^2(Q_T)}^2 \leq C \left(\|\rho_p^{-1} \mathcal{J}(y, \mathbf{p})\|_{L^2(Q_T)}^2 + \|\rho_{p,2}^{-1} \mathcal{I}(y, \mathbf{p})\|_{L^2(Q_T)}^2 + \|\rho_{p,0}^{-1} y\|_{L^2(q_T)}^2 \right),$$

H_0^1 – L^2 first order formulation

First order formulation involving y and the flux $\mathbf{p} = c(x)\nabla y$.

$$\begin{cases} \mathcal{I}(y, \mathbf{p}) := y_t - \nabla \cdot \mathbf{p} + d y = f, & \mathcal{J}(y, \mathbf{p}) := c(x)\nabla y - \mathbf{p} = \mathbf{0} & \text{in } Q_T, \\ y = 0 & & \text{on } \Sigma_T, \\ y(x, 0) = y_0(x) & & \text{in } \Omega. \end{cases} \quad (68)$$

$$(y_0, f) \in L^2(\Omega) \times L^2(Q_T) \implies p \in \mathbf{L}^2(Q_T), y \in L^2(0, T, H_0^1(\Omega)), y_t \in L^2(0, T, H^{-1}(\Omega))$$

► Inverse Problem : **Distributed observation** on $q_T = \omega \times (0, T), \omega \subset \Omega$

$$\begin{cases} X = L^2(q_T), \\ \text{Given } (y_{obs}, f) \in (L^2(q_T), X), \text{ find } (y, \mathbf{p}) \text{ s.t. } \{(68) \text{ and } y - y_{obs} = 0 \text{ on } q_T\} \end{cases}$$


Key tool to set up the variational method: A global Carleman estimate obtained from [Immanuvilov, Puel, Yamamoto, 2010]: $\exists C > 0$

$$\|\rho_{p,0}^{-1} y\|_{L^2(Q_T)}^2 \leq C \left(\|\rho_p^{-1} \mathcal{J}(y, \mathbf{p})\|_{L^2(Q_T)}^2 + \|\rho_{p,2}^{-1} \mathcal{I}(y, \mathbf{p})\|_{L^2(Q_T)}^2 + \|\rho_{p,0}^{-1} y\|_{L^2(q_T)}^2 \right),$$


$N = 1$ - Heat eq. Comparison with the standard method

$$y_0(x) = \sin(\pi x)^{20}, \quad Q_T = (0, 1) \times (0, T), \quad q_T = (0.7, 0.8) \times (0, T), \quad T = 1/2$$

$$\min_{y_{0h}} \left(J_h(y_{0h}) + \frac{h^2}{2} \|y_{0h}\|_{L^2(\Omega)}^2 \right) \quad \text{vs.} \quad \min_{\lambda_h} J_r^{**}(\lambda_h) \quad \text{over } \Lambda_h \quad (69)$$





$$\frac{\|y - y_h\|_{L^2(Q_T)}}{\|y\|_{L^2(Q_T)}} \approx 5.86 \times 10^{-2},$$



$$\frac{\|y - y_h\|_{L^2(Q_T)}}{\|y\|_{L^2(Q_T)}} \approx 7.70 \times 10^{-2}$$



$N = 1$ - Comparison with the standard method

$$y_0(x) = \sin(\pi x)^{20}, \quad Q_T = (0, 1) \times (0, T), \quad q_T = (0.7, 0.8) \times (0, T), \quad T = 1/2$$



Restriction at $(0, 1) \times \{0\}$

$N = 1$ - Comparison with the standard method



Evolution of the relative residu $\frac{\|g^k\|}{\|g^0\|}$ w.r.t. iterate k

The minimization of J w.r.t. y_0 requires 452 (conjugate gradient) iterates

The minimization of J_{λ}^{**} w.r.t. λ requires 4 (conjugate gradient) iterates !

Summary

THE VARIATIONAL APPROACH CAN BE USED IN THE CONTEXT OF MANY OTHER CONTROLLABLE SYSTEMS FOR WHICH APPROPRIATE CARLEMAN ESTIMATES ARE AVAILABLE.

CONFORMAL TIME-SPACE FINITE ELEMENTS APPROXIMATIONS LEAD TO STRONG CONVERGENCE RESULTS WITH RESPECT TO h .

THIS APPROACH MAY BE APPLIED FOR INVERSE PROBLEMS, OBSERVATION PROBLEMS, RECONSTRUCTION OF DATA,

EASILY COMBINED WITH MESH ADAPTIVITY, POSSIBLY VERY USEFUL IN THE PARABOLIC SITUATION

Ongoing works

- ▶ Extension to sparse controls (L^1 term in the cost)
- ▶ Average controllability ²²
- ▶ Approximation of observability constants C_{obs} (to infer or not observability property).

see The 2016 case of the transient heat equation (Martinez-Frutos et al., Class 20, May 13, etc)

$$\begin{cases} y_t - \Delta y = u(x, t) = 0, & (x, t) \in (0, 1) \times (0, T), \quad y = 0 \text{ on } \partial\Omega \times (0, T) \\ y(x, 0) = y_0(x), \quad x \in \Omega \\ y(x, T) = y_1(x) \in L^2(\Omega) \end{cases} \quad (20)$$

Extension of the L^2 case to L^1 (see [20]) and L^1 (see [21]) and of L^2 (see [22]).

The L^1 case is done by PD (see [23] and [24]).

$$\begin{cases} y_t - \Delta y - (X + \gamma) \nabla y = 0, & (x, t) \in \Omega \times (0, T), \quad \Omega \subset \mathbb{R}^d \\ y = 0 \text{ on } \partial\Omega \times (0, T) \end{cases} \quad (21)$$

Extension of the L^2 case to L^1 (see [25] and [26]).

²²Martinez-Frutos, Kessler, M, Periago, Robust optimal Robin boundary control for the transient heat equation with random input data, (2016). 

Ongoing works

- ▶ Extension to sparse controls (L^1 term in the cost)
- ▶ Average controllability ²²
- ▶ Approximation of observability constants C_{obs} (to infer or not observability property).

see The 2016 case of the transient heat equation (Martinez-Frutos et al., Class 20, May 13, etc)

$$\begin{cases} y_t - \Delta y = u(x, t) = 0, & (x, t) \in (0, 1) \times (0, T), \quad y = 0 \text{ on } \partial\Omega \times (0, T) \\ y(x, 0) = y_0(x), \quad x \in \Omega \\ y(x, T) = y_1(x) \in L^2(\Omega) \end{cases} \quad (20)$$

Extension of the L^2 cost $J(u) = \int_0^T \int_{\Omega} |u(x, t)|^2 dx dt$ and $T \in [0, \infty)$ and $y_0, y_1 \in L^2(\Omega)$

The associated adjoint PDE is $y_t - \Delta y = 0$ with $y_0 = 0$

$$\begin{cases} y_t - \Delta y - (X + \lambda) \nabla \cdot \nabla y = 0, & (x, t) \in \Omega \times (0, T), \quad \Omega \subset \mathbb{R}^d \\ y = 0 \text{ on } \partial\Omega \times (0, T) \\ y(x, T) = y_1(x) \in L^2(\Omega) \end{cases}$$

Estimation of the $C_{obs}(T, \lambda) \approx \frac{1}{\lambda} \|\mathbb{1}_{\Omega}\|_{L^2(\Omega)}^2 \approx \lambda^{-1} |\Omega|$

²²Martinez-Frutos, Kessler, M, Periago, Robust optimal Robin boundary control for the transient heat equation with random input data, (2016). 

Ongoing works

- ▶ Extension to sparse controls (L^1 term in the cost)
- ▶ **Average controllability** ²²
- ▶ Approximation of observability constants C_{obs} (to infer or not observability property).

▶ The L^1 case of the boundary control problem (Luenberger, 1997, Chap. 10, Sec. 2)

$$\begin{aligned} \min_{u \in L^1(\Sigma)} & \int_0^T \int_{\Sigma} |u| \, d\sigma \, dt \\ \text{s.t.} & \quad \begin{cases} \partial_t \theta - \Delta \theta = 0 & \text{in } \Omega \times (0, T) \\ \theta = 0 & \text{on } \Gamma \times (0, T) \\ \theta = \theta_0 & \text{in } \Omega \end{cases} \\ & \quad \theta(T) \geq \theta^* \end{aligned}$$

▶ Approximation of the observability constant C_{obs} (see [10])

▶ Approximation of the observability constant C_{obs} (see [10])

▶ Approximation of the observability constant C_{obs} (see [10])

▶ Approximation of the observability constant C_{obs} (see [10])

▶ Approximation of the observability constant C_{obs} (see [10])

²²Martinez-Frutos, Kessler, M, Periago, Robust optimal Robin boundary control for the transient heat equation with random input data, (2016).

Ongoing works

- ▶ Extension to sparse controls (L^1 term in the cost)
- ▶ **Average controllability** ²²
- ▶ Approximation of observability constants C_{obs} (to infer or not observability property).

▶ The L^1 case of the boundary control problem is addressed in [Martinez-Frutos, Class 20, 2016, 13, 142](#).

$$\begin{aligned} \min_{u \in L^1(\Sigma)} & \int_{\Omega} |u| \, dx + \int_{\Omega} |y - y_d| \, dx \\ \text{s.t.} & \quad \begin{cases} \partial_t y - \Delta y = 0 & \text{in } \Omega \times (0, T) \\ y = 0 & \text{on } \Gamma \times (0, T) \\ y = 0 & \text{at } t = 0 \end{cases} \end{aligned}$$

▶ [Martinez-Frutos, Kessler, Periago, SIAM J. Control Optim. 54\(2\) \(2016\)](#)

▶ [Martinez-Frutos, Kessler, Periago, SIAM J. Control Optim. 54\(2\) \(2016\)](#)

▶ [Martinez-Frutos, Kessler, Periago, SIAM J. Control Optim. 54\(2\) \(2016\)](#)

▶ [Martinez-Frutos, Kessler, Periago, SIAM J. Control Optim. 54\(2\) \(2016\)](#)

▶ [Martinez-Frutos, Kessler, Periago, SIAM J. Control Optim. 54\(2\) \(2016\)](#)

²²Martinez-Frutos, Kessler, M, Periago, Robust optimal Robin boundary control for the transient heat equation with random input data, (2016).

Ongoing works

- ▶ Extension to sparse controls (L^1 term in the cost)
- ▶ Average controllability²²
- ▶ Approximation of observability constants C_{obs} (to infer or not observability property).
 - ▶ The stiff case of the transport-diffusion equation [Guerrero-Coron 06, Glass 09, Lissy'13, etc]

$$\begin{cases} y_t - \epsilon y_{xx} + My_x = 0, & Q_T := (0, 1) \times (0, T), \quad M \in \{-1, 1\} \\ y(0, t) = v_\epsilon(t), y(1, t) = 0, \\ y(x, 0) = y_0(x) \in L^2(0, 1) \end{cases} \quad (70)$$

Estimation of $\lim_{\epsilon \rightarrow 0^+} C_{obs}(\epsilon, T, M)$ w.r.t $T \in [|M|^{-1}, \infty)$ and $M \in \{-1, 1\}$!?

- ▶ The elasticity system in 2D (controllable with 2 controls)

$$\begin{cases} \mathbf{y}_{tt} - \mu \Delta \mathbf{y} - (\lambda + \mu) \nabla \operatorname{div} \mathbf{y} = 0, & (\mathbf{x}, t) \in \Omega \times (0, T), \quad \Omega \subset \mathbb{R}^2, \\ \mathbf{y} = \mathbf{f} = (f_1, f_2) \mathbf{1}_{\Gamma_0}, & (\mathbf{x}, t) \in \partial\Omega \times (0, T). \end{cases}$$

Estimation of $\lim C_{obs}(f_1, f_2)$ as $\|f_1\|_{L^2(\Gamma_0 \times (0, T))} \rightarrow 0^+$!?

²²Martinez-Frutos, Kessler, M, Periago, Robust optimal Robin boundary control for the transient heat equation with random input data, (2016).

Ongoing works

- ▶ Extension to sparse controls (L^1 term in the cost)
- ▶ Average controllability²²
- ▶ Approximation of observability constants C_{obs} (to infer or not observability property).
 - ▶ The stiff case of the transport-diffusion equation [Guerrero-Coron 06, Glass 09, Lissy'13, etc]

$$\begin{cases} y_t - \epsilon y_{xx} + My_x = 0, & Q_T := (0, 1) \times (0, T), \quad M \in \{-1, 1\} \\ y(0, t) = v_\epsilon(t), y(1, t) = 0, \\ y(x, 0) = y_0(x) \in L^2(0, 1) \end{cases} \quad (70)$$

Estimation of $\lim_{\epsilon \rightarrow 0^+} C_{obs}(\epsilon, T, M)$ w.r.t $T \in [|M|^{-1}, \infty)$ and $M \in \{-1, 1\}$!?

- ▶ The elasticity system in 2D (controllable with 2 controls)

$$\begin{cases} \mathbf{y}_{tt} - \mu \Delta \mathbf{y} - (\lambda + \mu) \nabla \operatorname{div} \mathbf{y} = 0, & (\mathbf{x}, t) \in \Omega \times (0, T), \quad \Omega \subset \mathbb{R}^2, \\ \mathbf{y} = \mathbf{f} = (f_1, f_2) \mathbf{1}_{\Gamma_0}, & (\mathbf{x}, t) \in \partial\Omega \times (0, T). \end{cases}$$

Estimation of $\lim C_{obs}(f_1, f_2)$ as $\|f_1\|_{L^2(\Gamma_0 \times (0, T))} \rightarrow 0^+$!?

²²Martinez-Frutos, Kessler, M, Periago, Robust optimal Robin boundary control for the transient heat equation with random input data, (2016).

Ongoing works

- ▶ Extension to sparse controls (L^1 term in the cost)
- ▶ Average controllability²²
- ▶ Approximation of observability constants C_{obs} (to infer or not observability property).
 - ▶ The stiff case of the transport-diffusion equation [Guerrero-Coron 06, Glass 09, Lissy'13, etc]

$$\begin{cases} y_t - \epsilon y_{xx} + My_x = 0, & Q_T := (0, 1) \times (0, T), \quad M \in \{-1, 1\} \\ y(0, t) = v_\epsilon(t), y(1, t) = 0, \\ y(x, 0) = y_0(x) \in L^2(0, 1) \end{cases} \quad (70)$$

Estimation of $\lim_{\epsilon \rightarrow 0^+} C_{obs}(\epsilon, T, M)$ w.r.t $T \in [|M|^{-1}, \infty)$ and $M \in \{-1, 1\}$!?

- ▶ The elasticity system in 2D (controllable with 2 controls)

$$\begin{cases} \mathbf{y}_{tt} - \mu \Delta \mathbf{y} - (\lambda + \mu) \nabla \operatorname{div} \mathbf{y} = 0, & (\mathbf{x}, t) \in \Omega \times (0, T), \quad \Omega \subset \mathbb{R}^2, \\ \mathbf{y} = \mathbf{f} = (f_1, f_2) \mathbf{1}_{\Gamma_0}, & (\mathbf{x}, t) \in \partial\Omega \times (0, T). \end{cases}$$

Estimation of $\lim C_{obs}(f_1, f_2)$ as $\|f_1\|_{L^2(\Gamma_0 \times (0, T))} \rightarrow 0^+$!?

²²Martinez-Frutos, Kessler, M, Periago, Robust optimal Robin boundary control for the transient heat equation with random input data, (2016).

The End

NADA MAS !

ENRIQUE, GRACIAS POR TODO Y FELIZ CUMPLEAÑOS !!!!

THANK YOU VERY MUCH FOR YOUR ATTENTION