

UNIFORM STABILIZATION OF A NUMERICAL APPROXIMATION OF A LOCALLY DAMPED WAVE EQUATION *

ARNAUD MÜNCH¹ AND ADEMIR FERNANDO PAZOTO²

Abstract. This work is devoted to the analysis of a viscous finite-difference space semi-discretization of a locally damped wave equation in a regular 2-D domain. The damping term is supported in a suitable subset of the domain, so that the energy of solutions of the damped continuous wave equation decays exponentially to zero as time goes to infinity. Using discrete multiplier techniques, we prove that adding a suitable vanishing numerical viscosity term leads to a uniform (with respect to the mesh size) exponential decay of the energy for the solutions of the numerical scheme. The numerical viscosity term damps out the high frequency numerical spurious oscillations while the convergence of the scheme towards the original damped wave equation is kept, which guarantees that the low frequencies are damped correctly. Numerical experiments are presented and confirm these theoretical results. These results extend those by Tcheugoué-Tébou and Zuazua in [23] where the 1-D case was addressed as well the square domain in 2-D. The methods and results in this paper extend to smooth domains in any space dimension.

Résumé. Ce travail concerne l'analyse d'un schéma aux différences finies semi-discret en espace approchant une équation des ondes amorties localement et posée sur un domaine 2-D. Le terme d'amortissement est défini sur un sous-ensemble du domaine de façon à ce que l'énergie associée au système continu décroisse exponentiellement vers zéro lorsque le temps tends vers l'infini. En utilisant une technique des multiplicateurs discrètes, on montre que l'ajout d'un terme de viscosité permet de garantir la décroissance exponentielle uniformément (vis à vis de la taille du maillage) de l'énergie associée au système semi-discret. Le terme de viscosité permet de capter les solutions numériques haute fréquence tout en préservant la convergence du schéma vers le système continu. Des essais numériques sont présentés et confirment ces résultats théoriques. Ces résultats étendent ceux de Tcheugoué-Tébou and Zuazua [23] aux cas de domaines 2-D réguliers quelconques.

1991 Mathematics Subject Classification. 65M06.

Keywords and phrases: wave equation, stabilization, finite difference, viscous terms

** This work was done while the authors were visiting the Department of Mathematics of the Universidad Autónoma de Madrid. The first author was supported by Grants HPRN-CT-2002-00284 of the European program New materials, adaptive systems and their nonlinearities: modelling, control and numerical simulation. The second author was supported by Grants SB2001-0158 of the MECD (Spain) and BFM 2002-03345 of the MCYT (Spain).*

¹ Departamento de Matemáticas, Universidad Autónoma de Madrid, 28049, Madrid, Spain, email:arnaud.munch@uam.es

² Instituto de Matemática, Universidade Federal do Rio de Janeiro, 21940-970, Rio de Janeiro, Brasil, email:ademir@acd.ufrj.br

INTRODUCTION AND SETTING OF THE PROBLEM

Let us consider the following damped wave equation defined on $\Omega \times [0, \infty)$, Ω being a bounded domain in \mathbb{R}^2 of class C^2 :

$$\begin{cases} y'' - \Delta y + a(\mathbf{x})y' = 0 & \text{in } \Omega \times (0, \infty), \\ y = 0 & \text{on } \Gamma \equiv \partial\Omega \times (0, \infty), \\ y(\mathbf{x}, 0) = y_0(\mathbf{x}) & \text{in } \Omega, \\ y'(\mathbf{x}, 0) = y_1(\mathbf{x}) & \text{in } \Omega, \end{cases} \quad (1)$$

where the symbol $'$ designates the partial differentiation with respect to time. The damping potential $a = a(\mathbf{x})$ is assumed to be a bounded nonnegative function defined on Ω . In addition, we suppose that there exists a positive constant α_0 such that

$$a(\mathbf{x}) \geq \alpha_0 \text{ a.e. } \mathbf{x} \in \omega \text{ where } \omega \text{ is a nonempty open subset of } \Omega. \quad (2)$$

System (1) is well-posed in the space $H_0^1(\Omega) \times L^2(\Omega)$. Indeed, given $\{y_0, y_1\} \in H_0^1(\Omega) \times L^2(\Omega)$, there exists a unique solution of (1) with (see [15])

$$y \in \mathcal{C}([0, \infty); H_0^1(\Omega)) \cap \mathcal{C}^1([0, \infty); L^2(\Omega)). \quad (3)$$

The energy of the system (1) is given by

$$E(t) = \frac{1}{2} \int_{\Omega} \{|y'(\mathbf{x}, t)|^2 + |\nabla y(\mathbf{x}, t)|^2\} dx, \quad \forall t \geq 0. \quad (4)$$

Moreover, the following dissipation law holds

$$E'(t) = - \int_{\Omega} a(\mathbf{x})|y'(\mathbf{x}, t)|^2 dx, \quad \forall t \geq 0. \quad (5)$$

Consequently, E is a non increasing function of the time variable t and therefore a rate of decay is expected.

When $a(\mathbf{x}) \geq \alpha_0 > 0$ a.e. in Ω , it is straightforward to see that the energy decays uniformly exponentially as t goes to infinity: there exist two positive constants C and α such that

$$E(t) \leq C e^{-\alpha t} E(0), \quad \forall t \geq 0. \quad (6)$$

The problem of stabilization when the damping is effective only on a subset of Ω is much more subtle. This problem has been extensively investigated in the context of wave equations and there is a large literature on the subject. For instance, Dafermos [6], Haraux [9] and Slemrod [24] used La Salle's invariance principle to obtain asymptotic stability properties. More recently, assuming that the dissipation is effective in a suitable subset of the domain where the equation holds, decay rates have been obtained (see for instance Nakao [18] and Zuazua [25]). In particular, when ω is a neighborhood of the subset of the boundary $\partial\Omega$ as follows

$$\omega \text{ is the intersection of } \Omega \text{ and a neighborhood of } \Gamma_0, \quad (7)$$

where

$$\Gamma_0 = \{\mathbf{x} \in \partial\Omega; \mathbf{x} \cdot \boldsymbol{\nu} > 0\}, \quad (8)$$

and $\boldsymbol{\nu}$ the unit normal pointing into the exterior of Ω then, the uniform exponential decay (6) holds ([8], [16], [22], [25]).

In view of (5), the problem of the exponential decay of E can be stated in the following equivalent form: is it possible to find $T > 0$ and $C > 0$ such that

$$E(0) \leq C \int_0^T \int_{\Omega} a(\mathbf{x}) |y'(\mathbf{x}, t)|^2 dx dt \quad (9)$$

holds for every finite energy solution of (1)? More precisely, (9) combined with (5) and the semigroup property, allows to obtain the exponential decay property (6). However, we emphasize that, in the general multi-dimensional setting, inequalities of the form (9) are valid if and only if the subset ω where the damping is effective satisfies a suitable Geometric Control Condition (GCC) (see [3]). Roughly speaking, the GCC requires that every ray of geometric optics reaches the region in which the damping mechanism is effective in a uniform time, a property that holds in the particular case where ω is given by (7-8).

We address in this work the numerical approximation of this problem which is by now well-know to be extremely sensitive: for instance, the most classical finite-difference semi-discrete numerical approximation of the first equation of (1)

$$y''_{j,k}(t) - \frac{y_{j+1,k}(t) - 2y_{j,k}(t) + y_{j-1,k}(t)}{h_1^2} - \frac{y_{j,k+1}(t) - 2y_{j,k}(t) + y_{j,k-1}(t)}{h_2^2} + a_{j,k} y'_{j,k}(t) = 0 \quad (10)$$

leads to a semi-discrete observability equality similar to (9) where the constant C_{h_1, h_2} depending now on the mesh size h_1, h_2 (defined in details in section 2) may blows-up as h_1, h_2 go to zero, depending mainly on the regularity of the initial conditions. Therefore, the exponential decay rate α_{h_1, h_2} of the discretized energy is not uniform with respect to the discretization parameters. This numerical pathology is independent of the consistence and stability properties of the scheme. This is due to the fact that the interaction of waves with a numerical mesh produces dispersion phenomena and spurious high frequency oscillations. Because of this nonphysical interaction the velocity of propagation of such numerical waves, the so called group velocity may converge to zero when the wavelength of solutions is of order of the mesh size $O(h_1, h_2)$ leading to a time of observability of order $1/O(h_1, h_2)$. Such a pathology was suggested by numerical simulations in a preliminary work in [2]. According to the close links between stabilization and exact controllability [12], same bad behaviors appear for the numerical schemes devoted to exact controllability problems (we refer the reader to [27] and the references therein for an extended survey on this issue).

The problem of numerical approximation and stabilization has been addressed in [23] in 1-D and 2-D in the interval and the unit square domain, respectively. It was shown that an estimate similar to (9) is valid for the “finite-difference space semi-discretization” of (1), which is uniform with respect to the mesh size, provided a suitable viscosity term is added. More precisely, the authors replaced the equation (10) by the following one:

$$y''_{j,k}(t) - \frac{y_{j+1,k}(t) - 2y_{j,k}(t) + y_{j-1,k}(t)}{h_1^2} - \frac{y_{j,k+1}(t) - 2y_{j,k}(t) + y_{j,k-1}(t)}{h_2^2} + a_{j,k} y'_{j,k}(t) - h_1^2 \left(\frac{y'_{j+1,k}(t) - 2y'_{j,k}(t) + y'_{j-1,k}(t)}{h_1^2} \right) - h_2^2 \left(\frac{y'_{j,k+1}(t) - 2y'_{j,k}(t) + y'_{j,k-1}(t)}{h_2^2} \right) = 0 \quad (11)$$

In this work, we treat an important case left open in [23]. It is the one in which the model under consideration is posed in a regular 2-D bounded domain. This introduces important additional technical difficulties related with the analysis near the boundary. In this respect, we introduce a sequence of discretized domains $(\Omega_{h_1, h_2})_{h_1, h_2}$ converging to Ω (in a sense that will be made precise later) when the mesh parameters h_1 and h_2 go to zero (see Figure 1). Then, we associate to the first equation of (1) the finite-difference semi-discretization equation (11) where $y_{j,k}(t)$ denotes the approximation of $y(\cdot, t)$ at the node (jh_1, kh_2) of Ω_{h_1, h_2} . Following the methodology developed in [23], we show the efficiency of this scheme : we first prove that the energy associated with this numerical scheme decays exponentially, with a rate independent of the mesh size (see Theorem 2.1,

section 2). To obtain the result we combine discrete multiplier methods introduced in [11] and compactness-uniqueness arguments. Let us note that a discrete version of multipliers was also developed in [11, 26] to address the issue of boundary observability in 1-D and the 2-D square domain. Then, we prove that the solution of the discretized system defined on Ω_{h_1, h_2} converges to the solution of (1) defined on Ω (see theorem 3.1, section 3). Finally, we present some numerical experiments in order to confirm these theoretical results (section 4).




FIGURE 1. General domain Ω and two different discretizations Ω_{h_1, h_2} .

1. DISCRETIZATION OF THE DOMAIN Ω AND THE FINITE DIFFERENT SCHEME

1.1. Discretization of the domain Ω

We introduce in this paragraph a spatial discretization of the domain Ω . Without loss of generality, we assume that the domain Ω is included in the unit square box: $\Omega \subseteq B \equiv]0, 1[\times]0, 1[$. For non negatives integer J and K , we then set

$$h_1 = \frac{1}{J+1}, \quad h_2 = \frac{1}{K+1}, \quad (12)$$

and consider the two following uniform subdivisions of the interval $[0, 1]$

$$\begin{cases} 0 = x_{1,0} < x_{1,1} < \dots < x_{1,J} < x_{1,J+1} = 1, \\ 0 = x_{2,0} < x_{2,1} < \dots < x_{2,K} < x_{2,K+1} = 1, \end{cases} \quad (13)$$

where $x_{1,j} = jh_1$ and $x_{2,k} = kh_2$ for $j \in [0, J+1]$ and $k \in [0, K+1]$. We associate to the grid $G = (x_{1,j}, x_{2,k})_{j,k}$ a discretization \mathbf{B}_{h_1, h_2} of B as follow:

$$\mathbf{B}_{h_1, h_2} = \cup_{j \in [0, J], k \in [0, K]} B_{j,k},$$

where

$$B_{j,k} =]x_{1,j}, x_{1,j+1}[\times]x_{2,k}, x_{2,k+1}[\quad (14)$$

represents the cell of area $h_1 h_2$. Naturally, in the simple case of the unit square, the discretization is exact in the sense that $\mathbf{B}_{h_1, h_2} = B$. We then define Ω_{h_1, h_2} as the union of all the cells $B_{j,k}$ strictly included in Ω :

$$\Omega_{h_1, h_2} = \cup_{j,k} B_{j,k}, \quad \forall j \in [0, J], k \in [0, K] \quad \text{such that} \quad B_{j,k} \subset \Omega \quad (15)$$

implying in particular the inclusion $\Omega_{h_1, h_2} \subset \Omega$ for all h_1 and h_2 . More precisely, we can write Ω_{h_1, h_2} in the following way

$$\Omega_{h_1, h_2} = \cup_{j=0, J} \cup_{k \in \mathcal{I}_j} B_{j,k}, \quad (16)$$

where, for j from 0 to J , \mathcal{I}_j designs the subset of $[0, K]$ defined by

$$\mathcal{I}_j = \left\{ k \in [0, K] : B_{j,k} \subset \Omega_{h_1, h_2} \right\}, \quad j \in [0, J]. \quad (17)$$

Therefore, $k \in \mathcal{I}_j \iff B_{j,k} \in \Omega_{h_1, h_2}$. Now, in order to state the next definition, we extend \mathcal{I}_j to $j = -1$ and $j = J + 1$ putting $\mathcal{I}_{-1} = \emptyset$ and $\mathcal{I}_{J+1} = \emptyset$. Then, proceeding as before, we define the approximation ω_{h_1, h_2} of the open subset ω as the union of all the cells $B_{j,k}$ included in ω .

Let us now describe the boundary of Ω_{h_1, h_2} which plays a crucial role in the context of stabilization. According to (16), the boundary is composed of segments $\tilde{I}(j, k)$ and $\hat{I}(j, k)$ parallel to the axis Ox_1 or to Ox_2 , respectively:

$$\tilde{I}(j, k) = \{j\} \times [k, k + 1] \quad \text{and} \quad \hat{I}(j, k) = [j, j + 1] \times \{k\}, \quad (18)$$

for $j \in [0, J]$ and $k \in [0, K]$. Thus,

$$\tilde{I}(j, k) \subset \partial\Omega_{h_1, h_2} \quad \text{if} \quad (k \in \mathcal{I}_j \text{ and } k \notin \mathcal{I}_{j-1}) \quad \text{or} \quad (k \notin \mathcal{I}_j \text{ and } k \in \mathcal{I}_{j-1}) \quad (19)$$

and

$$\hat{I}(j, k) \subset \partial\Omega_{h_1, h_2} \quad \text{if} \quad (k \in \mathcal{I}_j \text{ and } k - 1 \notin \mathcal{I}_j) \quad \text{or} \quad (k \notin \mathcal{I}_j \text{ and } k - 1 \in \mathcal{I}_j). \quad (20)$$

In the sequel, we will need a finer distinction of the elements of the boundary of Ω_{h_1, h_2} :

Definition 1.1 (Boundary of Ω_{h_1, h_2}). We introduce the following subsets of $\mathbb{N} \times \mathbb{N}$:

- $\tilde{I}^+ = \{(j, k) \text{ such that } k \in \mathcal{I}_{j-1}, k \notin \mathcal{I}_j, j \in [1, J + 1]\}$;
- $\tilde{I}^- = \{(j, k) \text{ such that } k \in \mathcal{I}_j, k \notin \mathcal{I}_{j-1}, j \in [0, J]\}$;
- $\hat{I}^+ = \{(j, k) \text{ such that } k - 1 \in \mathcal{I}_j, k \notin \mathcal{I}_j, j \in [0, J]\}$;
- $\hat{I}^- = \{(j, k) \text{ such that } k - 1 \notin \mathcal{I}_j, k \in \mathcal{I}_j, j \in [0, J]\}$

describing the nodes of $\partial\Omega_{h_1, h_2}$.

One element of each of these four subsets is described in Figure 2.

Remark 1.2. In the sequel, in order to simplify the notation, we will identify each of these four subsets by the corresponding union of segments of the boundary: for instance, we identify \tilde{I}^+ with $\{\mathbf{x} \in \tilde{I}(j, k), \forall (j, k) \in \tilde{I}^+\}$.

To illustrate the parametrization above we write it explicitly for the simple polygonal domain Ω_{h_1, h_2} described in the Figure 3 where $J = 8$ and $K = 5$. The objects corresponding to this example are summarized on the table 1 for which the subsets \tilde{I}^+ and \hat{I}^+ are

$$\begin{cases} \tilde{I}^+ = \{(3, 0), (3, 1), (3, 5), (7, 3), (9, 0), (9, 1), (9, 2), (9, 4), (9, 5)\}, \\ \hat{I}^+ = \{(0, 4), (3, 5), (4, 5), (7, 3), (8, 3), (9, 3), (1, 6), (2, 6), (5, 6), (6, 6), (7, 6), (8, 6)\}. \end{cases}$$





FIGURE 2. Description of the four types of segments of the boundary: $\tilde{I}^+(j, k)$ (left up), $\tilde{I}^-(j, k)$ (right up), $\hat{I}^+(j, k)$ (left down) and $\hat{I}^-(j, k)$ (right down).

j	$n(j)$	\mathcal{I}_j
0	1	{1, 2, 3}
1	1	{0, 1, 2, 3, 4, 5}
2	1	{0, 1, 2, 3, 4, 5}
3	1	{2, 3, 4}
4	1	{2, 3, 4}
5	1	{0, 1, 2, 3, 4, 5}
6	1	{0, 1, 2, 3, 4, 5}
7	2	{0, 1, 2} \cup {4, 5}
8	2	{0, 1, 2} \cup {4, 5}

TABLE 1. Indexes corresponding to the parametrization of Ω_{h_1, h_2} in Figure 3.

1.2. Finite difference scheme

Thanks to the parametrization described above, we are now in the position to introduce and study a discretized version of the system (1). We consider the following semi-discrete finite difference scheme resulting from the usual 5-points approximation (of order 2) in space of the laplacian:

FIGURE 3. Polygonal domain Ω_{h_1, h_2} with $J = 8$ and $K = 5$.

$$\left\{ \begin{array}{l} y''_{j,k} - \frac{y_{j+1,k} - 2y_{j,k} + y_{j-1,k}}{h_1^2} - \frac{y_{j,k+1} - 2y_{j,k} + y_{j,k-1}}{h_2^2} + a_{j,k}y'_{j,k} \\ \quad - h_1^2 \left(\frac{y'_{j+1,k} - 2y'_{j,k} + y'_{j-1,k}}{h_1^2} \right) - h_2^2 \left(\frac{y'_{j,k+1} - 2y'_{j,k} + y'_{j,k-1}}{h_2^2} \right) = 0 \\ \quad \text{for } (j, k) \in \Omega_{h_1, h_2} \text{ and } t \in (0, \infty), \\ y_{j,k}(t) = 0 \text{ for } (j, k) \in \partial\Omega_{h_1, h_2} \text{ and } t \in (0, \infty), \\ y_{j,k}(t) = (y_0)_{j,k}, \quad y'_{j,k}(t) = (y_1)_{j,k} \text{ for } (j, k) \in \Omega_{h_1, h_2} \text{ and } t = 0. \end{array} \right. \quad (21)$$

which differs from the most classical finite-difference scheme by the addition of viscous terms. We shall see below that this numerical approximation scheme converges in the energy space to the continuous damped wave equation (1).

Here, $y_{j,k}$ stands for the approximation of the solution y at the mesh point $(x_{1,j}, x_{2,k})$ and at time $t \in (0, \infty)$ and $(y_0)_{j,k}$ and $(y_1)_{j,k}$ the approximations of the initial data $y_0(\mathbf{x})$ and $y_1(\mathbf{x})$ at time $t = 0$. Furthermore, the notation for $(j, k) \in \Omega_{h_1, h_2}$ means for $j \in [0, J], k \in \mathcal{I}_j$, i.e. for the indexes (j, k) such that the nodes $(x_{1,j}, x_{2,k})$ are in Ω_{h_1, h_2} . In the same manner, the notation for $(j, k) \in \partial\Omega_{h_1, h_2}$ means for $(j, k) \in \tilde{I}^+ \cup \tilde{I}^- \cup \hat{I}^+ \cup \hat{I}^-$. $a_{j,k}$ denotes the approximation of the damping at the mesh point $(x_{1,j}, x_{2,k})$ and we assume that there exists a positive constant α_0 such that

$$a_{j,k} \geq \alpha_0 \quad \forall (j, k) \in \omega_{h_1, h_2}. \quad (22)$$

According to (7), ω_{h_1, h_2} defined in the section 1 is as follow

$$\omega_{h_1, h_2} \text{ is the intersection of } \Omega_{h_1, h_2} \text{ and a neighborhood of } \tilde{I}^+ \cup \hat{I}^+. \quad (23)$$

We can associate to this scheme the following energy:

$$E_{h_1, h_2}(t) = \frac{h_1 h_2}{2} \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \left\{ (y'_{j,k})^2 + \left(\frac{y_{j+1,k} - y_{j,k}}{h_1} \right)^2 + \left(\frac{y_{j,k+1} - y_{j,k}}{h_2} \right)^2 \right\}, \quad (24)$$

which appears to be the semi-discrete version of the energy E in (4) of the solutions of the continuous wave equation. Moreover, multiplying the first equation in (21) by $y'_{j,k}$ and summing over j and k , one can show that the discretized energy E_{h_1, h_2} is a non increasing function of time. More precisely,

$$\begin{aligned} E'_{h_1, h_2}(t) &= -h_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} a_{j,k} (y'_{j,k})^2 \\ &\quad - h_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \left\{ h_1^2 \left(\frac{y'_{j+1,k} - y'_{j,k}}{h_1} \right)^2 + h_2^2 \left(\frac{y'_{j,k+1} - y'_{j,k}}{h_2} \right)^2 \right\}. \end{aligned} \quad (25)$$

which is a semi-discrete version of the energy dissipation law (5).

Furthermore, in order to obtain the exponential decay of solutions of the damped semi-discrete system (21) it is necessary and sufficient to prove the existence of a time $T > 0$ and a constant $C > 0$ independent of h_1 and h_2 such that the following Observability Inequality holds

$$E_{h_1, h_2}(0) \leq C h_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T a_{j,k} (y'_{j,k})^2 dt, \quad (26)$$

for every solution of (21). In the next section, we will prove this inequality using a discrete multiplier method.

We highlight the fact that the constant C in (26) may be not independent of h_1, h_2 for some solutions of the usual scheme without extra viscous terms. More precisely, the usual scheme creates high frequency spurious solutions that propagate very slowly (with group velocity of the order of the step size h_1 and h_2) making the observability property (26) impossible to hold uniformly in h_1 and h_2 . The purpose of the viscous terms is to precisely absorb these high frequency solutions for which the quantities $y'_{j+1,k} - 2y'_{j,k} + y'_{j-1,k}$ and $y'_{j,k+1} - 2y'_{j,k} + y'_{j,k-1}$ are not negligible.

Remark 1.3. The first equation in (21) can be seen as a semi-discrete version of the following viscous wave equation defined on Ω :

$$y'' - \Delta y + a(x)y' - h_1^2 \frac{\partial^2}{\partial x_1^2} y' - h_2^2 \frac{\partial^2}{\partial x_2^2} y' = 0. \quad (27)$$

Therefore, the extra viscous terms introduced are closely connected to the Tychonoff regularization technique [7] and also to the techniques introduced in [13, 17] which consist to add extra corrected terms proportionnal to h_1^2, h_2^2 in order to restore uniform observability property.

In the following, we denote by C a positive constant that may vary from line to line but is independent on h_1 and h_2 .

2. UNIFORM STABILIZATION OF THE ENERGY

This section is devoted to show that the energy associated to (21) decays exponentially, the rate being uniform with respect to the mesh size. Our main result reads as follows:

According to (29) we have

$$\begin{aligned}
E_{h_1, h_2}(0) &\leq Ch_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left[a_{j,k}(y'_{j,k})^2 + h_1^2 \left(\frac{y'_{j+1,k} - y'_{j,k}}{h_1} \right)^2 + h_2^2 \left(\frac{y'_{j,k+1} - y'_{j,k}}{h_2} \right)^2 \right. \\
&\quad \left. + a_{j,k}(z'_{j,k})^2 + h_1^2 \left(\frac{z'_{j+1,k} - z'_{j,k}}{h_1} \right)^2 + h_2^2 \left(\frac{z'_{j,k+1} - z'_{j,k}}{h_2} \right)^2 \right] dt.
\end{aligned} \tag{32}$$

Let us now assume for the moment that the following inequality holds:

$$\begin{aligned}
&\sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left[a_{j,k}(z'_{j,k})^2 + h_1^2 \left(\frac{z'_{j+1,k} - z'_{j,k}}{h_1} \right)^2 + h_2^2 \left(\frac{z'_{j,k+1} - z'_{j,k}}{h_2} \right)^2 \right] dt \\
&\leq C \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left[a_{j,k}(y'_{j,k})^2 + h_1^2 \left(\frac{y'_{j+1,k} - y'_{j,k}}{h_1} \right)^2 + h_2^2 \left(\frac{y'_{j,k+1} - y'_{j,k}}{h_2} \right)^2 \right] dt.
\end{aligned} \tag{33}$$

Combining (32) and (33), it follows that

$$\begin{aligned}
E_{h_1, h_2}(T) &\leq E_{h_1, h_2}(0) \\
&\leq Ch_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left[a_{j,k}(y'_{j,k})^2 + h_1^2 \left(\frac{y'_{j+1,k} - y'_{j,k}}{h_1} \right)^2 + h_2^2 \left(\frac{y'_{j,k+1} - y'_{j,k}}{h_2} \right)^2 \right] dt.
\end{aligned} \tag{34}$$

Now, integrating in $(0, T)$ the energy dissipation law (25) for solutions of (21), it follows that

$$\begin{aligned}
E_{h_1, h_2}(T) - E_{h_1, h_2}(0) \\
&= -h_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left[a_{j,k}(y'_{j,k})^2 + h_1^2 \left(\frac{y'_{j+1,k} - y'_{j,k}}{h_1} \right)^2 + h_2^2 \left(\frac{y'_{j,k+1} - y'_{j,k}}{h_2} \right)^2 \right] dt.
\end{aligned} \tag{35}$$

Combining (34) and (35), we deduce that

$$E_{h_1, h_2}(T) \leq \gamma E_{h_1, h_2}(0) \tag{36}$$

with $\gamma = \frac{C}{C+1}$. Inequality (36) together with the semigroup property imply that

$$E_{h_1, h_2}(t) \leq \frac{1}{\gamma} \exp^{\frac{-|\log(\gamma)|}{T}t} E_{h_1, h_2}(0), \quad \forall t > 0, \quad \text{and } 0 < h_1, h_2 < 1. \tag{37}$$

This yields the desired exponential decay.

To conclude, let us now check (33). Multiplying the equation satisfied by z_{h_1, h_2} by $z'_{j,k}$, we get

$$\begin{aligned}
F'_{h_1, h_2}(t) &= h_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \left(h_1 \left(\frac{y'_{j+1, k} - y'_{j, k}}{h_1} \right) + h_2 \left(\frac{y'_{j, k+1} - y'_{j, k}}{h_2} \right) \right) z'_{j, k} \\
&\quad - h_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \left(h_1 \left(\frac{y'_{j+1, k} - y'_{j, k}}{h_1} \right) z'_{j+1, k} + h_2 \left(\frac{y'_{j, k+1} - y'_{j, k}}{h_2} \right) z'_{j, k+1} \right) \\
&\quad - h_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} a_{j, k} y'_{j, k} z'_{j, k} \\
&\leq F_{h_1, h_2}(t) + C h_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \left(h_1^2 \left(\frac{y'_{j+1, k} - y'_{j, k}}{h_1} \right)^2 + h_2^2 \left(\frac{y'_{j, k+1} - y'_{j, k}}{h_2} \right)^2 + a_{j, k} (y'_{j, k})^2 \right),
\end{aligned} \tag{38}$$

where F_{h_1, h_2} denotes the energy of the solution of system (31).

Integrating (38) in $[0, T]$, taking into account that $F_{h_1, h_2}(0) = 0$, using the Gronwall lemma and the fact that

$$\begin{aligned}
&h_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left(h_1^2 \left(\frac{z_{j+1, k} - z_{j, k}}{h_1} \right)^2 + h_2^2 \left(\frac{z_{j, k+1} - z_{j, k}}{h_2} \right)^2 \right) dt \\
&\quad + h_1 h_2 \sum_{j=1}^J \sum_{k \in \mathcal{I}_j} \int_0^T a_{j, k} (z_{j, k}(t))^2 dt \leq C \int_0^T F_{h_1, h_2}(t) dt,
\end{aligned} \tag{39}$$

inequality (33) immediately follows.

Remark 2.2. In the case where the damping is effective on the whole domain ($\omega = \Omega$), we obtain easily that

$$\sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left[h_1^2 \left(\frac{u'_{j+1, k} - u'_{j, k}}{h_1} \right)^2 + h_2^2 \left(\frac{u'_{j, k+1} - u'_{j, k}}{h_2} \right)^2 \right] dt \leq C \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T a_{j, k} (u'_{j, k})^2 dt \tag{40}$$

implying that the proof of (29) is reduced to the proof of (26). Therefore, in this case, the use of extra viscous terms is no longer necessary.

Step 2: *Energy identity for the conservative discretized equation (30)*

The purpose of this step is to derive an energy identity for the conservative discretized system (30), which is the basic tool for proving an *a priori* asymptotic estimate on the solution of (21). We state the result in the following lemma:

Lemma 2.3. *The semi-discretized energy E_{h_1, h_2} satisfies the following relation:*

$$\begin{aligned}
TE_{h_1, h_2}(0) &= \int_0^T E_{h_1, h_2}(t) dt \\
&= -\frac{h_1 h_2}{2} \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} u'_{j,k} \left(j h_1 \left(\frac{u_{j+1,k} - u_{j-1,k}}{h_1} \right) + \frac{u_{j,k}}{2} \right) \Big|_0^T \\
&\quad - \frac{h_1 h_2}{2} \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} u'_{j,k} \left(k h_2 \left(\frac{u_{j,k+1} - u_{j,k-1}}{h_2} \right) + \frac{u_{j,k}}{2} \right) \Big|_0^T \\
&\quad - \frac{h_2}{2} \sum_{(j,k) \in \tilde{I}^-} j h_1 \int_0^T \left(\frac{u_{j+1,k}}{h_1} \right)^2 dt + \frac{h_2}{2} \sum_{(j,k) \in \tilde{I}^+} j h_1 \int_0^T \left(\frac{u_{j-1,k}}{h_1} \right)^2 dt \\
&\quad - \frac{h_1}{2} \sum_{(j,k) \in \hat{I}^-} k h_2 \int_0^T \left(\frac{u_{j,k+1}}{h_2} \right)^2 dt + \frac{h_1}{2} \sum_{(j,k) \in \hat{I}^+} k h_2 \int_0^T \left(\frac{u_{j,k-1}}{h_2} \right)^2 dt \\
&\quad - \frac{h_1 h_2}{2} \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left\{ \left(\frac{u_{j+1,k} - u_{j,k}}{h_1} \right)^2 + \left(\frac{u_{j,k+1} - u_{j,k}}{h_2} \right)^2 \right\} dt \\
&\quad + \frac{h_1 h_2}{2} \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left\{ \left(\frac{u_{j+1,k+1} - u_{j,k+1}}{h_1} \right) \left(\frac{u_{j+1,k} - u_{j,k}}{h_1} \right) \right\} dt \\
&\quad + \frac{h_1 h_2}{2} \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left\{ \left(\frac{u_{j+1,k+1} - u_{j+1,k}}{h_2} \right) \left(\frac{u_{j,k+1} - u_{j,k}}{h_2} \right) \right\} dt \\
&\quad + \frac{h_1 h_2}{4} \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left\{ h_1^2 \left(\frac{u'_{j+1,k} - u'_{j,k}}{h_1} \right)^2 + h_2^2 \left(\frac{u'_{j,k+1} - u'_{j,k}}{h_2} \right)^2 \right\} dt.
\end{aligned} \tag{41}$$

Remark 2.4. The identity (41) is analogous - on a semi discrete level - to the well-known relation

$$TE(0) = - \int_{\Omega} u_t \left(\mathbf{x} \cdot \nabla u + \frac{u}{2} \right) \Big|_0^T dx + \frac{1}{2} \int_0^T \int_{\partial\Omega} \mathbf{x} \cdot \boldsymbol{\nu} \left| \frac{\partial u}{\partial \nu} \right|^2 d\nu dt \tag{42}$$

where u is solution the continuous conservative wave equation (1) with $a \equiv 0$. This relation may be obtained using the multipliers $\mathbf{x} \cdot \nabla u$ and u (see [12] for instance). More precisely, the two first sums of (41) clearly approximate the first integral of (42). Furthermore, the third and fourth sums approximate the boundary terms of (42). To see this, it is convenient to split the boundary integral into two integrals in the subsets where $\mathbf{x} \cdot \boldsymbol{\nu} > 0$ and $\mathbf{x} \cdot \boldsymbol{\nu} < 0$, respectively. Due the Dirichlet boundary condition, the approximation of, for instance, $\frac{\partial u}{\partial \nu}$ on the point $(x_{1,j}, x_{2,k})$ of \tilde{I}^+ is simply $-\frac{u_{j-1,k}}{h_1}$. Furthermore, $\tilde{I}^+ \cup \hat{I}^+$ coincides with Γ_0 while $\tilde{I}^- \cup \hat{I}^-$ with $\partial\Omega \setminus \Gamma_0$. However, (41) contains some terms (the last four) that do not appear in the continuous case. It is precisely in order to absorb these terms and more specifically the positives ones that adding the numerical viscosity term is needed.

Proof of lemma 2.3 - Let us introduce $M_{j,k}$

$$M_{j,k} = j \left(\frac{u_{j+1,k} - u_{j-1,k}}{2} \right) + k \left(\frac{u_{j,k+1} - u_{j,k-1}}{2} \right), \tag{43}$$

which corresponds to the semi-discrete version of the multiplier $\mathbf{x} \cdot \nabla u$. Following the developments in [23], we multiply the first equation of (30) by $M_{j,k}$, take the sum over j and k , and integrate by parts in $[0, T]$. \square

Step 3: *Absorption of the term at time $t = 0, T$ and the crossed one*

Now we are going to estimate the term at time $t = 0, T$ as well as the crossed terms entering in the right hand side of (41).

Using Young inequality as well as (24), we obtain (see the appendix of [23] for more details concerning the unit square domain)

$$\begin{aligned} & \left| -\frac{h_1 h_2}{2} \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} u'_{j,k} \left(j h_1 \left(\frac{u_{j+1,k} - u_{j-1,k}}{h_1} \right) + \frac{u_{j,k}}{2} \right) \right|_0^T \\ & \quad - \frac{h_1 h_2}{2} \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} u'_{j,k} \left(k h_2 \left(\frac{u_{j,k+1} - u_{j,k-1}}{h_1} \right) + \frac{u_{j,k}}{2} \right) \right|_0^T \\ & \leq \left(2\sqrt{2} + \frac{\sqrt{2}}{4} \max(h_1^2, h_2^2) \right) E_{h_1, h_2}(0). \end{aligned} \quad (44)$$

Also, we obtain

$$\begin{aligned} & -\frac{h_1 h_2}{2} \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left\{ \left(\frac{u_{j+1,k} - u_{j,k}}{h_1^2} \right)^2 + \left(\frac{u_{j,k+1} - u_{j,k}}{h_2^2} \right)^2 \right\} dt \\ & + \frac{h_1 h_2}{2} \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left\{ \left(\frac{u_{j+1,k+1} - u_{j,k+1}}{h_1} \right) \left(\frac{u_{j+1,k} - u_{j,k}}{h_1} \right) \right\} dt \\ & + \frac{h_1 h_2}{2} \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left\{ \left(\frac{u_{j+1,k+1} - u_{j+1,k}}{h_2} \right) \left(\frac{u_{j,k+1} - u_{j,k}}{h_2} \right) \right\} dt \leq 0. \end{aligned} \quad (45)$$

The proof of this inequality relies essentially on the inequality $-(a^2 + b^2) + 2ab \leq 0$, for all $a, b \in \mathbb{R}$. Reporting (44) and the above inequality in (41), we get for all $T > 2\sqrt{2} + \frac{\sqrt{2}}{4} \max(h_1^2, h_2^2)$

$$\begin{aligned} & \left(T - 2\sqrt{2} - \frac{\sqrt{2}}{4} \max(h_1^2, h_2^2) \right) E_{h_1, h_2}(0) \\ & \leq \frac{h_1 h_2}{2} \sum_{(j,k) \in \tilde{\mathcal{I}}^+} j \int_0^T \left(\frac{u_{j-1,k}}{h_1} \right)^2 dt + \frac{h_1 h_2}{2} \sum_{(j,k) \in \hat{\mathcal{I}}^+} k \int_0^T \left(\frac{u_{j,k-1}}{h_2} \right)^2 dt \\ & + \frac{h_1 h_2}{4} \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left\{ h_1^2 \left(\frac{u'_{j+1,k} - u'_{j,k}}{h_1} \right)^2 + h_2^2 \left(\frac{u'_{j,k+1} - u'_{j,k}}{h_2} \right)^2 \right\} dt. \end{aligned} \quad (46)$$

In (46) the boundary terms corresponding to the subsets $\tilde{\mathcal{I}}^-$ and $\hat{\mathcal{I}}^-$ have been neglected since they are negative. The next steps are devoted to the absorption of the remaining boundary terms in the right hand side of (46).

Step 4: *Absorption of the boundary terms*

Let us introduce the function $\mathbf{f} = (f^1, f^2) \in (W^{1,\infty}(\Omega))^2$ of support $\hat{\omega}$ localized on the neighborhood of Γ_0 (defined in (8)) and included in ω ; more precisely, we write :

$$\begin{cases} f^1(\mathbf{x}) = 1 \text{ on } \{\mathbf{x} \in \Gamma_0, \nu(\mathbf{x}) = (1, 0)\}, \\ f^2(\mathbf{x}) = 1 \text{ on } \{\mathbf{x} \in \Gamma_0, \nu(\mathbf{x}) = (0, 1)\}, \\ \mathbf{f}(\mathbf{x}) = \mathbf{0} \text{ in } \Omega \setminus \mathcal{V}_r(\Gamma_0), \end{cases} \quad (47)$$

where $\mathcal{V}_r(\Gamma_0)$ denotes a neighborhood of Γ_0 for the euclidean distance,

$$\mathcal{V}_r(\Gamma_0) = \{\mathbf{x} \in \Omega ; d(\mathbf{x}, \Gamma_0) \leq r\}, \quad (48)$$

and r a sufficiently small positive real. We then define the discrete cut-off function of support the grid $G(\hat{\omega}_{h_1, h_2})$ of $\hat{\omega}_{h_1, h_2}$ defined as the union of cells $B_{j,k}$ contained in $\hat{\omega}$:

$$\begin{cases} f_{j,k}^1 = 1 & \text{if } (j, k) \in \tilde{I}^+, & f_{j,k}^2 = 1 & \text{if } (j, k) \in \hat{I}^+, \\ \mathbf{f}_{j,k} = 0 & \text{if } (j, k) \notin G(\Omega_{h_1, h_2}) \setminus (\mathcal{V}_r(\tilde{I}^+ \cup \hat{I}^+)). \end{cases} \quad (49)$$

We also introduce a smooth nonnegative function $g = g(t)$ such that

$$g(0) = g(T) = 0, \quad g(t) = 1 \text{ in } [\delta, T - \delta] \quad (50)$$

where $\delta > 0$ is sufficiently small. Now, we multiply the first equation in (30) by

$$f_{j,k}^1 \left(\frac{u_{j+1,k} - u_{j-1,k}}{2h_1} \right) g(t) + f_{j,k}^2 \left(\frac{u_{j,k+1} - u_{j,k-1}}{2h_2} \right) g(t) \quad (51)$$

and proceed as in the first step. We find

$$\begin{aligned} & \frac{1}{2} \sum_{(j,k) \in \tilde{I}^+} j \int_0^T g(t) \left(\frac{u_{j-1,k}}{h_1} \right)^2 dt + \frac{1}{2} \sum_{(j,k) \in \hat{I}^+} k \int_0^T g(t) \left(\frac{u_{j,k-1}}{h_2} \right)^2 dt \\ &= \frac{1}{2} \sum_{(j,k) \in \hat{\omega}_{h_1, h_2}} \left(\frac{f_{j+1,k}^1 - f_{j,k}^1}{h_1} \int_0^T g(t) u'_{j+1,k} u'_{j,k} dt + \frac{f_{j,k+1}^2 - f_{j,k}^2}{h_2} \int_0^T g(t) u'_{j,k+1} u'_{j,k} dt \right) \\ & \quad - \sum_{(j,k) \in \hat{\omega}_{h_1, h_2}} \left(f_{j,k}^1 \int_0^T g'(t) u'_{j,k}(t) \left(\frac{u_{j+1,k} - u_{j-1,k}}{2h_1} \right) dt + f_{j,k}^2 \int_0^T g'(t) u'_{j,k} \left(\frac{u_{j,k+1} - u_{j,k-1}}{2h_2} \right) dt \right) \\ & \quad + \frac{1}{2} \sum_{(j,k) \in \hat{\omega}_{h_1, h_2}} \frac{f_{j+1,k}^1 - f_{j,k}^1}{h_1} \int_0^T g(t) \left(\frac{u_{j+1,k} - u_{j,k}}{h_1} \right)^2 dt \\ & \quad + \frac{1}{2} \sum_{(j,k) \in \hat{\omega}_{h_1, h_2}} \frac{f_{j,k+1}^2 - f_{j,k}^2}{h_2} \int_0^T g(t) \left(\frac{u_{j,k+1} - u_{j,k}}{h_2} \right)^2 dt \\ & \quad - \sum_{(j,k) \in \hat{\omega}_{h_1, h_2}} f_{j,k}^2 \int_0^T g(t) \left(\frac{u_{j+1,k} - 2u_{j,k} + u_{j-1,k}}{h_1^2} \right) \left(\frac{u_{j,k+1} - u_{j,k-1}}{2h_2} \right) dt \\ & \quad - \sum_{(j,k) \in \hat{\omega}_{h_1, h_2}} f_{j,k}^1 \int_0^T g(t) \left(\frac{u_{j,k+1} - 2u_{j,k} + u_{j,k-1}}{h_2^2} \right) \left(\frac{u_{j+1,k} - u_{j-1,k}}{2h_1} \right) dt. \end{aligned} \quad (52)$$

From the relation (52), we have the following result:

Lemma 2.5. *The bounded terms can be bounded as follows:*

$$\begin{aligned}
& \frac{1}{2} \sum_{(j,k) \in \tilde{I}^-} j \int_{\delta}^{T-\delta} \left(\frac{u_{j-1,k}}{h_1} \right)^2 dt + \frac{1}{2} \sum_{(j,k) \in \tilde{I}^+} k \int_{\delta}^{T-\delta} \left(\frac{u_{j,k-1}}{h_2} \right)^2 dt \\
& \leq C \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} a_{j,k} \int_0^T (u'_{j,k}(t))^2 dt + C \sum_{(j,k) \in \tilde{\omega}_{h_1, h_2}} \int_0^T g(t) (u_{j,k}(t))^2 dt \\
& + C \sum_{(j,k) \in \tilde{\omega}_{h_1, h_2}} \int_0^T g(t) \left\{ \left(\frac{u_{j+1,k} - u_{j,k}}{h_1} \right)^2 + \left(\frac{u_{j,k+1} - u_{j,k}}{h_2} \right)^2 \right\} dt.
\end{aligned} \tag{53}$$

Proof of lemma 2.5 - Performing as in [23] the result follows. \square

Reporting (53) in (46) with $T > 2\sqrt{2} + \frac{\sqrt{2}}{4} \max(h_1^2, h_2^2)$ and δ small enough such that $T - 2\delta > 0$, we obtain, using the invariance by the translation in time,

$$\begin{aligned}
E_{h_1, h_2}(0) & \leq Ch_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T a_{j,k} (u'_{j,k}(t))^2 dt \\
& + Ch_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left\{ h_1^2 \left(\frac{u'_{j+1,k} - u'_{j,k}}{h_1} \right)^2 + h_2^2 \left(\frac{u'_{j,k+1} - u'_{j,k}}{h_2} \right)^2 \right\} dt \\
& + Ch_1 h_2 \sum_{(j,k) \in \tilde{\omega}_{h_1, h_2}} \int_0^T g(t) (u_{j,k}(t))^2 dt \\
& + Ch_1 h_2 \sum_{(j,k) \in \tilde{\omega}_{h_1, h_2}} \int_0^T g(t) \left\{ \left(\frac{u_{j+1,k} - u_{j,k}}{h_1} \right)^2 + \left(\frac{u_{j,k+1} - u_{j,k}}{h_2} \right)^2 \right\} dt.
\end{aligned} \tag{54}$$

It remains to show that the last two terms in the right hand side of (54) may be uniformly absorbed:

Step 5: *Absorption of the discrete gradient term*

To absorb the last term in (54) which corresponds to the energy localized in the discrete gradient of $\tilde{\omega}_{h_1, h_2}$, we use a localized version of the formula of equipartition of energy. We introduce the function $\eta \in W^{1, \infty}(\Omega)$ with the support ω of the damping a containing strictly the support $\tilde{\omega}$ of the function \mathbf{f} . As a consequence, proceeding as in step 4 (see also [23]), an approximation $(\eta_{j,k})_{j,k}$ of η may be defined as follows:

$$\eta_{j,k} = 1 \quad \text{if } (j,k) \in \tilde{I}^+ \cup \hat{I}^+, \quad \eta_{j,k} = 0 \quad \text{if } (j,k) \notin G(\omega_{h_1, h_2}). \tag{55}$$

Then, multiplying the first equation of (30) by $g(t)\eta_{j,k}^2 u_{j,k}$ (with g defined in (50)) and proceeding as in steps 1 and 4, we obtain (see [23]) using $\tilde{\omega}_{h_1, h_2} \subset \omega_{h_1, h_2}$ that:

$$\begin{aligned}
E_{h_1, h_2}(0) & \leq Ch_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T a_{j,k} (u'_{j,k}(t))^2 dt + Ch_1 h_2 \sum_{(j,k) \in \omega_{h_1, h_2}} \int_0^T g(t) (u_{j,k}(t))^2 dt \\
& + Ch_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left\{ h_1^2 \left(\frac{u'_{j+1,k} - u'_{j,k}}{h_1} \right)^2 + h_2^2 \left(\frac{u'_{j,k+1} - u'_{j,k}}{h_2} \right)^2 \right\} dt,
\end{aligned} \tag{56}$$

for any $T > 2\sqrt{2}$.

Step 6: Conclusion

To conclude the proof it is sufficient to show that there exists a constant $C > 0$ such that

$$\begin{aligned} \sum_{(j,k) \in \omega_{h_1, h_2}} \int_0^T g(t)(u_{j,k}(t))^2 dt &\leq C \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T a_{j,k}(u'_{j,k}(t))^2 dt \\ &+ Ch_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \int_0^T \left\{ h_1^2 \left(\frac{u'_{j+1,k} - u'_{j,k}}{h_1} \right)^2 + h_2^2 \left(\frac{u'_{j,k+1} - u'_{j,k}}{h_2} \right)^2 \right\} dt. \end{aligned} \quad (57)$$

To do this we may argue by means of a classical compactness uniqueness argument as in Proposition 4.1 in [26]. According to this argument, (57) reduces to an unique continuation property for the solution of the continuous wave equation. But, for that to be true, we need a convergence result guaranteeing that the solutions of the discrete system converge to those of the wave equation. This is done in Theorem 3.1 of section 3. Since the analysis described above is analogous to the one developed in [23], we omit the details. This completes the proof of Theorem 2.1. \square

3. CONVERGENCE OF THE NUMERICAL SCHEME

The aim of this section is to prove a convergence result of the solution of the semi-discrete scheme (21) to the solution of system (1). This allows to conclude that the presence of the viscous terms acts only as a regularization term to damp out the high frequency components and does not change the limit behavior. This will also permit to pass to the limit in the equation (28) of Theorem 2.1 to find again the exponential decay of the energy E (equation (6)). Before stating the convergence result, let us introduce the two following extensions P_{h_1, h_2} and Q_{h_1, h_2} defined by:

$$P_{h_1, h_2} \mathbf{v}_{h_1, h_2} = \begin{cases} \text{the continuous function linear in each cell } B_{j,k} \text{ such that} \\ P_{h_1, h_2} \mathbf{v}_{h_1, h_2}(x_{1,j}, x_{2,k}) = v_{j,k}, \\ P_{h_1, h_2} \mathbf{v}_{h_1, h_2}(x_{1,j+1}, x_{2,k}) = v_{j+1,k}, \\ P_{h_1, h_2} \mathbf{v}_{h_1, h_2}(x_{1,j}, x_{2,k+1}) = v_{j,k+1}, \\ P_{h_1, h_2} \mathbf{v}_{h_1, h_2}(x_{1,j+1}, x_{2,k+1}) = v_{j+1,k+1}, \quad j \in [0, J], k \in \mathcal{I}_j. \end{cases} \quad (58)$$

$$Q_{h_1, h_2} \mathbf{v}_{h_1, h_2} = \begin{cases} \text{the step function defined in each square} \\ \left((j - \frac{1}{2})h_1, (j + \frac{1}{2})h_1 \right) \times \left((k - \frac{1}{2})h_2, (k + \frac{1}{2})h_2 \right) \cap \Omega \\ \text{by } Q_{h_1, h_2} \mathbf{v}_{h_1, h_2}(\mathbf{x}) = v_{j,k}, \quad j \in [0, J], k \in \mathcal{I}_j. \end{cases} \quad (59)$$

We now state our convergence result:

Theorem 3.1 (Convergence of \mathbf{y}_{h_1, h_2}). *Let \mathbf{y}_{h_1, h_2} denote the solution of (21). Assume that \mathbf{a}_{h_1, h_2} , $(\mathbf{y}_0)_{h_1, h_2}$ and $(\mathbf{y}_1)_{h_1, h_2}$ are such that there is a nonnegative constant C independent of h_1, h_2 with*

$$\begin{cases} E_{h_1, h_2}(0) \leq C, \quad Q_{h_1, h_2} \mathbf{a}_{h_1, h_2} \rightarrow a \text{ weakly } * \text{ in } L^\infty(\Omega), \\ P_{h_1, h_2}(\mathbf{y}_0)_{h_1, h_2} \rightarrow y_0 \text{ weakly in } H_0^1(\Omega), \quad Q_{h_1, h_2}(\mathbf{y}_1)_{h_1, h_2} \rightarrow y_1 \text{ weakly in } L^2(\Omega). \end{cases} \quad (60)$$

Then, we have

$$\begin{cases} P_{h_1, h_2} \mathbf{y}_{h_1, h_2} \rightarrow y \text{ weakly } * \text{ in } L^\infty(0, \infty; H_0^1(\Omega)), \\ Q_{h_1, h_2} \frac{\partial \mathbf{y}_{h_1, h_2}}{\partial t} \rightarrow \frac{\partial y}{\partial t} \text{ weakly } * \text{ in } L^\infty(0, \infty; L^2(\Omega)), \end{cases} \quad (61)$$

where y is the solution of system (1).

Proof: We decompose the proof in several steps.

Step 1: *A priori estimates and weak convergence*

Straightforward computations first lead to the two following equalities:

$$\begin{aligned} & \int_{\Omega_{h_1, h_2}} \nabla P_{h_1, h_2} \mathbf{v}_{h_1, h_2}(\mathbf{x}) \cdot \nabla P_{h_1, h_2} \mathbf{w}_{h_1, h_2}(\mathbf{x}) dx \\ &= h_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \left(\frac{v_{j+1, k} - v_{j, k}}{h_1} \right) \left(\frac{w_{j+1, k} - w_{j, k}}{h_1} \right) \\ & \quad + h_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} \left(\frac{v_{j, k+1} - v_{j, k}}{h_2} \right) \left(\frac{w_{j, k+1} - w_{j, k}}{h_2} \right), \end{aligned} \quad (62)$$

$$\int_{\Omega_{h_1, h_2}} Q_{h_1, h_2} \mathbf{v}_{h_1, h_2}(\mathbf{x}) Q_{h_1, h_2} \mathbf{w}_{h_1, h_2}(\mathbf{x}) dx = h_1 h_2 \sum_{j=0}^J \sum_{k \in \mathcal{I}_j} v_{j, k} w_{j, k}.$$

Then, according to the definition (24), we get, for every $t \geq 0$,

$$E_{h_1, h_2}(t) = \frac{1}{2} (\|P_{h_1, h_2} \mathbf{y}_{h_1, h_2}(t)\|_{H_0^1(\Omega_{h_1, h_2})}^2 + \|Q_{h_1, h_2} \mathbf{y}'_{h_1, h_2}(t)\|_{L^2(\Omega_{h_1, h_2})}^2). \quad (63)$$

Then, thanks to (60), (63) and the decreasing character of E_{h_1, h_2} , we deduce that $P_{h_1, h_2} \mathbf{y}_{h_1, h_2}$ is bounded in $L^\infty(0, \infty; H_0^1(\Omega_{h_1, h_2})) \cap W^{1, \infty}(0, \infty; L^2(\Omega_{h_1, h_2}))$, while $Q_{h_1, h_2} \mathbf{y}'_{h_1, h_2}$ is bounded in $L^\infty(0, \infty; L^2(\Omega_{h_1, h_2}))$. Then, since

$$\|Q_{h_1, h_2} \mathbf{y}_{h_1, h_2}\|_{L^2(\Omega_{h_1, h_2})} \leq \|(P_{h_1, h_2} - Q_{h_1, h_2}) \mathbf{y}_{h_1, h_2}\|_{L^2(\Omega_{h_1, h_2})} + \|P_{h_1, h_2} \mathbf{y}_{h_1, h_2}\|_{L^2(\Omega_{h_1, h_2})} \quad (64)$$

and E_{h_1, h_2} is bounded, straightforward computations leads to

$$\|P_{h_1, h_2} \mathbf{y}_{h_1, h_2} - Q_{h_1, h_2} \mathbf{y}_{h_1, h_2}\|_{L^2(\Omega_{h_1, h_2})} = O(h_1 h_2), \forall t \geq 0, \quad (65)$$

what allows to deduce that $Q_{h_1, h_2} \mathbf{y}_{h_1, h_2}$ is bounded in $L^\infty(0, \infty; L^2(\Omega_{h_1, h_2}))$. In the same way, multiplying the first equation of (21) by $y_{j, k}$, integrating by part over $[0, T]$ and rearranging the index, one can also show that $P_{h_1, h_2} \mathbf{y}'_{h_1, h_2}$ is bounded in $L^\infty(0, \infty; L^2(\Omega_{h_1, h_2}))$ such that $P_{h_1, h_2} \mathbf{y}_{h_1, h_2}$ is also bounded in $W^{1, \infty}(0, \infty; L^2(\Omega_{h_1, h_2}))$.

Furthermore, taking into account that solutions both in the continuous and the semi-discrete case satisfy Dirichlet boundary conditions, we can extend them by zero to the whole unit box B . The extension operator is denoted by the symbol $\tilde{\cdot}$. Consequently, we can consider each \mathbf{y}_{h_1, h_2} , as well as the solution of (1), as solution of $H_0^1(B)$, and therefore the boundedness of $E_{h_1, h_2}(t)$ mentioned above still holds in the domain B , independent

of h_1 and h_2 . Thus, we can extract subsequences (that we still denote by $\tilde{\mathbf{y}}_{h_1, h_2}$), such that

$$\begin{cases} P_{h_1, h_2} \tilde{\mathbf{y}}_{h_1, h_2} \rightarrow \tilde{y} & \text{weakly } * \text{ in } L^\infty(0, \infty; H_0^1(B)), \\ P_{h_1, h_2} \tilde{\mathbf{y}}'_{h_1, h_2} \rightarrow \tilde{y}' & \text{weakly } * \text{ in } L^\infty(0, \infty; L^2(B)), \\ P_{h_1, h_2} \tilde{\mathbf{y}}_{h_1, h_2} \rightarrow \tilde{y} & \text{strongly in } L^2_{loc}(0, \infty; L^2(B)), \\ Q_{h_1, h_2} \tilde{\mathbf{y}}_{h_1, h_2} \rightarrow \tilde{y} & \text{weakly } * \text{ in } L^\infty(0, \infty; L^2(B)), \\ Q_{h_1, h_2} \tilde{\mathbf{y}}'_{h_1, h_2} \rightarrow \tilde{y}' & \text{weakly } * \text{ in } L^\infty(0, \infty; L^2(B)), \\ h_i P_{h_1, h_2} \tilde{\mathbf{y}}'_{h_1, h_2} \rightarrow 0 & \text{weakly } * \text{ in } L^2(0, \infty; H_0^1(B)), \quad i = 1, 2. \end{cases} \quad (66)$$

The last convergence in (66) follows from the second one and the boundness of the sequence $\{h_i P_{h_1, h_2}\}$, $i = 1, 2$, in the space $L^2(0, \infty; H_0^1(B))$ easily obtained by multiplying the first equation of (21) by $\tilde{y}'_{j, k}$, integration by parts and rearranging the index.

Note that in (66), we implicit claim that the limits of $P_{h_1, h_2} \tilde{\mathbf{y}}_{h_1, h_2}$ and $Q_{h_1, h_2} \tilde{\mathbf{y}}_{h_1, h_2}$ are the same which is a consequence of (65).

Step 2: *Characterization of the limit \tilde{y}*

Let us now show that the limit \tilde{y} is the solution of (1). To this end, let $w \in \mathcal{D}(\Omega \times (0, \infty))$ and extend it by zero to the whole B . We set $\tilde{\mathbf{w}}_{h_1, h_2} = (\tilde{w}_{j, k})_{j, k}$ where $\tilde{w}_{j, k} = \tilde{w}(x_{1, j}, x_{2, k})$. Multiplying the first equation of (21) by $\tilde{w}_{j, k}$, integrating by parts in $(0, \infty)$ and taking the sum, we find

$$\begin{aligned} & \sum_{j=0}^J \sum_{k=0}^K \int_0^\infty \tilde{y}_{j, k} \tilde{w}''_{j, k} dt - \sum_{j=0}^J \sum_{k=0}^K \int_0^\infty \tilde{a}_{j, k} \tilde{y}_{j, k} \tilde{w}'_{j, k} dt \\ & + \sum_{j=0}^J \sum_{k=0}^K \int_0^\infty \left[\left(\frac{\tilde{y}_{j+1, k} - \tilde{y}_{j, k}}{h_1} \right) \left(\frac{\tilde{w}_{j+1, k} - \tilde{w}_{j, k}}{h_1} \right) + \left(\frac{\tilde{y}_{j, k+1} - \tilde{y}_{j, k}}{h_2} \right) \left(\frac{\tilde{w}_{j, k+1} - \tilde{w}_{j, k}}{h_2} \right) \right] dt \\ & + \sum_{j=0}^J \sum_{k=0}^K \int_0^\infty \left[h_1^2 \left(\frac{\tilde{y}'_{j+1, k} - \tilde{y}'_{j, k}}{h_1} \right) \left(\frac{\tilde{w}_{j+1, k} - \tilde{w}_{j, k}}{h_1} \right) \right. \\ & \quad \left. + h_2^2 \left(\frac{\tilde{y}'_{j, k+1} - \tilde{y}'_{j, k}}{h_2} \right) \left(\frac{\tilde{w}_{j, k+1} - \tilde{w}_{j, k}}{h_2} \right) \right] dt = 0. \end{aligned} \quad (67)$$

Using the definitions of P_{h_1, h_2} and Q_{h_1, h_2} , we can check that (67) is equivalent to

$$\begin{aligned} & \int_0^\infty \int_B Q_{h_1, h_2} \tilde{\mathbf{y}}_{h_1, h_2} Q_{h_1, h_2} \tilde{\mathbf{w}}''_{h_1, h_2} dx dt \\ & + \int_0^\infty \int_B \nabla P_{h_1, h_2} \tilde{\mathbf{y}}_{h_1, h_2} \cdot \nabla P_{h_1, h_2} \tilde{\mathbf{w}}_{h_1, h_2} dx dt \\ & - \int_0^\infty \int_B Q_{h_1, h_2} \tilde{\mathbf{y}}_{h_1, h_2} Q_{h_1, h_2} \tilde{\mathbf{w}}'_{h_1, h_2} Q_{h_1, h_2} \tilde{\mathbf{a}}_{h_1, h_2} dx dt \\ & + h_1 h_2 \sum_{j=0}^J \sum_{k=0}^K \int_0^\infty \left[h_1^2 \left(\frac{\tilde{y}'_{j+1, k} - \tilde{y}'_{j, k}}{h_1} \right) \left(\frac{\tilde{w}_{j+1, k} - \tilde{w}_{j, k}}{h_1} \right) \right. \\ & \quad \left. + h_2^2 \left(\frac{\tilde{y}'_{j, k+1} - \tilde{y}'_{j, k}}{h_2} \right) \left(\frac{\tilde{w}_{j, k+1} - \tilde{w}_{j, k}}{h_2} \right) \right] dt = 0. \end{aligned} \quad (68)$$

Now, we observe that, if we show that for every $w \in \mathcal{D}(\Omega \times (0, \infty))$

$$\begin{cases} P_{h_1, h_2} \tilde{\mathbf{w}}_{h_1, h_2} \rightarrow \tilde{w} \text{ strongly in } L^2(0, \infty; H_0^1(B)), \\ Q_{h_1, h_2} \tilde{\mathbf{w}}_{h_1, h_2} \rightarrow \tilde{w} \text{ strongly in } L^2(0, \infty; L^2(B)), \end{cases} \quad (69)$$

then, we will be able to pass to the limit in (68) leading with (66) to

$$\int_0^\infty \int_{B \cap \text{supp } w} \tilde{y} w'' dx dt + \int_0^\infty \int_{B \cap \text{supp } w} \nabla \tilde{y} \cdot \nabla w dx dt - \int_0^\infty \int_{B \cap \text{supp } w} \tilde{y} w' a dx dt = 0, \quad (70)$$

where $\text{supp } w$ denotes the support of $w \in \mathcal{D}(\Omega \times (0, \infty))$, what guaranties that

$$\int_0^\infty \int_\Omega \tilde{y} w'' dx dt + \int_0^\infty \int_\Omega \nabla \tilde{y} \cdot \nabla w dx dt - \int_0^\infty \int_\Omega \tilde{y} w' a dx dt = 0 \quad (71)$$

for all $w \in \mathcal{D}(\Omega \times (0, \infty))$.

Step 3: *Property of $\tilde{y}(t)$ for all $t \geq 0$*

Let us first show that $\tilde{y}(t) \in H_0^1(\Omega)$ for all $t \geq 0$. Since $P_{h_1, h_2} \mathbf{y}_{h_1, h_2}$ belongs to $H_0^1(\Omega_{h_1, h_2}) \subset H_0^1(B)$, the limit $\tilde{y}(t)$ belongs to $H_0^1(B)$. Furthermore, since $\partial\Omega$ is Lipschitz by assumption and also $\partial\Omega_{h_1, h_2}$ by construction, it possess the so-called ϵ -cone property, introduced in [4]. This implies that the bounded sequence $(\Omega_{h_1, h_2})_{h_1, h_2}$ defined, for each h_1, h_2 fixed, as the largest domain included in Ω and union of cell $B_{j, k}$ converges towards Ω with respect to the complementary-Hausdorff topology (see [20]). This property implies, in particular, that the limit of the sequence $P_{h_1, h_2} \mathbf{y}_{h_1, h_2}$ defined on Ω_{h_1, h_2} extended to zero on $B \setminus \Omega$ belongs to $H_0^1(\Omega)$ for any time $t \geq 0$. Actually, in our situation - (Ω_{h_1, h_2}) is an increasing sequence include in Ω - we could obtain the same property using more classical results of spectral analysis (see [10] and the references therein).

Let us now show that $\tilde{y}(0) = y_0$ and $\tilde{y}'(0) = y_1$. For this purpose, let $v \in \mathcal{D}(\Omega)$ and $l \in \mathcal{D}([0, \infty))$. Then, multiplying the first equation of (21) by $v_{j, k} l$, integrating by parts on $[0, \infty)$ and taking the sum, we find

$$\begin{aligned} & - \sum_{j=0}^J \sum_{k=0}^K (y_1)_{j, k} \tilde{v}_{j, k} l(0) + \sum_{j=0}^J \sum_{k=0}^K (y_0)_{j, k} \tilde{v}_{j, k} l'(0) - \sum_{j=0}^J \sum_{k=0}^K \tilde{a}_{j, k} (y_0)_{j, k} \tilde{v}_{j, k} l(0) \\ & + \sum_{j=0}^J \sum_{k=0}^K \int_0^\infty \tilde{y}_{j, k} \tilde{v}_{j, k} l'' dt - \sum_{j=0}^J \sum_{k=0}^K \int_0^\infty \tilde{a}_{j, k} \tilde{y}_{j, k} \tilde{v}_{j, k} l' dt \\ & + \sum_{j=0}^J \sum_{k=0}^K \int_0^\infty \left[\left(\frac{\tilde{y}_{j+1, k} - \tilde{y}_{j, k}}{h_1} \right) \left(\frac{\tilde{v}_{j+1, k} - \tilde{v}_{j, k}}{h_1} \right) + \left(\frac{\tilde{y}_{j, k+1} - \tilde{y}_{j, k}}{h_2} \right) \left(\frac{\tilde{v}_{j, k+1} - \tilde{v}_{j, k}}{h_2} \right) \right] l dt \\ & + \sum_{j=0}^J \sum_{k=0}^K \int_0^\infty \left[h_1^2 \left(\frac{\tilde{y}'_{j+1, k} - \tilde{y}'_{j, k}}{h_1} \right) \left(\frac{\tilde{v}_{j+1, k} - \tilde{v}_{j, k}}{h_1} \right) \right. \\ & \quad \left. + h_2^2 \left(\frac{\tilde{y}'_{j, k+1} - \tilde{y}'_{j, k}}{h_2} \right) \left(\frac{\tilde{v}_{j, k+1} - \tilde{v}_{j, k}}{h_2} \right) \right] l dt = 0. \end{aligned} \quad (72)$$

Using the definitions of P_{h_1, h_2} and Q_{h_1, h_2} and proceeding as in the last steps, we can check that (72) is equivalent to

$$\begin{aligned}
& -l(0) \int_B Q_{h_1, h_1}(\mathbf{y}_1)_{h_1, h_2} Q_{h_1, h_2} \tilde{\mathbf{v}}_{h_1, h_2} dx + l'(0) \int_B Q_{h_1, h_2}(\mathbf{y}_0)_{h_1, h_2} Q_{h_1, h_2} \tilde{\mathbf{v}}_{h_1, h_2} dx \\
& -l(0) \int_B Q_{h_1, h_2}(\mathbf{y}_0)_{h_1, h_2} Q_{h_1, h_2} \tilde{\mathbf{v}}_{h_1, h_2} Q_{h_1, h_2} \tilde{\mathbf{a}}_{h_1, h_2} dx \\
& + \int_0^\infty \int_B Q_{h_1, h_2} \tilde{\mathbf{y}}_{h_1, h_2} Q_{h_1, h_2} \tilde{\mathbf{v}}_{h_1, h_2} l'' dx dt \\
& + \int_0^\infty \int_B \nabla P_{h_1, h_2} \tilde{\mathbf{y}}_{h_1, h_2} \cdot \nabla P_{h_1, h_2} \tilde{\mathbf{v}}_{h_1, h_2} l dx dt \\
& - \int_0^\infty \int_B Q_{h_1, h_2} \tilde{\mathbf{y}}_{h_1, h_2} Q_{h_1, h_2} \tilde{\mathbf{v}}_{h_1, h_2} Q_{h_1, h_2} \tilde{\mathbf{a}}_{h_1, h_2} l' dx dt \\
& + h_1 h_2 \sum_{j=0}^J \sum_{k=0}^K \int_0^\infty \left[h_1^2 \left(\frac{\tilde{y}'_{j+1, k} - \tilde{y}'_{j, k}}{h_1} \right) \left(\frac{\tilde{v}_{j+1, k} - \tilde{v}_{j, k}}{h_1} \right) \right. \\
& \quad \left. + h_2^2 \left(\frac{\tilde{y}'_{j, k+1} - \tilde{y}'_{j, k}}{h_2} \right) \left(\frac{\tilde{v}_{j, k+1} - \tilde{v}_{j, k}}{h_2} \right) \right] l dt = 0.
\end{aligned} \tag{73}$$

Passing to the limit as $(h_1, h_2) \rightarrow (0, 0)$ in (73), we get

$$\begin{aligned}
& -l(0) \int_\Omega y_1 v dx + l'(0) \int_\Omega y_0 v dx - l(0) \int_\Omega y_0 v a dx \\
& + \int_0^\infty \int_\Omega y v l'' dx dt + \int_0^\infty \int_\Omega \nabla y \cdot \nabla v l dx dt - \int_0^\infty \int_B y v a l' dx dt = 0
\end{aligned} \tag{74}$$

from which we derive $\tilde{y}(0) = y_0$ and $\tilde{y}'(0) = y_1$. Since system (1) has a unique solution, we conclude that the convergence results in (66) hold for the whole sequence $\{(h_1, h_2)\}$.

Step 3: Proof of (69)

To conclude the proof, let us prove (69). First observe that, for all $w \in \mathcal{D}(\Omega \times (0, \infty))$, we have

$$\begin{aligned}
& \int_0^\infty \int_B |\nabla(P_{h_1, h_2} \tilde{\mathbf{w}}_{h_1, h_2}) - \nabla \tilde{w}|^2 dx dt = \\
& h_1 h_2 \sum_{j=0}^J \sum_{k=0}^K \int_0^\infty \left[\left(\frac{\tilde{w}_{j+1, k} - \tilde{w}_{j, k}}{h_1} \right)^2 + \left(\frac{\tilde{w}_{j, k+1} - \tilde{w}_{j, k}}{h_2} \right)^2 \right] dt + \int_0^\infty \int_B |\nabla \tilde{w}|^2 dx dt \\
& - 2h_1 h_2 \sum_{j=0}^J \sum_{k=0}^K \int_0^\infty \left\{ \left(\frac{\tilde{w}_{j+1, k} - \tilde{w}_{j, k}}{h_1} \right) \int_0^1 \int_0^1 \frac{\partial \tilde{w}}{\partial x_1}((j + s_1)h_1, x_2) ds_1 dx_2 \right. \\
& \left. + \left(\frac{\tilde{w}_{j, k+1} - \tilde{w}_{j, k}}{h_2} \right) \int_0^1 \int_0^1 \frac{\partial \tilde{w}}{\partial x_2}(x_1, (k + s_2)h_2) ds_2 dx_1 \right\} dt, \quad \text{where } s = (s_1, s_2),
\end{aligned} \tag{75}$$

so that using the properties of Riemann sums, it follows that

$$\lim_{(h_1, h_2) \rightarrow 0} \int_0^\infty \int_\Omega |\nabla(P_{h_1, h_2} \mathbf{w}_{h_1, h_2}) - \nabla w|^2 dx dt = 0, \tag{76}$$

leading to the first convergence in (69). The second one is obtained using the same argument and with

$$\begin{aligned}
\int_0^\infty \int_B (Q_{h_1, h_2} \tilde{w}_{h_1, h_2} - \tilde{w})^2 dx dt &= h_1 h_2 \sum_{j=1}^J \sum_{k=0}^K \int_0^\infty |\tilde{w}(jh_1, kh_2, t)|^2 dt + \int_0^\infty \int_B \tilde{w}^2 dx dt \\
- 2h_1 h_2 \sum_{j=0}^J \sum_{k=0}^K \int_0^\infty &\left\{ \int_0^{\frac{1}{2}} \int_0^{\frac{1}{2}} \tilde{w}_{j,k} \tilde{w} \left((j+s_1)h_1, (k+s_2)h_2 \right) ds_1 ds_2 \right. \\
&+ \int_0^{\frac{1}{2}} \int_{\frac{1}{2}}^1 \tilde{w}_{j+1,k} \tilde{w} \left((j+s_1)h_1, (k+s_2)h_2 \right) ds_1 ds_2 \\
&+ \int_{\frac{1}{2}}^1 \int_0^{\frac{1}{2}} \tilde{w}_{j,k+1} \tilde{w} \left((j+s_1)h_1, (k+s_2)h_2 \right) ds_1 ds_2 \\
&\left. + \int_{\frac{1}{2}}^1 \int_{\frac{1}{2}}^1 \tilde{w}_{j+1,k+1} \tilde{w} \left((j+s_1)h_1, (k+s_2)h_2 \right) ds_1 ds_2 \right\} dt
\end{aligned} \tag{77}$$

Again, using the properties of Riemann sums, we obtain the second convergence in (69). This concludes the proof of Theorem 3.1. \square

Remark 3.2. The convergence of the finite difference space semi-discretization without numerical viscous term was previously proved in [21]. In this work the authors use the approach considered here to construct a uniform exponentially stable approximations for an abstract class of second order evolution equations with bounded feedback controls.

Remark 3.3. Assuming strong convergence of the initial data instead of (60), one may also show, by a compactness property related to the Hausdorff topology, that convergence (61) holds for a more strong topology (see for instance [14], chapter 4) leading to

$$E_{h_1, h_2}(t) \rightarrow E(t), \quad \forall t \geq 0, \quad \text{when } (h_1, h_2) \rightarrow (0, 0). \tag{78}$$

Therefore, up to this additional strong convergence assumption, we may take the limit in the relation (28) to reobtain the inequality (6) for weak solution of (1) defined on Ω , Ω being the limit for the Hausdorff metric of a bounded sequence Ω_{h_1, h_2} . Indeed, assume that the initial data satisfies

$$E_{h_1, h_2}(0) \rightarrow E(0) \text{ as } (h_1, h_2) \rightarrow (0, 0). \tag{79}$$

Since $E_{h_1, h_2}(t)$ decreases to zero as $t \rightarrow \infty$, it follows from (25) that

$$E_{h_1, h_2}(0) = h_1 h_2 \sum_{j=0}^J \sum_{k \in I_j} \int_0^\infty a_{j,k} (y'_{j,k})^2 dt + h_1 h_2 \sum_{j=0}^J \sum_{k \in I_j} \int_0^\infty \left\{ h_1^2 \left(\frac{y'_{j+1,k} - y'_{j,k}}{h_1} \right)^2 + h_2^2 \left(\frac{y'_{j+1,k} - y'_{j,k}}{h_1} \right)^2 \right\} dt. \tag{80}$$

On the other hand, being Ω of C^2 class, $E(t) \rightarrow 0$ as $t \rightarrow \infty$, and therefore

$$E(0) = \int_0^\infty \int_\Omega a(\mathbf{x}) |y_t|^2 dx dt. \tag{81}$$

Thus, from (79)-(81) it follows that

$$\begin{aligned} \limsup_{(h_1, h_2) \rightarrow (0, 0)} h_1 h_2 \sum_{j=0}^J \sum_{k \in I_j} \int_0^\infty a_{j,k}(y'_{j,k})^2 dt &= \limsup_{(h_1, h_2) \rightarrow (0, 0)} \int_0^\infty \int_\Omega Q_{h_1, h_2} a_{h_1, h_2} |Q_{h_1, h_2} y'_{h_1, h_2}|^2 dx dt \\ &\leq \int_0^\infty \int_\Omega a(\mathbf{x}) |y_t|^2 dx dt, \end{aligned} \quad (82)$$

which combined with the weak convergence proved in Theorem 3.1 yields

$$h_1 h_2 \sum_{j=0}^J \sum_{k \in I_j} a_{j,k}(y'_{j,k})^2 \rightarrow \int_\Omega a(\mathbf{x}) |y_t|^2 dx \text{ strongly in } L^2(0, \infty). \quad (83)$$

Consequently,

$$h_i P_{h_1, h_2} y'_{h_1, h_2} \rightarrow 0 \text{ strongly in } L^2(0, +\infty; H_0^1(\Omega)), \quad i = 1, 2. \quad (84)$$

Moreover, thanks to

$$E(0) = E(t) + \int_0^t \int_\Omega a(\mathbf{x}) |y_t|^2 dx dt, \quad \forall t > 0, \quad (85)$$

we have

$$\begin{aligned} |E_{h_1, h_2}(t) - E(t)| &\leq |E_{h_1, h_2}(0) - E(0)| + h_1 h_2 \int_0^t \int_\Omega |\nabla P_{h_1, h_2} y'_{h_1, h_2}|^2 dx dt \\ &\quad + \left| h_1 h_2 \sum_{j=0}^J \sum_{k \in I_j} \int_0^t a_{j,k}(y'_{j,k})^2 dt - \int_0^t \int_\Omega a(\mathbf{x}) |y_t|^2 dx dt \right| \\ &\leq |E_{h_1, h_2}(0) - E(0)| + h_1 h_2 \int_0^\infty \int_\Omega |\nabla P_{h_1, h_2} y'_{h_1, h_2}|^2 dx dt \\ &\quad + \int_0^t \left| h_1 h_2 \sum_{j=0}^J \sum_{k \in I_j} a_{j,k}(y'_{j,k})^2 - \int_\Omega a(\mathbf{x}) |y_t|^2 dx \right| dt \\ &\leq |E_{h_1, h_2}(0) - E(0)| + h_1 h_2 \int_0^\infty \int_\Omega |\nabla P_{h_1, h_2} y'_{h_1, h_2}|^2 dx dt \\ &\quad + \int_0^\infty \left| h_1 h_2 \sum_{j=0}^J \sum_{k \in I_j} a_{j,k}(y'_{j,k})^2 - \int_\Omega a(\mathbf{x}) |y_t|^2 dx \right| dt \end{aligned} \quad (86)$$

and (78) follows.

4. NUMERICAL EXPERIMENTS

The aim of this section is to present some simple numerical experiments in order to confirm the theoretical results that indicate the need and efficiency of adding an extra viscous damping term. This is done over a fully-discrete approximation.

A rigorous analysis of the results in this paper to the fully discrete case remains to be done. However, the recent results in [17, 19] on the discrete Ingham inequality and its applications to fully-discrete approximation of the 1-D wave equation suggest that, very likely, the viscous term will suffice to guarantee the uniform decay property to hold in this case too.

4.1. Description of a fully discrete finite-difference scheme

We describe a fully discrete - in space and also in time - scheme associated to system (27). In order to simplify the formulation, we assume now that $h_1 = h_2 \equiv h$. Let us denote by $y_{j,k}^n$ the approximation of the solution y of (27) at the point of coordinates $(x_{1,j}, x_{2,k})$ and at time $n\Delta t$:

$$y_{j,k}^n \approx y(x_{1,j}, x_{2,k}, n\Delta t). \quad (87)$$

Here and in the sequel Δt denotes the time-step and n a nonnegative integer in the set $\{0, N\}$. N and Δt are defined such that $T = N\Delta t$. Therefore, we consider the following finite-difference approximation for the time derivative :

$$\frac{\partial^2 y}{\partial t^2}(\mathbf{x}, t) \approx \frac{y(\mathbf{x}, t + \Delta t) - 2y(\mathbf{x}, t) + y(\mathbf{x}, t - \Delta t)}{(\Delta t)^2}, \quad \frac{\partial y}{\partial t}(\mathbf{x}, t) \approx \frac{y(\mathbf{x}, t + \Delta t) - y(\mathbf{x}, t - \Delta t)}{2\Delta t}. \quad (88)$$

Then, we introduce the following fully discrete approximation of (27):

$$\begin{cases} (M_1(y^{n+1}))_{j,k} = (M_2(y^n))_{j,k} + (M_3(y^{n-1}))_{j,k}, \forall (j, k) \in \Omega_{\mathbf{h}_1, \mathbf{h}_2}, \forall n \in [0, N] \\ y_{j,k}^n = 0, \quad \forall (j, k) \in \partial\Omega_{\mathbf{h}_1, \mathbf{h}_2}, \quad \forall n \in [0, N], \end{cases} \quad (89)$$

with

$$\begin{cases} (M_1(y))_{j,k} \equiv (1 + a_{j,k} \frac{\Delta t}{2})y_{j,k} - \frac{\Delta t}{2}(M(y))_{j,k} \\ (M_2(y))_{j,k} \equiv 2y_{j,k} + r^2(M(y))_{j,k} \\ (M_3(y))_{j,k} \equiv (-1 + a_{j,k} \frac{\Delta t}{2})y_{j,k} - \frac{\Delta t}{2}(M(y))_{j,k} \\ (M(y))_{j,k} = y_{j+1,k} + y_{j-1,k} + y_{j,k+1} + y_{j,k-1} - 4y_{j,k} \end{cases} \quad (90)$$

and $r = \Delta t/h$. To initialize (89), we use

$$y_{j,k}^0 = y_0(x_{1,j}, x_{2,k}), \quad \frac{y_{j,k}^1 - y_{j,k}^{-1}}{2\Delta t} = y_1(x_{1,j}, x_{2,k}), \quad \forall (j, k) \in \partial\Omega_{\mathbf{h}_1, \mathbf{h}_2}. \quad (91)$$

Writing the first relation of (89) for $n = 0$, we obtain that the approximation of the solution y at time Δt is

$$y_{j,k}^1 = \frac{1}{2} \left(M_2(y_0(x_{1,j}, x_{2,k})) - 2\Delta t M_3(y_1(x_{1,j}, x_{2,k})) \right) \quad (92)$$

Remark 4.1.

- In the absence of damping and viscous terms, the scheme above reduces to $(M_1(y))_{j,k} = (M_3(y))_{j,k} = y_{j,k}$ which coincides with the classical finite difference (explicit) scheme for the wave equation.
- Note that M_1 is not a diagonal matrix because of the presence of the viscous damping term. Therefore, the scheme (89) is implicit.

Proposition 4.2. *The implicit scheme (89) is consistent of order 2 in time and space and is stable under the CFL condition $\Delta t \leq h/\sqrt{2}$. Therefore, the scheme is convergent.*

Proof. The first assertion is a direct consequence of (88). Concerning the stability condition, let us assume that the function $a(\mathbf{x}) \equiv a$ is constant in Ω and then perform a Von Neumann analysis. We assume that $y_{j,k}^n = \hat{y}^n e^{i(j j_{x_1} + k k_{x_2})\pi h}$, with $j_{x_1} \in [1, J], k_{x_2} \in [1, K]$ and replace this expression in (89). We obtain the following relation

$$\hat{y}^{n+1} \left(1 + C \frac{\Delta t}{2} \right) + \hat{y}^n \left(r^2 \xi^2 - 2 \right) + \hat{y}^{n-1} \left(1 - C \frac{\Delta t}{2} \right) = 0 \quad (93)$$

with

$$C \equiv a + \xi^2; \quad \xi^2 \equiv 4 \left(\sin^2(k_{x_1} \pi h/2) + \sin^2(j_{x_2} \pi h/2) \right). \quad (94)$$

According to the Von Neumann analysis, the scheme (89) is stable if the modulus of each root of the following polynomial is less than one ;

$$X^2 \left(1 + C \frac{\Delta t}{2} \right) + X \left(r^2 \xi^2 - 2 \right) + \left(1 - C \frac{\Delta t}{2} \right) = 0. \quad (95)$$

This is the case if

$$2r^2 \xi^2 (-4 + r^2 \xi^2) - 4(a + \xi^2) \Delta t \leq 0. \quad (96)$$

Since $a \geq 0$, this inequality holds if $r^2 \xi^2 \leq 4$, i.e., if $r^2 \leq 1/2$ which leads to the condition $\Delta t \leq h/\sqrt{2}$. From the Lax theorem (see [5]), we conclude that the scheme (89) is convergent. \square

Remark 4.3. • The case where $a(\mathbf{x})$ is non constant can be treated using an energy method ([5]). This leads to the same CFL condition.

- Using the approximation $\frac{\partial y}{\partial t}(\mathbf{x}, t) \approx (y(\mathbf{x}, t) - y(\mathbf{x}, t - \Delta t))/\Delta t$ instead of (88) leads to the following explicit scheme :

$$\begin{cases} y_{j,k}^{n+1} = (M_2(y^n))_{j,k} + (M_3(y^{n-1}))_{j,k}, & (j, k) \in \Omega_{h_1, h_2}, \forall n \in [0, N] \\ y_{j,k}^n = 0, & \forall (j, k) \in \partial \Omega_{h_1, h_2}, \quad \forall n \in [0, N], \end{cases} \quad (97)$$

with

$$\begin{cases} (M_2(y))_{j,k} \equiv (2 - \Delta t a_{j,k}) y_{j,k} + (r^2 + \Delta t) (M(y))_{j,k} \\ (M_3(y))_{j,k} \equiv (-1 + \Delta t a_{j,k}) y_{j,k} - (M(y))_{j,k} \end{cases} \quad (98)$$

and (92) being replaced by

$$y_{j,k}^1 = y_0(x_{1,j}, x_{2,k}) + \Delta t y_1(x_{1,j}, x_{2,k}). \quad (99)$$

However, this scheme is only of order one in time and is stable under a more restrictive CFL condition depending on a .

4.2. Spectrum of the discrete damped wave operator

In this section, we study the influence of the viscous terms on the spectrum of the discrete damped wave operator

$$\mathcal{A}_h^{(p)} = \begin{pmatrix} 0 & \mathbf{I}_{h,\Omega} \\ \Delta_h & -a \mathbf{I}_{h,\omega} + p h^2 \Delta_h \end{pmatrix} \quad (100)$$

We designate by $\sigma(\mathcal{A}_h^{(p)})$ this spectrum and we recall that - at the continuous level - the exponential decay rate verifies the inequality: $\alpha \leq -\sup\{Re(\lambda), \lambda \in \sigma(\mathcal{A})\}$. When $p = 1$, the viscous terms are used. We consider the geometry of the figure 3 included in the unit box (see also Figure 4). The domain ω is represented by Figure 4.

For $a(\mathbf{x}) = 10 \mathbb{1}_\omega(\mathbf{x})$ and $h = 1/20$, Figures 5 depict the discrete spectrum $\sigma(\mathcal{A}_h^{(p)})$ for $p = 0$ and $p = 1$. As well-known, ω being fixed, the profile of the spectrum strongly depends on the function a . However, whatever a , we observe that without viscosity terms, the spectral abscissa $-\max\{Re(\lambda), \lambda \in \sigma(\mathcal{A}_h^{(0)})\}$ goes to zero with h . On the contrary, this quantity remains uniformly bounded by below when the viscosity terms are used (see Table 2). This clearly indicates the necessity of introducing viscosity term in order to restore the uniform exponential decay property. Moreover, in agreement with the Remark 2.2, we observe any difference on the spectral abscissa with and without viscosity term when the damping function is effective on the whole domain, i.e. when $\omega = \Omega$. We obtain $-\max\{Re(\lambda), \lambda \in \sigma(\mathcal{A}_h^{(p)})\} \geq 5$ for all $h > 0$ and $p = 0, 1$. Clearly, in this case, whatever the speed




FIGURE 4. Domain $\Omega \subset (0, 1)^2$ (left) and support $\omega \subset \Omega$ (right) of the damping function a (right).

of the wave packets of high frequencies, they reach the region ω in a finite time. Furthermore, we refer the reader to [1] for the computation of the spectrum for geometrically complex domain in \mathbb{R}^2 .




FIGURE 5. Location of the eigenvalues of $\mathcal{A}_h^{(0)}$ (left - without viscosity terms) and $\mathcal{A}_h^{(1)}$ (right - with viscosity terms).

	$h = 1/20$	$h = 1/40$	$h = 1/80$	$h = 1/160$
without viscosity terms	0.01532	0.00934	0.00312	0.00023
with viscosity terms	0.2321	0.1924	0.1902	0.1898

TABLE 2. $-\max\{Re(\lambda), \lambda \in \sigma(\mathcal{A}_h^{(p)})\}$ without ($p = 0$) and with ($p = 1$) viscosity terms.

4.3. Numerical examples

In this section, we present some numerical experiments for three different initial conditions. The first one concerns the simplest regular initial condition involving only the first component of the frequency mode (see eq. (101)). The second one is a very singular one involving a discontinuous initial condition y_0 . For these two cases, we consider the domain Ω - part of the unit square - described by Figure 4-left. Finally, we take $a(\mathbf{x}) = 10 \mathbf{1}_\omega(\mathbf{x})$

where $\omega \subset \Omega$ is described by Figure 4-right. The third example concerns a very oscillating function y_0 defined on the unit square.

4.3.1. Regular initial conditions

Let us designate by D the disc of center $(1/2, 1/2)$ and radius $\sqrt{0.02}$. We check that $D \subset \Omega \setminus \omega$. We then define the following initial conditions in $H_0^1(\Omega) \times L^2(\Omega)$:

$$y_0(x_1, x_2) = 100 \left[0.02 - (x_1 - 1/2)^2 - (x_2 - 1/2)^2 \right] \mathbb{1}_D(x_1, x_2), \quad y_1(x_1, x_2) = 0. \quad (101)$$

On Ω , the function y_0 is positive and continuous.

For this function and this geometry, we check - in the conservative case - the order 2 of the convergence of the scheme (89). Figure 6 depicts the relative error of the energy $|E_h(0) - E_{ref}(0)|/E_{ref}(0)$ with respect to h (in a log-log scale): we obtain $|E_h(0) - E_{ref}(0)| = O(h^{1.92})$. The value of reference $E_{ref}(0)$ is obtained with $h = 1/500$.




FIGURE 6. $\log(|E_h(0) - E_{ref}(0)|/E_{ref}(0))$ vs. $-\log(h)$.




FIGURE 7. $\log(E_h(t))$ vs. t , $h = 1/100$.

Figure 7 describes the evolution of the energy for the conservative case, the case with velocity damping and the one with both velocity damping and viscous term. The scheme clearly reproduces the expected properties: in the conservative case, E_h is constant in time whereas in the other cases, the decay rate is exponential. In the damped case (without viscous term), the energy E_h is initially constant during a small period: this period corresponds to the time needed for the wave to reach ω . We also observe that the viscous term has a low impact on the decay rate. The regularity of the function y_0 implies that $h^2 \Delta y$ is negligible with respect to the other terms in eq.(27): when h goes to zero, both decay rates converge to the same value (see Table 3).

	$h = 1/50$	$h = 1/100$	$h = 1/150$	$h = 1/200$
without viscosity terms	-1.16484	-1.19264	-1.19521	-1.19655
with viscosity terms	-1.23830	-1.20136	-1.19892	-1.19723

TABLE 3. Exponential decay rate observed in the time interval $[0, 3]$ vs. h .

Finally, Figure 8 depicts the evolution of the solution for $t \in [0, 9/4]$ obtained with $h = 1/100$, highlighting the damping mechanism.




FIGURE 8. Solution y_h at time $t = 0, 1/4, 2/4, 3/4, 4/4, 5/4, 6/4, 7/4, 8/4, 9/4$ - $E_h(0) \approx 11.98$, $E_h(1/4) \approx 11.38$, $E_h(2/4) \approx 8.83$, $E_h(3/4) \approx 5.56$, $E_h(4/4) \approx 3.95$, $E_h(5/4) \approx 3.18$, $E_h(6/4) \approx 2.41$, $E_h(7/4) \approx 1.72$, $E_h(8/4) \approx 1.21$, $E_h(9/4) \approx 0.95$.

4.3.2. Irregular initial conditions

The previous section does not highlight the necessity of the viscous term. With the same geometry, we now consider the discontinuous initial conditions :

$$y_0(\mathbf{x}) = 2 \mathbb{1}_D(\mathbf{x}) \quad ; \quad y_1(\mathbf{x}) = 0. \quad (102)$$

In this case, the viscous term $h^2 \Delta y$ is not negligible. Figure 9 and Figure 10 represent the evolution of E_h for $t \in [0, 3]$ with and without extra viscous terms with mesh size $h = 1/100$. In both cases, we observe an exponential decay rate. However, contrary to the first example, the decay rates are different (see Table 4). The viscous term permits to damped out the high frequencies components of the solution enhanced by the irregularity of y_0 . Without viscous term, the time needed for all the wave packets to reach ω is larger (and dependent of h). After this time (large enough), the energy in both cases are similar. Finally, in order to appreciate the regularizing effect of the term $h^2 \Delta y$, Figure 11 depicts in both cases the solution $\mathbf{y}_h(x_1, 0.4)$ at time $t = 2$.




FIGURE 9. $E_h(t)$ vs. $t \in [0, 6]$, $h = 1/100$.




FIGURE 10. $\log(E_h(t))$ vs. $t \in [0, 2]$, $h = 1/100$.

	$h = 1/50$	$h = 1/100$	$h = 1/150$	$h = 1/200$
without viscosity terms	-0.94462	-0.92221	-0.90134	-0.90012
with viscosity terms	-1.84121	-2.00718	-2.13322	-2.155835

TABLE 4. Exponential decay rate observed in the time interval $[0, 2]$ vs. h .

4.3.3. Influence of the high frequencies

In the extreme case where the solution y is only represented by high frequencies components, the term $h^2 \Delta y$ becomes preponderant. On the unit square, let us now consider the very oscillating initial condition (but still continuous) :

$$y_0(\mathbf{x}) = \sin(-\pi x_1 \frac{1}{h}) \sin(-\pi x_2 \frac{1}{h}) - \sin(\pi x_1 (1 - \frac{1}{h})) \sin(\pi x_2 (1 - \frac{1}{h})) \quad ; \quad y_1(\mathbf{x}) = 0 \quad (103)$$




FIGURE 11. $\mathbf{y}_h(x_1, 0.4, t = 2)$ vs. $x_1 \in [0, 1]$ for $h = 1/100$ with and without extra viscous terms.

and $a(\mathbf{x}) \equiv 20 \mathbb{1}_\omega(\mathbf{x})$ where $\omega \equiv \{x_1 \in [0.9, 1], x_2 \in [0, 1]\} \cup \{x_1 \in [0, 1], x_2 \in [0.9, 1]\}$. These initial conditions lead to wave packets with group velocity of the order of the mesh size h and they are the main obstacle for the uniform stabilization property to hold. We observe a clear difference on the behavior of the solution \mathbf{y}_h obtained with or without viscous damping. Without extra viscous terms, the velocity damping has no effect: computations lead to a rate of decay slowly converging toward zero when h goes to zero. On the contrary, when the viscous term is added, we obtain a rate of decay near -7 (see figure 12 and Table 5). This is also illustrated in Figures 13 where we draw the evolution of $\mathbf{y}_h(x_1, 1/2, t)$ for $t \in [0, 2]$. Finally, it is worth mentioning that the use of the viscous scheme leads to a significant increase of the computational cost.




FIGURE 12. $\log(E_h(t))$ vs. t ; $h = 1/60$.




FIGURE 13. $y_h(x_1, 1/2, t)$ without (left) and with (right) extra viscous terms for $t = 0.8, 1.2, 1.6, 2$.

	$h = 1/20$	$h = 1/40$	$h = 1/60$	$h = 1/80$	$h = 1/100$
without viscosity terms	-0.4256	-0.4042	-0.3911	-0.3815	-0.3700
with viscosity terms	-5.8714	-6.4738	-6.7190	-6.8379	-6.9098

TABLE 5. Exponential decay rate observed in the time interval $[0, 2]$ vs. h .

Acknowledgements.

We would like to thank the Referee of this journal for the constructive criticism which allow us to improve our earlier version. Professor Enrique Zuazua is also gratefully acknowledged for fruitful discussions and valuable comments on preliminary version of this work .

REFERENCES

- [1] M. Asch, G. Lebeau, *The spectrum of the damped wave operator for geometrically complex domain in \mathbb{R}^2* , Experimental Mathematics **12(2)** (2003).
- [2] H. T. Banks, K. Ito, B. Wang, *Exponentially stable approximations of weakly damped wave equations*, Ser. Num. Math. **100**, 1-33, Birkhäuser (1990).
- [3] C. Bardos, G. Lebeau, J. Rauch, *Sharp sufficient conditions for the observation, control and stabilization from the boundary*, SIAM J. Control and Opt., **30**, 1024-1065 (1992).
- [4] D. Chenais, *On the existence of a solution in a domain identification problem*, J. Math. Anal. Appl., **52**, 189-219 (1975).
- [5] G.C. Cohen, *Higher-order Numerical Methods for Transient Wave Equations*, Scientific Computation, Springer, (2002).
- [6] C. M. Dafermos, *On the existence and the asymptotic stability of solutions to the equations of linear thermoelasticity*, Arch. Rational Mech. Anal., **29**, 241-271 (1968).
- [7] R. Glowinski, C. H. Li, J.-L. Lions, *A numerical approach to the exact boundary controllability of the wave equation (I). Dirichlet Controls: Description of the numerical methods*, Japan. J. Appl. Math., **7**, 1-76 (1990).
- [8] A. Haraux, *Une remarque sur la stabilisation de certains systèmes du deuxième ordre en temps*, Portug. Math., **46**, 245-258 (1989).
- [9] A. Haraux, *Stabilization of trajectories for some weakly damped hyperbolic equations*, J. Differential Equations, **59**, 145-154 (1985).
- [10] A. Henrot, *Continuity with respect to the domain for the Laplacian : a survey*, Control and Cybernetics, **23(3)**, 427-443 (1994).
- [11] J.A. Infante and E. Zuazua, *Boundary observability for the space-discretizations of the 1-D wave equation*, Math. Model. Num. An. **33**, 407-438 (1999).
- [12] V. Komornik, *Exact controllability and Stabilization - The multiplier method*, J. Wiley and Masson (1994).
- [13] S. Krenk, *Dispersion-corrected explicit integration of the wave equation*, Comput. Methods. Appl. Mech. Engrg., **191**, 975-987 (2001).
- [14] J.L. Lions, *Quelques méthodes de résolution des problèmes aux limites non linéaires*, Dunod-Gauthier-Villars, Paris (1969).
- [15] J.L. Lions and E. Magenes, *Problèmes aux limites non homogènes et applications, vol. 1*, Dunod, Paris (1968).
- [16] J. Lagnese, *Control of wave processes with distributed control supported on a subregion*, SIAM J. Control and Opt., **21**, 68-85 (1983).
- [17] A. Münch, *A uniformly controllable and implicit scheme for the 1-D wave equation*, Mathematical Modelling and Numerical Analysis, **39(2)**, 377-418, (2005).
- [18] M. Nakao, *Decay of solutions of the wave equation with a local degenerate dissipation*, Israel J. Math., **95**, 25-42 (1996).
- [19] M. Negreanu and E. Zuazua, *Discrete Ingham inequalities and applications*, C.R. Acad. Sci. Paris, **338**, 281-286 (2004).
- [20] O. Pironneau, *Optimal shape design for elliptic systems*, New York, Springer (1984).
- [21] K. Ramdani, T. Takahashi and M. Tucsnak, *Uniformly exponentially stable approximations for a class of second order evolution equations: Application to the optimal controle of flexible structures*, Technical report, Prépublications de l'Institut Elie Cartan, 27 (2003).
- [22] L.R. Tcheugoué-Tébou, *Stabilization of the wave equation with localized nonlinear damping*, Journal of Differential Equations **145**, 502-524 (1998).

- [23] L.R. Tcheugoué-Tébou and E. Zuazua, *Uniform exponential long time decay for the space semi-discretization of a locally damped wave equation via an artificial numerical viscosity*, Numer. Math. **95(3)**, 563-598 (2003).
- [24] M. Slemrod, *Weak asymptotic decay via a "Relaxed Invariance Principle" for a wave equation with nonlinear, non-monotone damping*, Proc. Royal Soc. Edinburgh, **113**, 87-97 (1989).
- [25] E. Zuazua, *Exponential decay for the semilinear wave equation with locally distributed damping*, Comm. in Partial Differential Equation **15**, 205-235 (1990).
- [26] E. Zuazua, *Boundary observability for finite-difference space semi-discretizations of the 2-D wave equation in the square*, J. Math. Pures. Appl. **78**, 523-563 (1999).
- [27] E. Zuazua, *Propagation, Observation, Control and Numerical Approximation of Waves*, SIAM Review **47(2)**, 197-243 (2005).