

COMPLEX ANALOGUES OF THE HALF-CLASSICAL GEOMETRY

TEODOR BANICA AND JULIEN BICHON

ABSTRACT. Under very strong axioms, there is precisely one real noncommutative geometry between the classical one and the free one, namely the half-classical one, coming from the relations $abc = cba$. We discuss here the complex analogues of this geometry, notably with a study of the geometry coming from the commutation relations between all the variables $\{ab^*, a^*b\}$, that we believe to be the “correct” one.

INTRODUCTION

The present paper is motivated by some rigidity phenomena appearing in real non-commutative algebraic geometry. It is known indeed that, when imposing the strongest possible axioms, there are only three such geometries, namely the classical one, the free one, and an intermediate one, called “half-classical”. Motivated by this fact, we started in [4] an investigation of the possible half-classical complex geometries. In this paper we will finish the work started there, by identifying the “standard” such geometry.

Let us first explain the above-mentioned rigidity phenomenon, from the real case. Consider the unit sphere, the standard cube, and the orthogonal group:

$$\begin{aligned} S_{\mathbb{R}}^{N-1} &= \left\{ x \in \mathbb{R}^N \mid \sum_i x_i^2 = 1 \right\} \\ T_N &= \left\{ x \in \mathbb{R}^N \mid x_i = \pm \frac{1}{\sqrt{N}} \right\} \\ O_N &= \left\{ U \in M_N(\mathbb{R}) \mid U^t = U^{-1} \right\} \end{aligned}$$

Quite remarkably, there is a full set of connections between these objects:

- (1) O_N is the isometry group of $S_{\mathbb{R}}^{N-1}$, and $S_{\mathbb{R}}^{N-1}$ appears as $O_N(\circ)$, where $\circ = (1, 0, \dots, 0)$. In addition, we have an embedding $O_N \subset \sqrt{N} \cdot S_{\mathbb{R}}^{N^2-1}$.
- (2) T_N appears inside $S_{\mathbb{R}}^{N-1}$ by setting $|x_1| = \dots = |x_N|$. Conversely, $S_{\mathbb{R}}^{N-1}$ appears from $T_N \subset \mathbb{R}^N$ by “deleting” this relation, while still keeping $\sum_i x_i^2 = 1$.
- (3) $T_N \simeq \mathbb{Z}_2^N$ is a maximal compact abelian subgroup of O_N , and the group O_N itself can be reconstructed from this subgroup, by using various methods.

2010 *Mathematics Subject Classification.* 14A22 (16S38, 46L65).

Key words and phrases. Quantum isometry, Noncommutative manifold.

In order to construct noncommutative versions of the above objects, we use the operator algebra formalism. The idea indeed is that whenever we have a unital C^* -algebra A , we can write $A = C(X)$, with X being a noncommutative compact space. This is supported by a non-trivial theorem of Gelfand, which states that when A is commutative, the formula $A = C(X)$ holds indeed, with X being a classical space, called spectrum of A .

So, let us define the free sphere, cube, and orthogonal group, by setting:

$$\begin{aligned} C(S_{\mathbb{R},+}^{N-1}) &= C^* \left((x_i)_{i=1,\dots,N} \mid x_i = x_i^*, \sum_i x_i^2 = 1 \right) \\ C(T_N^+) &= C^* \left((x_i)_{i=1,\dots,N} \mid x_i = x_i^*, x_i^2 = \frac{1}{N} \right) \\ C(O_N^+) &= C^* \left((u_{ij})_{i,j=1,\dots,N} \mid u_{ij} = u_{ij}^*, u^t = u^{-1} \right) \end{aligned}$$

Observe that $u_i = \sqrt{N}x_i \in C(T_N^+)$ are subject to the relations $u_i = u_i^* = u_i^{-1}$. Thus, T_N^+ appears as abstract dual of the discrete group \mathbb{Z}_2^{*N} . As for O_N^+ , this is a compact quantum group in the sense of Woronowicz [25], with structural maps as follows:

$$\Delta(u_{ij}) = \sum_k u_{ik} \otimes u_{kj} \quad , \quad \varepsilon(u_{ij}) = \delta_{ij} \quad , \quad S(u_{ij}) = u_{ji}$$

In analogy with what happens in the classical case, we have:

- (1) O_N^+ is the quantum isometry group of $S_{\mathbb{R},+}^{N-1}$, and $S_{\mathbb{R},+}^{N-1}$ appears as an homogeneous space over O_N^+ . In addition, we have an embedding $O_N^+ \subset \sqrt{N} \cdot S_{\mathbb{R},+}^{N^2-1}$.
- (2) T_N^+ appears inside $S_{\mathbb{R},+}^{N-1}$ by setting $x_1^2 = \dots = x_N^2$. Conversely, $S_{\mathbb{R},+}^{N-1}$ appears from T_N^+ by “deleting” this relation, while still keeping $\sum_i x_i^2 = 1$.
- (3) $T_N^+ \simeq \widehat{\mathbb{Z}_2^{*N}}$ is a maximal group dual subgroup of O_N^+ , and O_N^+ itself can be reconstructed from this subgroup, via representation theory methods.

We refer to [6], [7], [23], [24] and to the lecture notes [3] for details.

Let us try now to understand the possible “intermediate liberations”. At the sphere and cube level, there is a lot of freedom in dealing with this question. However, at the quantum group level, things are quite rigid. So, as a first good question, we would like to find the intermediate quantum groups, as follows:

$$O_N \subset G \subset O_N^+ \quad (*)$$

In order to deal with this problem, let us recall Brauer’s theorem [14]. Given a pairing $\pi \in P_2(k, l)$, between an upper row of k points, and a lower row of l points, we set:

$$T_\pi(e_{i_1} \otimes \dots \otimes e_{i_k}) = \sum_{j_1 \dots j_l} \delta_\pi \begin{pmatrix} i_1 & \dots & i_k \\ j_1 & \dots & j_l \end{pmatrix} e_{j_1} \otimes \dots \otimes e_{j_l}$$


The theorem states that the intertwining spaces $Hom(u^{\otimes k}, u^{\otimes l})$ for the orthogonal group O_N are precisely those spanned by these maps T_π . In addition, a free version of this result is available, stating that for O_N^+ , the intertwining spaces are spanned as well by the maps T_π , but this time with π being a noncrossing pairing, $\pi \in NC_2$. See [6].

Based on these results, let us call a quantum group $O_N \subset G \subset O_N^+$ easy when the following holds, for a certain category of pairings, $NC_2 \subset D \subset P_2$:

$$Hom(u^{\otimes k}, u^{\otimes l}) = span \left(T_\pi \mid \pi \in D(k, l) \right)$$

Here the categorical operations are the horizontal and vertical concatenation, and the upside-down turning of the pairings. These operations ensure the fact that the spaces $span \left(T_\pi \mid \pi \in D(k, l) \right)$ form a tensor C^* -category, and as a consequence of Woronowicz's Tannakian duality [26], each such category produces a quantum group. See [8], [18].

We are now ready to go back to (*). If we restrict attention to the easy case, we just have to find the intermediate categories $NC_2 \subset D \subset P_2$. And here, there is only one solution, namely the category P_2^* generated by the following “crossing”:



Due to the uniqueness result, we will call this diagram “half-classical crossing”, and the corresponding relation, namely $abc = cba$, “half-commutation relation”. See [9].

Summarizing, we have now an answer to (*), which is in addition unique, in the easy quantum group setting. We should mention that, conjecturally, this solution is actually unique, in the arbitrary compact quantum group setting. See [5].

So, let us go ahead now, and construct our various geometric objects, as follows:

$$\begin{aligned} C(S_{\mathbb{R},*}^{N-1}) &= C(S_{\mathbb{R},+}^{N-1}) / \langle abc = cba \mid \forall a, b, c \in \{x_i\} \rangle \\ C(T_N^*) &= C(T_N^+) / \langle abc = cba \mid \forall a, b, c \in \{x_i\} \rangle \\ C(O_N^*) &= C(O_N^+) / \langle abc = cba \mid \forall a, b, c \in \{u_{ij}\} \rangle \end{aligned}$$

As in the classical and free cases, we have correspondences between these objects, with statements (1-3) as above. For details here, we refer to [3], [7], [9].

In view of the uniqueness results mentioned above, we can stop here the axiomatization work, because this is the third and last possible “geometry”. So, what is left to do now is to start the actual geometric work. We would like to understand the structure and geometry of the various algebraic manifolds $X \subset S_{\mathbb{R},*}^{N-1}, X \subset S_{\mathbb{R},+}^{N-1}$. We have as well the important question of finding explicit matrix models for the coordinates of such manifolds. The whole subject here is still very young, and we refer to [2], [4], [12].

The aim of the present paper is to clarify what happens in the complex geometry setting. Available here are the classical theory, having symmetry group U_N , and its free version, with symmetry group U_N^+ . At the level of intermediate geometries, however, the situation is quite complicated, because the following question has many answers:

$$U_N \subset G \subset U_N^+ \quad (**)$$

Our goal here will be not that of solving this problem, but rather of trying to identify the “main” solution. For this purpose, we will use an axiomatic approach. We will first axiomatize the triples (S, T, G) which are subject to correspondences (1-3) as above, and then we will examine the various complex analogues of the triple $(S_{\mathbb{R},*}^{N-1}, T_N^*, O_N^*)$.

The paper is organized as follows: 1-2 contain preliminaries and generalities, and in 3-4 we construct the complex half-classical geometry.

1. FORMALISM

We are interested here in the noncommutative analogues of the real algebraic manifolds $X \subset S_{\mathbb{C}}^{N-1}$. Our starting point will be the following well-known fact:

Proposition 1.1. *Consider a real algebraic manifold $X \subset S_{\mathbb{C}}^{N-1}$, appearing as:*

$$X = \left\{ x \in \mathbb{C}^N \mid \sum_i |x_i|^2 = 1, P_\alpha(x_1, \dots, x_N) = 0 \right\}$$

The algebra of continuous functions $f : X \rightarrow \mathbb{C}$ is then given by

$$C(X) = C_{comm}^* \left(x_1, \dots, x_N \mid \sum_i x_i x_i^* = 1, P_\alpha(x_1, \dots, x_N) = 0 \right)$$

where by C_{comm}^ we mean universal commutative C^* -algebra.*

Proof. Observe first that the universal algebra in the statement is well-defined, because $\sum_i x_i x_i^* = 1$ gives $\|x_i\| \leq 1$ for any i , and so the biggest norm is bounded. If we denote by A this algebra, we have an arrow $A \rightarrow C(X)$. Conversely, by Gelfand duality we have $A = C(X')$ for a certain compact space X' . The coordinates x_i produce an embedding $X' \subset \mathbb{C}^N$, then the condition $\sum_i x_i x_i^* = 1$ gives $X' \subset S_{\mathbb{C}}^{N-1}$, and finally the conditions $P_\alpha(x_1, \dots, x_N) = 0$ give $X' \subset X$. Thus we have $X = X'$, as claimed. \square

The above result suggests to construct a free version X^+ , simply by removing the commutativity assumption from the presentation of $C(X)$. However, this is quite tricky, because the relations $P_\alpha = 0$ must not include, or imply, the commutativity.

In practice, this method works in a number of situations. We have:

Definition 1.2. *The free complex sphere and free unitary group are constructed as*

$$\begin{aligned} C(S_{\mathbb{C},+}^{N-1}) &= C^* \left((x_i)_{i=1,\dots,N} \mid \sum_i x_i x_i^* = \sum_i x_i^* x_i = 1 \right) \\ C(U_N^+) &= C^* \left((u_{ij})_{i=1,\dots,N} \mid u^* = u^{-1}, u^t = \bar{u}^{-1} \right) \end{aligned}$$

where on the right we have universal C^* -algebras.

As explained by Wang in [23], the above noncommutative space U_N^+ is a compact quantum group in the sense of Woronowicz [25], with structural maps as follows:

$$\Delta(u_{ij}) = \sum_k u_{ik} \otimes u_{kj} \quad , \quad \varepsilon(u_{ij}) = \delta_{ij} \quad , \quad S(u_{ij}) = u_{ji}^*$$

We have an action $U_N^+ \curvearrowright S_{\mathbb{C},+}^{N-1}$, whose properties are quite similar to that of the action $U_N \curvearrowright S_{\mathbb{C}}^{N-1}$. In order to explain this material, let us introduce a few more notions:

Definition 1.3. *Consider an algebraic submanifold $X \subset S_{\mathbb{C},+}^{N-1}$, i.e. a closed subset defined via algebraic relations, and a closed quantum subgroup $G \subset U_N^+$.*

- (1) *We say that we have an affine action $G \curvearrowright X$ when the formula $x_i \rightarrow \sum_j u_{ij} \otimes x_j$ defines a morphism of C^* -algebras $\Phi : C(X) \rightarrow C(G) \otimes C(X)$.*
- (2) *The biggest quantum subgroup $G \subset U_N^+$ acting affinely on X is denoted $G^+(X)$, and is called quantum isometry group of X .*

Observe that the morphism in (1) is automatically coassociative, $(\Phi \otimes id)\Phi = (id \otimes \Delta)\Phi$, and counital as well, $(id \otimes \varepsilon)\Phi = id$. When X, G are both classical such a morphism must appear by transposition from a usual affine group action $G \times X \rightarrow X$.

Regarding (2), it is routine to check that such a biggest quantum group exists indeed, simply by dividing $C(U_N^+)$ by the appropriate relations. See [3], [16], [17].

We will need as well the following constructions:

Definition 1.4. *Consider a subspace $S \subset S_{\mathbb{C},+}^{N-1}$, and a subgroup $G \subset U_N^+$.*

- (1) *The standard torus of S is the subspace $T \subset S$ obtained by setting, at the algebra level, $C(T) = C(S) / \langle x_i x_i^* = x_i^* x_i = 1/N \rangle$.*
- (2) *The diagonal torus of G is the subspace $\mathcal{T} \subset G$ obtained by setting, at the algebra level, $C(\mathcal{T}) = C(G) / \langle u_{ij} = 0 \mid \forall i \neq j \rangle$.*

For $S = S_{\mathbb{C}}^{N-1}$, the space constructed in (1) is the usual torus, $T = \mathbb{T}^N$. For $S = S_{\mathbb{C},+}^{N-1}$, the rescaled generators $u_i = \sqrt{N}x_i$ are subject to the relations $u_i^* = u_i^{-1}$, which produce the free group algebra $C^*(F_N)$. Thus, we obtain here a group dual, $T = \widehat{F_N}$.

Regarding (2), observe that $C(\mathcal{T})$ is generated by the variables $g_i = u_{ii}$, which are group-like. Thus $\mathcal{T} = \widehat{\Gamma}$, where Γ is the group generated by these variables. See [9].

We can now formulate our main definition, as follows:

Definition 1.5. *A noncommutative geometry consists of three spaces,*

- (1) *an intermediate algebraic manifold $S_{\mathbb{R}}^{N-1} \subset S \subset S_{\mathbb{C},+}^{N-1}$, called sphere,*
- (2) *an intermediate space $\widehat{\mathbb{Z}}_2^N \subset T \subset \widehat{F}_N$, called torus,*
- (3) *and an intermediate quantum group, $O_N \subset G \subset U_N^+$,*

such that the following two conditions are satisfied,

- (1) *G is the quantum isometry group of S ,*
- (2) *T is the standard torus of S , as well as the diagonal torus of G ,*

where, for the needs of the second axiom, we identify full and reduced group algebras.

The torus T will be called as well ‘‘cube’’ of the geometry, because in the real case, as explained below, we rather have a cube. The discrete dual $\Gamma = \widehat{T}$, which appears as an intermediate group $F_N \rightarrow \Gamma \rightarrow \mathbb{Z}_2^N$, will be called structural group of the geometry.

Here is useful reformulation of the axioms, in terms of S only:

Proposition 1.6. *An algebraic manifold $S_{\mathbb{R}}^{N-1} \subset S \subset S_{\mathbb{C},+}^{N-1}$, with standard torus denoted $T \subset S$, produces a noncommutative geometry precisely when:*

- (1) *We have an affine action $O_N \curvearrowright S$.*
- (2) *$\delta(x_i) = \sqrt{N}x_i \otimes x_i$ defines a morphism of algebras $\delta : C(S) \rightarrow C(T) \otimes C(S)$.*

If these conditions are satisfied, we say that S is a noncommutative sphere.

Proof. Given $S_{\mathbb{R}}^{N-1} \subset S \subset S_{\mathbb{C},+}^{N-1}$, consider its quantum isometry group $G \subset U_N^+$, and consider as well the standard torus $T \subset S$, and the diagonal torus $\mathcal{T} \subset G$.

Assuming that (S, T, G) is a noncommutative geometry, we have $O_N \subset G$, so the condition (1) is clear. Also, since we have $T = \mathcal{T}$, the morphism in (2) simply appears by composing the universal coaction with the diagonal torus quotient map:

$$\begin{aligned} C(S) &\rightarrow C(G) \otimes C(S) \rightarrow C(\mathcal{T}) \otimes C(S) = C(T) \otimes C(S) \\ x_i &\rightarrow \sum_j u_{ij} \otimes x_j \rightarrow g_i \otimes x_i = \sqrt{N}x_i \otimes x_i \end{aligned}$$

Conversely, assuming that (1,2) are satisfied, we must prove that we have $T = \mathcal{T}$. For this purpose, observe first that the map δ induces a morphism as follows:

$$\bar{\delta} : C(T) \rightarrow C(T) \otimes C(T) \quad , \quad x_i \rightarrow \sqrt{N}x_i \otimes x_i$$

Thus $C(T)$ is a cocommutative Hopf algebra, and its elements $\sqrt{N}x_i$ are group-like.

With this picture in mind, the map δ in the statement corresponds to an action $T \curvearrowright S$, and by universality of the quantum isometry group G , we obtain a map as follows:

$$\theta : C(\mathcal{T}) \rightarrow C(T) \quad , \quad u_{ii} \rightarrow \sqrt{N}x_i$$

In order to construct an inverse map, we will use a composition, as follows:

$$\begin{aligned} C(S) &\rightarrow C(G) \otimes C(S) \rightarrow C(G) \rightarrow C(G) \rightarrow C(\mathcal{T}) \\ x_i &\rightarrow \sum_j u_{ij} \otimes x_j \rightarrow u_{i1} \rightarrow \frac{1}{\sqrt{N}} \sum_j u_{ij} \rightarrow \frac{1}{\sqrt{N}} u_{ii} \end{aligned}$$

Here the first map is the universal coaction, and the second map comes from the evaluation at $(1, 0, \dots, 0) \in S_{\mathbb{R}}^{N-1} \subset S$, which gives a map $\varepsilon : C(S) \rightarrow \mathbb{C}$, $x_i \rightarrow \delta_{i1}$. Regarding the third map, this is the algebra automorphism $u_{i1} \rightarrow \sum_k m_{k1} u_{ik}$ constructed by using $O_N \subset G$, and an orthogonal matrix $M = (m_{ij})$ having $\frac{1}{\sqrt{N}}$ entries on its first column. Finally, the fourth map is the canonical quotient map.

Now since the elements $X_i = \frac{1}{\sqrt{N}} u_{ii}$ satisfy the relations $X_i X_i^* = X_i^* X_i = \frac{1}{N}$, the composition that we constructed factorizes into a map as follows:

$$\rho : C(T) \rightarrow C(\mathcal{T}) \quad , \quad x_i \rightarrow \frac{1}{\sqrt{N}} u_{ii}$$

It is clear that θ, ρ are inverse to each other, and this finishes the proof. □

At the level of basic examples, we have:

Proposition 1.7. *We have the following examples of geometries:*

- (1) *Real geometry:* $S = S_{\mathbb{R}}^{N-1}$, $T = \widehat{\mathbb{Z}_2^N}$, $G = O_N$.
- (2) *Complex geometry:* $S = S_{\mathbb{C}}^{N-1}$, $T = \widehat{\mathbb{Z}^N}$, $G = U_N$.
- (3) *Free real geometry:* $S = S_{\mathbb{R},+}^{N-1}$, $T = \widehat{\mathbb{Z}_2^{*N}}$, $G = O_N^+$.
- (4) *Free complex geometry:* $S = S_{\mathbb{C},+}^{N-1}$, $T = \widehat{F_N}$, $G = U_N^+$.

Proof. These results are well-known, and we refer to [1], [7] for details here. □

Based on these examples, here are now a few more general notions:

Definition 1.8. *A noncommutative geometry is called:*

- (1) *Real*, if $S \subset S_{\mathbb{R},+}^{N-1}$, $T \subset \widehat{\mathbb{Z}_2^{*N}}$, $G \subset O_N^+$.
- (2) *Complex*, if $S_{\mathbb{C}}^{N-1} \subset S$, $\widehat{\mathbb{Z}^N} \subset T$, $U_N \subset G$.
- (3) *Classical*, if $S \subset S_{\mathbb{C}}^{N-1}$, $T \subset \widehat{\mathbb{Z}^N}$, $G \subset U_N$.
- (4) *Free*, if $S_{\mathbb{R},+}^{N-1} \subset S$, $\widehat{\mathbb{Z}_2^{*N}} \subset T$, $O_N^+ \subset G$.

A geometry which is not real, nor complex, will be called hybrid. A geometry which is not classical, nor free, will be called intermediate. Examples here come from:

Definition 1.9. *We have spheres, tori, and quantum groups, as follows:*

- (1) $S_{\mathbb{R},*}^{N-1}, \widehat{\mathbb{Z}_2^{*N}}, O_N^*$, obtained respectively from $S_{\mathbb{R},+}^{N-1}, \widehat{\mathbb{Z}_2^{*N}}, O_N^+$ by imposing to the standard coordinates the relations $abc = cba$.
- (2) $S_{\mathbb{C},**}^{N-1}, \widehat{\mathbb{Z}^{\circ N}}, U_N^{**}$, obtained respectively from $S_{\mathbb{C},+}^{N-1}, \widehat{F_N}, U_N^+$ by imposing to the standard coordinates and their adjoints the relations $abc = cba$.

Here (1) is a well-established definition, coming from the work in [8], [9]. Regarding (2), we have there some temporary objects, coming from [13], which are somehow “minimal”, and which will be replaced with the correct ones, later on.

We can now formulate our first result, as follows:

Theorem 1.10. *We have geometries, whose unitary groups are as follows,*

$$\begin{array}{ccccc}
U_N & \longrightarrow & U_N^{**} & \longrightarrow & U_N^+ \\
\uparrow & & \uparrow & & \uparrow \\
\mathbb{T}O_N & \longrightarrow & \mathbb{T}O_N^* & \longrightarrow & \mathbb{T}O_N^+ \\
\uparrow & & \uparrow & & \uparrow \\
O_N & \longrightarrow & O_N^* & \longrightarrow & O_N^+
\end{array}$$

with the middle row spaces obtained from the upper ones via the relations $ab^* = a^*b$.

Proof. The results in the upper and lower row are well-known, see [1]. Regarding the middle row, consider the following quotient algebra:

$$C(\mathbb{T}O_N^+) = C(U_N^+) / \left\langle ab^* = a^*b \mid \forall a, b \in \{u_{ij}\} \right\rangle$$

Inside this algebra, observe that with $U_{ij} = \sum_a u_{ia} \otimes u_{aj}$ we have:

$$U_{ij}U_{kl}^* = \sum_{ab} u_{ia}u_{kb}^* \otimes u_{aj}u_{bl}^* = \sum_{ab} u_{ia}^*u_{kb} \otimes u_{aj}^*u_{bl} = U_{ij}^*U_{kl}$$

Thus we can construct a comultiplication morphism Δ , by mapping $u_{ij} \rightarrow U_{ij}$, and the existence of the counit ε and of the antipode S is clear too. Now with $\mathbb{T}O_N^+$ constructed as above, we can construct the other quantum groups as well, as follows:

$$\mathbb{T}O_N = \mathbb{T}O_N^+ \cap U_N \quad , \quad \mathbb{T}O_N^* = \mathbb{T}O_N^+ \cap U_N^*$$

For the spheres and tori, the discussion here parallels the one from the quantum group case. To be more precise, the definition of these objects is as follows:

$$\begin{aligned}
C(\mathbb{T}S_{\mathbb{R}, \times}^{N-1}) &= C(S_{\mathbb{C}, \times}^{N-1}) / \left\langle ab^* = a^*b \mid \forall a, b \in \{x_i\} \right\rangle \\
C(\widehat{\mathbb{T}\mathbb{Z}_2^{\times N}}) &= C(\widehat{\mathbb{Z}_2^{\times N}}) / \left\langle ab^* = a^*b \mid \forall a, b \in \{g_i\} \right\rangle
\end{aligned}$$

Regarding the axioms, let us prove now that the standard action $U_N^\times \curvearrowright S_{\mathbb{C}, \times}^{N-1}$ restricts to an action $\mathbb{T}O_N^\times \curvearrowright \mathbb{T}S_{\mathbb{R}, \times}^{N-1}$. With $X_i = \sum_a u_{ia} \otimes x_a$ we have:

$$X_i X_j^* = \sum_{ab} u_{ia}u_{jb}^* \otimes x_a x_b^* = \sum_{ab} u_{ia}^*u_{jb} \otimes x_a^* x_b = X_i^* X_j$$

Thus we can indeed define our coaction map, via $x_i \rightarrow X_i$. In order to prove now the universality, assume that we have an action $G \curvearrowright \mathbb{T}S_{\mathbb{R}, \times}^{N-1}$. With $X_i = \sum_a u_{ia} \otimes x_a$ as above we have $X_i X_j^* = X_i^* X_j$, and from this we obtain, via the symmetrization trick from [11],

that we have $u_{ia}u_{jb}^* = u_{ia}^*u_{jb}$ for any i, j, a, b , and so $G \subset \text{TO}_N^\times$. We refer here to [1] for the classical case, the proof in the half-classical and free cases being similar. \square

2. EASINESS, AMENABILITY

We will need the notion of easy quantum group, from [8], [15], [21], [22].

We denote by $P(k, l)$ the set of partitions between an upper row of k points, and a lower row of l points, with each leg colored black or white, and with k, l standing for the corresponding ‘‘colored integers’’. We have the following notion:

Definition 2.1. *A category of partitions is a collection of sets $D = \bigcup_{kl} D(k, l)$, with $D(k, l) \subset P(k, l)$, which contains the identity, and is stable under:*

- (1) *The horizontal concatenation operation \otimes .*
- (2) *The vertical concatenation \circ , after deleting closed strings in the middle.*
- (3) *The upside-down turning operation $*$ (with reversing of the colors).*

As explained in [22], such categories produce quantum groups. To be more precise, associated to any partition $\pi \in P(k, l)$ is the following linear map:

$$T_\pi(e_{i_1} \otimes \dots \otimes e_{i_k}) = \sum_{j_1 \dots j_l} \delta_\pi \begin{pmatrix} i_1 & \dots & i_k \\ j_1 & \dots & j_l \end{pmatrix} e_{j_1} \otimes \dots \otimes e_{j_l}$$

Here the Kronecker type symbol $\delta_\pi \in \{0, 1\}$ is by definition 1 if all the strings of π join pairs of equal indices, and is 0 otherwise. With this notion in hand, we have:


Definition 2.2. *A compact quantum group $G \subset U_N^+$ is called easy when*

$$\text{Hom}(u^{\otimes k}, u^{\otimes l}) = \text{span} \left(T_\pi \mid \pi \in D(k, l) \right)$$

for any k, l , for a certain category of partitions $D \subset P$.

In other words, the easiness condition states that Tannakian dual of G , also called Schur-Weyl dual, should come in the simplest possible way: from partitions.

In order to discuss some basic examples, consider the categories of pairings, and of noncrossing pairings, $NC_2 \subset P_2$. Consider as well the ‘‘color-matching’’ versions of these categories, $\mathcal{NC}_2 \subset \mathcal{P}_2$, the color-matching condition stating that the various types of strings (upper, lower, through) of our pairings must be colored as follows:



With these notions in hand, we have the following result:

Proposition 2.3. *We have easy quantum groups, as follows,*

$$\begin{array}{ccc}
 U_N & \longrightarrow & U_N^+ \\
 \uparrow & & \uparrow \\
 O_N & \longrightarrow & O_N^+
 \end{array}
 \quad \sim \quad
 \begin{array}{ccc}
 \mathcal{P}_2 & \longleftarrow & \mathcal{NC}_2 \\
 \downarrow & & \downarrow \\
 P_2 & \longleftarrow & NC_2
 \end{array}$$

with the diagram at right describing the corresponding categories of partitions.

Proof. Here the results on the right are the Brauer theorem for O_N , and for U_N , and the results on the left are free versions of Brauer's theorem, discussed in [6]. As a quick, partly heuristic explanation here, all these results follow from Tannakian duality:

(1) U_N^+ is defined via the relations $u^* = u^{-1}$, $u^t = \bar{u}^{-1}$, which tell us that the operators T_π , with $\pi = \begin{array}{c} \circ \\ \bullet \end{array}$ and $\pi = \begin{array}{c} \circ \\ \bullet \end{array}$, must be in the associated Tannakian category C . We therefore obtain $C = \text{span}(T_\pi | \pi \in D)$, with $D = \langle \begin{array}{c} \circ \\ \bullet \end{array}, \begin{array}{c} \circ \\ \bullet \end{array} \rangle = \mathcal{NC}_2$, as claimed.

(2) $O_N^+ \subset U_N^+$ is defined by imposing the relations $u_{ij} = \bar{u}_{ij}$, which tell us that the operators T_π , with $\pi = \begin{array}{c} \circ \\ \bullet \end{array}$ and $\pi = \begin{array}{c} \bullet \\ \circ \end{array}$, must be in the associated Tannakian category C . We therefore obtain $C = \text{span}(T_\pi | \pi \in D)$, with $D = \langle \mathcal{NC}_2, \begin{array}{c} \circ \\ \bullet \end{array}, \begin{array}{c} \bullet \\ \circ \end{array} \rangle = NC_2$, as claimed.

(3) $U_N \subset U_N^+$ is defined via the relations $[u_{ij}, u_{kl}] = 0$ and $[u_{ij}, \bar{u}_{kl}] = 0$, which tell us that the operators T_π , with $\pi = \begin{array}{c} \circ \\ \circ \end{array}$ and $\pi = \begin{array}{c} \bullet \\ \bullet \end{array}$, must be in the associated Tannakian category C . Thus $C = \text{span}(T_\pi | \pi \in D)$, with $D = \langle \mathcal{NC}_2, \begin{array}{c} \circ \\ \circ \end{array}, \begin{array}{c} \bullet \\ \bullet \end{array} \rangle = \mathcal{P}_2$, as claimed.

(4) Finally, in order to deal with O_N , we can use here the formula $O_N = O_N^+ \cap U_N$. At the categorical level, this tells us that the associated Tannakian category is given by $C = \text{span}(T_\pi | \pi \in D)$, with $D = \langle NC_2, \mathcal{P}_2 \rangle = P_2$, as claimed. \square

There are many other examples of easy quantum groups, as for instance the permutation group S_N and its free analogue S_N^+ , the corresponding categories being here the category of all partitions P , and the category of noncrossing partitions $NC \subset P$. See [8].

In the pairing case, however, the main examples remain those in Proposition 2.3. Now observe that the 4 quantum groups there are precisely the unitary groups of the 4 main geometries, from Proposition 1.7. We are therefore led to the following notion:

Definition 2.4. *A noncommutative geometry is called easy when its associated unitary group $O_N \subset G \subset U_N^+$ is easy, coming from a category of pairings $\mathcal{NC}_2 \subset D \subset P_2$.*

Our first task will be that of proving that the 9 geometries from Theorem 1.10 above are all easy. For this purpose, let $P_2^* \subset P_2$ be the category of pairings having the property that when flatenning the pairing, each string has an even number of points between its legs. Let also $\bar{P}_2 \subset P_2$ be the category of pairings having the property that when flatenning the pairing, the number of \circ symbols equals the number of \bullet symbols. Finally, let us define as well categories $\mathcal{P}_2^{**}, \bar{P}_2^*, \bar{NC}_2$ in the obvious way, by taking intersections.

With these notions in hand, we have the following result:

Proposition 2.5. *The basic 9 geometries are easy, with quantum groups as follows,*

$$\begin{array}{ccccc}
 U_N & \longrightarrow & U_N^{**} & \longrightarrow & U_N^+ \\
 \uparrow & & \uparrow & & \uparrow \\
 \mathbb{T}O_N & \longrightarrow & \mathbb{T}O_N^* & \longrightarrow & \mathbb{T}O_N^+ \\
 \uparrow & & \uparrow & & \uparrow \\
 O_N & \longrightarrow & O_N^* & \longrightarrow & O_N^+
 \end{array}
 \quad \sim \quad
 \begin{array}{ccccc}
 \mathcal{P}_2 & \longleftarrow & \mathcal{P}_2^{**} & \longleftarrow & \mathcal{N}C_2 \\
 \downarrow & & \downarrow & & \downarrow \\
 \bar{\mathcal{P}}_2 & \longleftarrow & \bar{\mathcal{P}}_2^* & \longleftarrow & \bar{\mathcal{N}}C_2 \\
 \downarrow & & \downarrow & & \downarrow \\
 P_2 & \longleftarrow & P_2^* & \longleftarrow & NC_2
 \end{array}$$


with the diagram at right describing the corresponding categories of partitions.

Proof. The idea is that of converting the defining relations for the quantum group into statements regarding certain operators of type T_π , and then computing the category of partitions generated by these defining partitions π . More precisely:

(1) First of all, the 4 results at the corners are known from Proposition 2.3. Also, the result for O_N^* is known from [9], and its unitary version, for U_N^{**} , is known from [1]. We are therefore left with proving the 3 results corresponding to the middle rows.

(2) As a first observation here, the result is clear for $\mathbb{T}O_N$, and this, by using an elementary approach. Indeed, if we denote the standard corepresentation by $u = zv$, with $z \in \mathbb{T}$ and with $v = \bar{v}$, then in order to have $Hom(u^{\otimes k}, u^{\otimes l}) \neq \emptyset$, the z variables must cancel, and in the case where they cancel, we obtain the same Hom-space as for O_N . Now since the cancelling property for the z variables corresponds precisely to the fact that k, l must have the same numbers of \circ symbols minus \bullet symbols, the associated Tannakian category must come from the category of pairings $\bar{P}_2 \subset P_2$, as claimed.

(3) In order to deal now with the free version $\mathbb{T}O_N^+$, no such shortcut is available here, and we must use the regular, abstract method. So, observe that the defining relations for this quantum group, namely $ab^* = a^*b$, correspond to the following diagram:



Thus the associated category of partitions is $D = \langle \mathcal{N}C_2, \begin{smallmatrix} \circ & \bullet \\ | & | \\ \bullet & \circ \end{smallmatrix} \rangle = \bar{\mathcal{N}}C_2$, as claimed.

(4) Finally, since we have $\mathbb{T}O_N^* = \mathbb{T}O_N^+ \cap U_N^{**}$, here the associated category of partitions follows to be $D = \langle \bar{\mathcal{N}}C_2, \mathcal{P}_2^{**} \rangle = \bar{P}_2^*$, and this finishes the proof. \square

Now back to the general case, our claim is that, for an easy geometry, there are a few simplifications in the axioms. We first have the following result:

Proposition 2.6. *For a geometry which is easy, coming from a category of pairings $\mathcal{NC}_2 \subset D \subset P_2$, the associated quantum group is given by*

$$C(G) = C(U_N^+) / \left\langle T_\pi \in \text{Hom}(u^{\otimes k}, u^{\otimes l}) \mid \forall k, l, \forall \pi \in D(k, l) \right\rangle$$

and the associated noncommutative torus is $T = \widehat{\Gamma}$, with:

$$\Gamma = F_N / \left\langle g_{i_1} \cdots g_{i_k} = g_{j_1} \cdots g_{j_l} \mid \forall i, j, k, l, \exists \pi \in D(k, l), \delta_\pi \begin{pmatrix} i \\ j \end{pmatrix} \neq 0 \right\rangle$$

Moreover, in both cases, we can just use partitions π which generate the category D .

Proof. The first assertion is well-known, see [8], [18]. If we denote by $g_i = u_{ii}$ the standard coordinates on the associated torus T , then we have, with $g = \text{diag}(g_1, \dots, g_N)$:

$$\begin{aligned} C(T) &= \left[C(U_N^+) / \left\langle T_\pi \in \text{Hom}(u^{\otimes k}, u^{\otimes l}) \mid \forall \pi \in D \right\rangle \right] / \left\langle u_{ij} = 0 \mid \forall i \neq j \right\rangle \\ &= \left[C(U_N^+) / \left\langle u_{ij} = 0 \mid \forall i \neq j \right\rangle \right] / \left\langle T_\pi \in \text{Hom}(u^{\otimes k}, u^{\otimes l}) \mid \forall \pi \in D \right\rangle \\ &= C^*(F_N) / \left\langle T_\pi \in \text{Hom}(g^{\otimes k}, g^{\otimes l}) \mid \forall \pi \in D \right\rangle \end{aligned}$$

The associated discrete group, $\Gamma = \widehat{T}$, is therefore given by:

$$\Gamma = F_N / \left\langle T_\pi \in \text{Hom}(g^{\otimes k}, g^{\otimes l}) \mid \forall \pi \in D \right\rangle$$

Now observe that, with $g = \text{diag}(g_1, \dots, g_N)$, we have:


$$\begin{aligned} T_\pi g^{\otimes k}(e_{i_1} \otimes \cdots \otimes e_{i_k}) &= \sum_{j_1 \dots j_l} \delta_\pi \begin{pmatrix} i_1 & \cdots & i_k \\ j_1 & \cdots & j_l \end{pmatrix} e_{j_1} \otimes \cdots \otimes e_{j_l} \cdot g_{i_1} \cdots g_{i_k} \\ g^{\otimes l} T_\pi(e_{i_1} \otimes \cdots \otimes e_{i_k}) &= \sum_{j_1 \dots j_l} \delta_\pi \begin{pmatrix} i_1 & \cdots & i_k \\ j_1 & \cdots & j_l \end{pmatrix} e_{j_1} \otimes \cdots \otimes e_{j_l} \cdot g_{j_1} \cdots g_{j_l} \end{aligned}$$

We conclude that the relation $T_\pi \in \text{Hom}(g^{\otimes k}, g^{\otimes l})$ reformulates as follows:

$$\delta_\pi \begin{pmatrix} i_1 & \cdots & i_k \\ j_1 & \cdots & j_l \end{pmatrix} \neq 0 \implies g_{i_1} \cdots g_{i_k} = g_{j_1} \cdots g_{j_l}$$

Thus we obtain the formula in the statement. Finally, the last assertion follows from Tannakian duality in the quantum group case, and then in the torus case as well. \square

We conjecture that, in the case of an easy geometry, the category D determines everything, and is determined by everything. Thus, in this case we should have full correspondences, between all the objects involved, with D being the central object:



Observe the similarity with the usual real and complex geometries, which have as well a central object, namely $\mathbb{R}^N, \mathbb{C}^N$. Thus, in a certain abstract sense, for an easy geometry, D is the analogue of the ambient space $\mathbb{R}^N, \mathbb{C}^N$, which cannot be axiomatized.

Here are now a few more abstract notions:

Definition 2.7. *A noncommutative geometry is called:*

- (1) *Amenable, when the discrete quantum group \widehat{G} is amenable.*
- (2) *Weakly amenable, when the discrete group $\Gamma = \widehat{T}$ is amenable.*

Here we use the usual amenability notion for the discrete groups, and for the discrete quantum groups. For the general theory regarding this latter notion, see [19].

We should mention that, for all the known examples of noncommutative geometries in our sense, the coamenability of G is equivalent to the coamenability of T . We conjecture that this should be true in general, but have no idea on how to prove this.

Let us discuss now more in detail the half-classical real geometry, associated to O_N^* . We recall that the projective version of a quantum subgroup $G \subset O_N^+$ is the quotient quantum group $G \rightarrow PG$ having $w_{ij,kl} = u_{ik}u_{jl}$ as fundamental corepresentation. In the classical case, $G \subset O_N$, we recover in this way the usual projective version.

We have the following uniqueness results:

Theorem 2.8. *We have the following results:*

- (1) *O_N^* is the unique easy quantum group $O_N \subset G \subset O_N^+$.*
- (2) *O_N^* is maximal coamenable, in the easy framework.*
- (3) *O_N^* is the biggest quantum group whose projective version PO_N^* is classical.*

Proof. These results are well-known, but we remind here the ideas of the proofs, because these will serve as inspiration for various unitary generalizations, to be done below:

(1) This result is from [9], the idea being that P_2^* is the unique intermediate category of partitions $NC_2 \subset D \subset P_2$. We should mention here that, conjecturally, O_N^* is the unique quantum group $O_N \subset G \subset O_N^+$, even without the easiness assumption. See [5].

(2) This follows from (1), because O_N^* is known to be coamenable, with this latter property coming from instance from an explicit random walk computation.

(3) This is well-known as well. Indeed, the relations $abc = cba$ are equivalent to the relations $abcd = cdab$, as shown by the following two computations:

$$[abc = cba] \implies [abcd = cbad = cdab]$$

$$[abcd = cdab] \implies \left[abc = \sum_d abcdd = \sum_d cdabd = \sum_d cbdda = cba \right]$$


Here we assume that all the variables are standard coordinates, and we have used the quadratic condition relating these coordinates, namely $\sum_d d^2 = 1$. \square

3. COMPLEX GEOMETRIES


In this section and in the next two ones we discuss the construction of the complex half-classical geometry. We will proceed in two steps:

(1) In this section we discuss a first extension of the U_N^{**} geometry, with unitary group denoted U_N^\times . This quantum group is the one constructed in [10], and denoted U_N^* there. The problem, however, is that this geometry is not amenable.


(2) In section 4 we construct and study the “correct” complex analogue of the O_N^* geometry, with unitary group denoted U_N^+ . This quantum group is the one constructed in [4], and denoted $U_{N,\infty}$ there. The enlarged picture will look as follows:



In order to get started, our first task is to look for extensions of the U_N^{**} geometry. This geometry is by definition easy, coming from the following diagrams:



These diagrams stand for the relations $abc = cba$, $abc^* = c^*ba$, and so on, up to $a^*b^*c^* = c^*b^*a^*$. There are some obvious equivalences between these relations, and by erasing the corresponding diagrams, we are led to three diagrams, namely:




In order to extend now the U_N^{**} geometry, the idea would be that of picking one of these diagrams, and using the corresponding relations. But here, we have:

Proposition 3.1. *Consider the following relations, with $a, b, c \in \{x_i\}$:*

- (1) $abc = cba$.
- (2) $ab^*c = cb^*a$
- (3) $abc^* = c^*ba$.

We have then (1) \iff (3) \implies (2).

Proof. The equivalence (1) \iff (3) follows from the following computations:



As for (1 + 3) \implies (2), this is best worked out algebraically, as follows:

$$ab^*c = \sum_d ab^*cdd^* = \sum_d adcb^*d^* = \sum_d cdab^*d^* = \sum_d cb^*add^* = cb^*a$$

Thus we have indeed (1) \iff (3) \implies (2), as claimed. □

Summarizing, we have only one good choice, and are led to the following definition:

Definition 3.2. *We have a sphere, a torus, and a quantum group, as follows:*

- (1) $S_{\mathbb{C},x}^{N-1} \subset S_{\mathbb{C},+}^{N-1}$, obtained via $ab^*c = cb^*a$, with $a, b, c \in \{x_i\}$.
- (2) $T_N^\times \subset \widehat{F}_N$, obtained via $ab^{-1}c = cb^{-1}a$, with $a, b, c \in \{g_i\}$.
- (3) $U_N^\times \subset U_N^+$, obtained via $ab^*c = cb^*a$, with $a, b, c \in \{u_{ij}\}$.

Observe that U_N^\times is indeed a quantum group, containing U_N^{**} . We will prove later on that we have indeed a noncommutative geometry, in the sense of Definition 1.5.

Let us clarify now a few more algebraic issues. The most elegant approach to the U_N^\times geometry is in fact via projective space theory, using the following notions:

Definition 3.3. *Given a subspace $X \subset S_{\mathbb{C},+}^{N-1}$, we define quotients as follows,*

- (1) *Left projective version: $X \rightarrow PX$, with coordinates $p_{ij} = x_i x_j^*$,*
- (2) *Right projective version: $X \rightarrow P'X$, with coordinates $q_{ij} = x_j^* x_i$,*
- (3) *Full projective version: $X \rightarrow \mathcal{P}X$, with coordinates p_{ij}, q_{ij} ,*

and we say that X is left/right/full half-classical when these spaces are classical.

Observe that in the classical case, $X \subset S_{\mathbb{C}}^{N-1}$, the three projective versions coincide, and equal the usual projective version, obtained by dividing under the action of \mathbb{T} .

In the real case, $X \subset S_{\mathbb{R},+}^{N-1}$, the three projective versions coincide as well, and X is left or right half-classical when $X \subset S_{\mathbb{R},*}^{N-1}$. This follows indeed from Theorem 2.9.

In relation now with $S_{\mathbb{C},\times}^{N-1}, S_{\mathbb{C},**}^{N-1}$, we can use the following simple fact, from [4]:

Proposition 3.4. *Let $X \subset S_{\mathbb{C},+}^{N-1}$, with coordinates x_1, \dots, x_N .*

- (1) *$X \subset S_{\mathbb{C},\times}^{N-1}$ precisely when $\{x_i x_j^*\}$ commute, and $\{x_i^* x_j\}$ commute as well.*
- (2) *$X \subset S_{\mathbb{C},**}^{N-1}$ precisely when the variables $\{x_i x_j, x_i x_j^*, x_i^* x_j, x_i^* x_j^*\}$ all commute.*

Proof. Regarding the first assertion, the implication “ \implies ” follows from:

$$ab^*cd^* = cb^*ad^* = cd^*ab^* \quad , \quad a^*bc^*d = c^*ba^*d = c^*da^*b$$

As for the implication “ \impliedby ”, this is obtained as follows, by using the commutation assumptions in the statement, and by summing over $e = x_i$:

$$ae^*eb^*c = ab^*ce^*e = ce^*ab^*e = cb^*ee^*a \implies ab^*c = cb^*a$$

The proof of the second assertion is similar, because we can remove all the $*$ signs, except for those concerning e^* , and use the above computations with $a, b, c, d \in \{x_i, x_i^*\}$. \square

With the above result in hand, we can now formulate:

Proposition 3.5. *We have the following results:*

- (1) *$S_{\mathbb{C},\times}^{N-1}$ is left and right half-classical, and is maximal with this property.*
- (2) *$S_{\mathbb{C},**}^{N-1}$ is fully half-classical.*
- (3) *$S_{\mathbb{R},*}^{N-1}$ is fully half-classical, and is maximal inside $S_{\mathbb{R},+}^{N-1}$ with this property.*

Proof. All these assertions follow indeed from Proposition 3.4. We will see later on, in connection with (2), that $S_{\mathbb{C},**}^{N-1}$ is far from being maximal, with the property there. \square

We still have an issue to be clarified, namely that of proving that, in the diagram drawn in the beginning of this section, U_N^\times sits indeed above $\mathbb{T}O_N^*$. But this comes from:

Proposition 3.6. *We have $\mathbb{T}O_N^+ \cap U_N^\times = \mathbb{T}O_N^*$, as quantum subgroups of U_N^+ .*

Proof. According to the definition of $\mathbb{T}O_N^*$, from section 1, this quantum group appears as $\mathbb{T}O_N^* = \mathbb{T}O_N^+ \cap U_N^{**}$. Thus, we must prove that we have $\mathbb{T}O_N^+ \cap U_N^\times \subset U_N^{**}$.

In terms of defining relations, we must prove that, from $ab^* = a^*b$ and $ab^*c = cb^*a$ for any $a, b, c \in \{u_{ij}\}$, we can deduce that we have $abc = cba$, for any $a, b, c \in \{u_{ij}, u_{ij}^*\}$.

But this is clear, because by using $ab^* = a^*b$, we can first obtain $a^*bc = cba^*$, and then, by using Proposition 3.1, we can obtain from this the other relations as well. \square

We recall that the free complexification of a compact quantum group G , with standard coordinates denoted v_{ij} , is the compact quantum group \tilde{G} corresponding to the subalgebra $C(\tilde{G}) \subset C(\mathbb{T}) * C(G)$ generated by the variables $u_{ij} = zv_{ij}$, where z is the standard generator of $C(\mathbb{T})$. Observe that \tilde{G} is indeed a quantum group, because it appears as a subgroup of $\mathbb{T} * G$, the quantum group associated to $C(\mathbb{T}) * C(G)$. See [20].

Following [2], we have the following result:

Proposition 3.7. *The quantum groups $O_N^*, U_N, U_N^{**}, U_N^\times$ have the following properties:*

- (1) *They have the same left projective version, equal to PU_N .*
- (2) *They have the same free complexification, equal to U_N^\times .*

Proof. With terminology and notations from [2], the idea is as follows:

(1) Here $PO_N^* = PU_N$ is known from Proposition 2.8, and it is clear as well that we have $PU_N \subset PU_N^{**} \subset PU_N^\times$. Now by using Proposition 3.4 (1), we conclude that PU_N^\times is classical, and so we must have $PU_N^\times \subset (PU_N^+)_{class}$. But this latter space $(PU_N^+)_{class}$ is known to be equal to PU_N , and this finishes the proof. See [2], [10].

(2) If we denote by v_{ij} the standard coordinates on $G = O_N^*, U_N, U_N^{**}, U_N^\times$, and by z the generator of a copy of $C(\mathbb{T})$, free from $C(G)$, then with $a, b, c \in \{v_{ij}\}$ we have:

$$(za)(zb)^*(zc) = zab^*c = zcb^*a = (zc)(zb)^*(za)$$

Thus we have $\tilde{G} \subset U_N^\times$. Conversely now, it follows from the general theory of the free complexifications of easy quantum groups [20] that both $K = \tilde{G}, U_N^\times$ should appear as free complexifications of certain intermediate easy quantum groups $O_N \subset H \subset O_N^+$. On the other hand, since we have $PH = P\tilde{H} = PK = PU_N$, the only choice here is $H = O_N^*$. Thus we have $\tilde{G} = U_N^\times = \tilde{O}_N^*$, and this finishes the proof. See [2]. \square

Let us prove now the quantum isometry group result. This is known since [2], but we have now a much simpler proof for this fact. We use the following notion:

Definition 3.8. *Given a closed subgroup $G \subset U_N^+$, and a closed subset $X \subset S_{\mathbb{C},+}^{N-1}$, we say that G acts projectively on X , and write $PG \curvearrowright PX$, when the formula*

$$\Phi(x_i x_j^*) = \sum_{ab} u_{ia} u_{jb}^* \otimes x_a x_b^*$$

defines a morphism of C^ -algebras $\Phi : C(PX) \rightarrow C(PG) \otimes C(PX)$.*

Observe the similarity with Definition 1.3 above, dealing with the affine case. As in the affine case, such a morphism is automatically coassociative and counital. Observe also that any affine action $G \curvearrowright X$ produces a projective action $PG \curvearrowright PX$.

We can now formulate our quantum isometry group results, as follows:

Proposition 3.9. *We have the following results:*

- (1) $PG \curvearrowright P_{\mathbb{C}}^{N-1}$ implies $G \subset U_N^{\times}$.
- (2) $G \curvearrowright S_{\mathbb{C}, \times}^{N-1}$ implies $G \subset U_N^{\times}$.

Proof. Since $G \curvearrowright S_{\mathbb{C}, \times}^{N-1}$ implies that we have $PG \curvearrowright PS_{\mathbb{C}, \times}^{N-1} = P_{\mathbb{C}}^{N-1}$, we just have to prove the first assertion. For this purpose, we use an old method from [11], as in [3].

Consider indeed a coaction map, written $\Phi(p_{ij}) = \sum_{ab} u_{ia} u_{jb}^* \otimes p_{ab}$, with $p_{ab} = z_a \bar{z}_b$. The idea will be that of using the formula $p_{ab} p_{cd} = p_{ad} p_{cb}$. We have:

$$\begin{aligned} \Phi(p_{ij} p_{kl}) &= \sum_{abcd} u_{ia} u_{jb}^* u_{kc} u_{ld}^* \otimes p_{ab} p_{cd} \\ \Phi(p_{il} p_{kj}) &= \sum_{abcd} u_{ia} u_{ld}^* u_{kc} u_{jb}^* \otimes p_{ad} p_{cb} \end{aligned}$$

The left terms being equal, and the last terms on the right being equal too, we deduce that, with $[a, b, c] = abc - cba$, we must have the following equality:

$$\sum_{abcd} u_{ia} [u_{jb}^*, u_{kc}, u_{ld}^*] \otimes p_{ab} p_{cd} = 0$$

Now since the quantities $p_{ab} p_{cd} = z_a \bar{z}_b z_c \bar{z}_d$ on the right depend only on the numbers $|\{a, c\}|, |\{b, d\}| \in \{1, 2\}$, and this dependence produces the only possible linear relations between the variables $p_{ab} p_{cd}$, we are led to $2 \times 2 = 4$ equations, as follows:

- (1) $u_{ia} [u_{jb}^*, u_{ka}, u_{lb}^*] = 0, \forall a, b$.
- (2) $u_{ia} [u_{jb}^*, u_{ka}, u_{ld}^*] + u_{ia} [u_{jd}^*, u_{ka}, u_{lb}^*] = 0, \forall a, \forall b \neq d$.
- (3) $u_{ia} [u_{jb}^*, u_{kc}, u_{lb}^*] + u_{ic} [u_{jb}^*, u_{ka}, u_{lb}^*] = 0, \forall a \neq c, \forall b$.
- (4) $u_{ia} ([u_{jb}^*, u_{kc}, u_{ld}^*] + [u_{jd}^*, u_{kc}, u_{lb}^*]) + u_{ic} ([u_{jb}^*, u_{ka}, u_{ld}^*] + [u_{jd}^*, u_{ka}, u_{lb}^*]) = 0, \forall a \neq c, \forall b \neq d$.

From (1,2) we conclude that (2) holds with no restriction on the indices. By multiplying now this formula to the left by u_{ia}^* , and then summing over i , we obtain:

$$[u_{jb}^*, u_{ka}, u_{ld}^*] + [u_{jd}^*, u_{ka}, u_{lb}^*] = 0$$

By applying now the antipode, then the involution, and finally by suitably relabelling all the indices, we successively obtain from this formula:

$$\begin{aligned} & [u_{dl}, u_{ak}^*, u_{bj}] + [u_{bl}, u_{ak}^*, u_{dj}] = 0 \\ \implies & [u_{dl}^*, u_{ak}, u_{bj}^*] + [u_{bl}^*, u_{ak}, u_{dj}^*] = 0 \\ \implies & [u_{ld}^*, u_{ka}, u_{jb}^*] + [u_{jd}^*, u_{ka}, u_{lb}^*] = 0 \end{aligned}$$

Now by comparing with the original relation, we conclude that we have:

$$[u_{jb}^*, u_{ka}^*, u_{ld}^*] = [u_{jd}^*, u_{ka}^*, u_{lb}^*] = 0$$

Thus we have reached to the formulae defining U_N^\times , and we are done. \square

We have now all the needed ingredients for proving:

Theorem 3.10. *The sphere $S_{\mathbb{R}, \times}^{N-1}$, the torus T_N^\times and the quantum group U_N^\times form a noncommutative geometry. This geometry is easy, and not amenable.*

Proof. The verification of all the axioms is standard, with the only non-trivial fact, namely the universality of the action $U_N^\times \curvearrowright S_{\mathbb{C}, \times}^{N-1}$, coming from Proposition 3.9 above.

The non-coamenability of U_N^\times can be checked by using a free complexification trick. We know that the discrete group $\Gamma_N^\times = \widehat{T_N^\times}$ is given by:

$$\Gamma_N^\times = \left\langle g_1, \dots, g_N \mid ab^{-1}c = cb^{-1}a, \forall a, b, c \in \{g_i\} \right\rangle$$

Now observe that, if we denote by h_1, \dots, h_N the standard generators of \mathbb{Z}^N , and by z the generator of a copy of \mathbb{Z} , which is free from \mathbb{Z}^N , then with $a, b, c \in \{h_i\}$ we have:

$$(za)(zb)^{-1}(zc) = zab^{-1}c = zcb^{-1}a = (zc)(zb)^{-1}(za)$$

We therefore have a group morphism $\Gamma_N^\times \rightarrow \mathbb{Z} * \mathbb{Z}^N$, given by $g_i \rightarrow zh_i$. Now observe that the image of this morphism contains the following two elements:

$$\begin{aligned} (zh_1)^{-1}(zh_2) &= h_1^{-1}h_2 \\ (zh_1)(zh_2)^{-1} &= zh_1h_2^{-1}z^* \end{aligned}$$

These elements being free, we obtain a copy of F_2 inside the image. Thus the image, and then Γ_N^\times , and then $\widehat{U_N^\times}$ itself, follow to be non-amenable as well. \square

The U_N^\times geometry can be further investigated by using various algebraic tricks, and notably the free complexification formula $U_N^\times = \widehat{U_N}$. We refer here to [2].

4. HALF-CLASSICAL GEOMETRY

In this section we introduce and study the U_N^* geometry, which is the “correct” complex half-classical one, in the sense that it is the biggest half-classical geometry.

In view of Definition 3.3 above, we can indeed formulate:

Definition 4.1. *We have a sphere, a torus, and a quantum group, as follows:*


- (1) $S_{\mathbb{C}, * }^{N-1} \subset S_{\mathbb{C}, + }^{N-1}$, obtained via “ ab^*, a^*b all commute”, with $a, b, c \in \{x_i\}$.
- (2) $T_N^* \subset \widehat{F_N}$, obtained via “ $ab^{-1}, a^{-1}b$ all commute”, with $a, b, c \in \{g_i\}$.
- (3) $U_N^* \subset U_N^+$, obtained via “ ab^*, a^*b all commute”, with $a, b, c \in \{u_{ij}\}$.

In other words, these are the biggest half-classical sphere, torus, and quantum group.

As a first remark, the real version of $S_{\mathbb{C},*}^{N-1}$, obtained by imposing the conditions $x_i = x_i^*$ to the standard coordinates, is the half-classical real sphere $S_{\mathbb{R},*}^{N-1}$. Observe also that we have inclusions as follows, coming from the various results in Proposition 3.4:

$$S_{\mathbb{C},**}^{N-1} \subset S_{\mathbb{C},*}^{N-1} \subset S_{\mathbb{C},\times}^{N-1}$$

Similar inclusions are valid for the tori, and for the quantum groups. Finally, observe that U_N^* is easy, coming from the following two diagrams:



In order to better understand the construction $X \rightarrow \mathcal{P}X$, and the definition of $S_{\mathbb{C},*}^{N-1}$ itself, let us perform now some explicit computations. We have here:

Proposition 4.2. *The projective versions of $S_{\mathbb{C},**}^{N-1} \subset S_{\mathbb{C},*}^{N-1} \subset S_{\mathbb{C},\times}^{N-1}$ are given by*

$$\begin{array}{ccccc} S_{\mathbb{C},**}^{N-1} & \longrightarrow & S_{\mathbb{C},*}^{N-1} & \longrightarrow & S_{\mathbb{C},\times}^{N-1} \\ \downarrow & & \downarrow & & \downarrow \\ P_{\mathbb{C}}^{N-1} \times P_{\mathbb{C}}^{N-1} & \longrightarrow & P_{\mathbb{C}}^{N-1} \times P_{\mathbb{C}}^{N-1} & \longrightarrow & P_{\mathbb{C}}^{N-1} \circ P_{\mathbb{C}}^{N-1} \end{array}$$

where the product on the bottom right is constructed by using a spinning free unitary.

Proof. We use the following presentation result, which comes from the Gelfand theorem, and from the fact that $P_{\mathbb{C}}^{N-1}$ is the space of rank 1 projections in $M_N(\mathbb{C})$:

$$C(P_{\mathbb{C}}^{N-1}) = C_{comm}^* \left\{ (p_{ij})_{i,j=1,\dots,N} \mid p = p^2 = p^*, \text{Tr}(p) = 1 \right\}$$

Let us first discuss the computation of the spaces $\mathcal{P}S_{\mathbb{C},**}^{N-1} \subset \mathcal{P}S_{\mathbb{C},*}^{N-1}$. We know that these spaces are both classical. We also know that the left and right components, in the sense of Definition 3.3 above, of these spaces are all equal to $P_{\mathbb{C}}^{N-1}$, for instance because their standard generators satisfy the above defining relations for $C(P_{\mathbb{C}}^{N-1})$.

In order to finish, it remains to prove that the subspaces $PS_{\mathbb{C},**}^{N-1}, P'S_{\mathbb{C},**}^{N-1} \subset \mathcal{P}S_{\mathbb{C},**}^{N-1}$, which are both isomorphic to $P_{\mathbb{C}}^{N-1}$, are in generic position. For this purpose, we can use a suitable matrix model, coming from [4]. Let indeed u_i, v_i be the standard coordinates of two independent copies of $S_{\mathbb{C}}^{N-1}$, and consider the following matrices:

$$X_i = \begin{pmatrix} 0 & u_i \\ v_i & 0 \end{pmatrix}, \quad X_i^* = \begin{pmatrix} 0 & \bar{v}_i \\ \bar{u}_i & 0 \end{pmatrix}$$

We have then the following formula, coming from $\sum_i |u_i|^2 = \sum_i |v_i|^2 = 1$:

$$\sum_i X_i X_i^* = \sum_i X_i^* X_i = 1$$

In addition, the relations $abc = cba$ hold as well, for any $a, b, c \in \{X_i, X_i^*\}$. Thus we have a matrix model, as follows:

$$C(S_{\mathbb{C},**}^{N-1}) \rightarrow M_2(C(S_{\mathbb{C}}^{N-1} \times S_{\mathbb{C}}^{N-1})) \quad , \quad x_i \rightarrow X_i$$

The point now is that, in this model, we have the following formulae:

$$X_i X_j^* = \begin{pmatrix} u_i \bar{u}_j & 0 \\ 0 & v_i \bar{v}_j \end{pmatrix} \quad , \quad X_j^* X_i = \begin{pmatrix} v_i \bar{v}_j & 0 \\ 0 & u_i \bar{u}_j \end{pmatrix}$$

Now since these matrices are conjugated by an order 2 automorphism, the algebra that they generate is isomorphic to $C(P_{\mathbb{C}}^{N-1} \times P_{\mathbb{C}}^{N-1})$, and this finishes the proof.

Finally, regarding the computation for $S_{\mathbb{C},\times}^{N-1}$, let us denote by $p_{ij} = z_i \bar{z}_j$ the standard coordinates on $P_{\mathbb{C}}^{N-1} = PS_{\mathbb{C}}^{N-1}$. In the usual free complexification model for $S_{\mathbb{C},\times}^{N-1}$, namely $\tilde{S}_{\mathbb{C}}^{N-1}$, we have then $x_i x_j^* = p_{ij}$, $x_j^* x_i = z p_{ij} z^*$, and this gives the result. \square

In the order to verify now the axioms, we use a modification of the proof for U_N^{\times} , from section 3 above. First, we have the following version of Definition 3.8 above:

Definition 4.3. *Given a closed subgroup $G \subset U_N^+$, and a closed subset $X \subset S_{\mathbb{C},+}^{N-1}$, we say that G acts fully projectively on X , and write $\mathcal{P}G \curvearrowright \mathcal{P}X$, when the formulae*

$$\Phi(x_i x_j^*) = \sum_{ab} u_{ia} u_{jb}^* \otimes x_a x_b^*$$

$$\Phi(x_i^* x_j) = \sum_{ab} u_{ia}^* u_{jb} \otimes x_a^* x_b$$

define a morphism of C^* -algebras $\Phi : C(\mathcal{P}X) \rightarrow C(\mathcal{P}G) \otimes C(\mathcal{P}X)$.

As in the affine case, such a morphism is automatically coassociative and counital. Observe also that any affine action $G \curvearrowright X$ produces a projective action $\mathcal{P}G \curvearrowright \mathcal{P}X$.

We can now formulate our quantum isometry group results, as follows:

Proposition 4.4. *We have the following results:*

- (1) *We have an affine action $U_N^* \curvearrowright S_{\mathbb{C},*}^{N-1}$.*
- (2) *$\mathcal{P}G \curvearrowright P_{\mathbb{C}}^{N-1} \times P_{\mathbb{C}}^{N-1}$ implies $G \subset U_N^*$.*
- (3) *$G \curvearrowright S_{\mathbb{C},*}^{N-1}$ implies $G \subset U_N^*$.*
- (4) *U_N^* is the quantum isometry group of $S_{\mathbb{C},*}^{N-1}$.*

Proof. Our first claim is that it is enough to prove (2). Indeed, in order to prove (1), observe that with $X_i = \sum_a u_{ia} \otimes x_a$, we have the following formulae:


$$X_i X_j^* = \sum_{ab} u_{ia} u_{jb}^* \otimes x_a x_b^* \quad , \quad X_j^* X_i = \sum_{ab} u_{jb}^* u_{ia} \otimes x_b^* x_a$$

Now since the various variables on the right pairwise commute, the variables on the left commute as well, and so we can define the action map, by $x_i \rightarrow X_i$.

The other remark is that since $G \curvearrowright S_{\mathbb{C},*}^{N-1}$ implies $\mathcal{P}G \curvearrowright PS_{\mathbb{C},\times}^{N-1} = P_{\mathbb{C}}^{N-1} \times P_{\mathbb{C}}^{N-1}$, we have (2) \implies (3). Finally, the implication (1 + 3) \implies (4) is trivial.

In order to prove now (2), observe that $\mathcal{P}G \curvearrowright P_{\mathbb{C}}^{N-1} \times P_{\mathbb{C}}^{N-1}$ implies $PG \curvearrowright P_{\mathbb{C}}^{N-1}$. Thus, we can use Proposition 3.9 (1), and we obtain $G \subset U_N^\times$.

We are therefore left with proving that $\mathcal{P}G \curvearrowright P_{\mathbb{C}}^{N-1} \times P_{\mathbb{C}}^{N-1}$ implies that the following diagram belongs to the Tannakian category of G :



For this purpose, consider indeed a coaction map, written as in Definition 4.3. By multiplying the two relations there, we obtain:

$$\begin{aligned} \Phi(x_i x_j^* x_k^* x_l) &= \sum_{abcd} u_{ia} u_{jb}^* u_{kc}^* u_{ld} \otimes x_a x_b^* x_c^* x_d \\ \Phi(x_k^* x_l x_i x_j^*) &= \sum_{abcd} u_{kc}^* u_{ld} u_{ia} u_{jb}^* \otimes x_c^* x_d x_a x_b^* \end{aligned}$$

Assuming now that x_1, \dots, x_N are the standard coordinates on $S_{\mathbb{C},*}^{N-1}$, the products of x variables at left are equal, and so are the products at right. Thus, we have:

$$\sum_{abcd} [u_{ia} u_{jb}^*, u_{kc}^* u_{ld}] \otimes x_a x_b^* x_c^* x_d = 0$$

Now recall that, in view of Proposition 4.2 above, we can write $x_a x_b^* x_c^* x_d = p_{ab} \otimes q_{dc}$, where p_{ab}, q_{cd} are the standard coordinates on $P_{\mathbb{C}}^{N-1}$. Thus, our formula becomes:

$$\sum_{abcd} [u_{ia} u_{jb}^*, u_{kc}^* u_{ld}] \otimes p_{ab} \otimes q_{dc} = 0$$

Now since the variables on the right are linearly independent, we obtain that all the commutators vanish, and this finishes the proof. \square

We can now formulate our main result, as follows:

Theorem 4.5. *We have a noncommutative geometry, with sphere $S_{\mathbb{C},*}^{N-1}$, torus T_N^* and quantum group U_N^* . This is the biggest geometry having a classical projective version.*

Proof. The verification of all the axioms is standard, with the only non-trivial fact, namely the universality of the action $U_N^* \curvearrowright S_{\mathbb{C},*}^{N-1}$, coming from Proposition 4.4 above. \square

In view of the above result, the U_N^* geometry seems to be the “correct” complex half-classical geometry. This geometry waits of course to be developed, notably with analogues of [12], [13]. Finally, we conjecture that this geometry is amenable, and is moreover maximal with this amenability property, at least in the easy framework.

REFERENCES

- [1] T. Banica, Liberations and twists of real and complex spheres, *J. Geom. Phys.* **96** (2015), 1–25.
- [2] T. Banica, Half-liberated manifolds, and their quantum isometries, *Glasg. Math. J.*, to appear.
- [3] T. Banica, Quantum isometries, noncommutative spheres, and related integrals, *Banach Center Publ.*, to appear.
- [4] T. Banica and J. Bichon, Matrix models for noncommutative algebraic manifolds, *J. Lond. Math. Soc.*, to appear.
- [5] T. Banica, J. Bichon, B. Collins and S. Curran, A maximality result for orthogonal quantum groups, *Comm. Algebra* **41** (2013), 656–665.
- [6] T. Banica and B. Collins, Integration over compact quantum groups, *Publ. Res. Inst. Math. Sci.* **43** (2007), 277–302.
- [7] T. Banica and D. Goswami, Quantum isometries and noncommutative spheres, *Comm. Math. Phys.* **298** (2010), 343–356.
- [8] T. Banica and R. Speicher, Liberation of orthogonal Lie groups, *Adv. Math.* **222** (2009), 1461–1501.
- [9] T. Banica and R. Vergnioux, Invariants of the half-liberated orthogonal group, *Ann. Inst. Fourier* **60** (2010), 2137–2164.
- [10] J. Bhowmick, F. D’Andrea, B. Das and L. Dabrowski, Quantum gauge symmetries in noncommutative geometry, *J. Noncommut. Geom.* **8** (2014), 433–471.
- [11] J. Bhowmick and D. Goswami, Quantum isometry groups: examples and computations, *Comm. Math. Phys.* **285** (2009), 421–444.
- [12] J. Bichon, Half-liberated real spheres and their subspaces, *Colloq. Math.* **144** (2016), 273–287.
- [13] J. Bichon and M. Dubois-Violette, Half-commutative orthogonal Hopf algebras, *Pacific J. Math.* **263** (2013), 13–28.
- [14] R. Brauer, On algebras which are connected with the semisimple continuous groups, *Ann. of Math.* **38** (1937), 857–872.
- [15] A. Freslon, On the partition approach to Schur-Weyl duality and free quantum groups, preprint 2014.
- [16] D. Goswami, Quantum group of isometries in classical and noncommutative geometry, *Comm. Math. Phys.* **285** (2009), 141–160.
- [17] D. Goswami, Existence and examples of quantum isometry groups for a class of compact metric spaces, *Adv. Math.* **280** (2015), 340–359.
- [18] S. Malacarne, Woronowicz’s Tannaka-Krein duality and free orthogonal quantum groups, preprint 2016.
- [19] S. Neshveyev and L. Tuset, Compact quantum groups and their representation categories, SMF (2013).
- [20] S. Raum, Isomorphisms and fusion rules of orthogonal free quantum groups and their complexifications, *Proc. Amer. Math. Soc.* **140** (2012), 3207–3218.

- [21] S. Raum and M. Weber, The full classification of orthogonal easy quantum groups, *Comm. Math. Phys.* **341** (2016), 751–779.
- [22] P. Tarrago and M. Weber, Unitary easy quantum groups: the free case and the group case, preprint 2015.
- [23] S. Wang, Free products of compact quantum groups, *Comm. Math. Phys.* **167** (1995), 671–692.
- [24] S. Wang, Quantum symmetry groups of finite spaces, *Comm. Math. Phys.* **195** (1998), 195–211.
- [25] S.L. Woronowicz, Compact matrix pseudogroups, *Comm. Math. Phys.* **111** (1987), 613–665.
- [26] S.L. Woronowicz, Tannaka-Krein duality for compact matrix pseudogroups. Twisted $SU(N)$ groups, *Invent. Math.* **93** (1988), 35–76.

T.B.: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CERGY-PONTOISE, F-95000 CERGY-PONTOISE, FRANCE. teodor.banica@u-cergy.fr

J.B.: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CLERMONT AUVERGNE, F-63178 AUBIERE, FRANCE. julien.bichon@uca.fr