

## Abstracts

### Adelic quadratic spaces

ÉRIC GAUDRON

(joint work with Gaël Rémond)

We examine the links between linear and quadratic equations through the search of algebraic solutions of small heights.

**0.1.** The starting point is a theorem by Cassels (1955, [1]) and Davenport (1957, [3]) which asserts that if  $q: \mathbb{Q}^n \rightarrow \mathbb{Q}$  is a non-zero isotropic quadratic form with integral coefficients  $(a_{i,j})_{i,j}$  then there exists a vector  $x = (x_1, \dots, x_n) \in \mathbb{Z}^n \setminus \{0\}$  such that  $q(x) = 0$  and

$$\sum_{i=1}^n x_i^2 \leq \left( 2\gamma_{n-1}^2 \sum_{i,j} a_{i,j}^2 \right)^{(n-1)/2}$$

( $\gamma_{n-1}$  is the Hermite constant). Our aim is to give a generalization of this statement in the context of rigid adelic spaces (introduced in the preceding talk by Gaël Rémond).

Let  $K$  be an algebraic extension of  $\mathbb{Q}$  and  $n$  be a positive integer. We denote by  $c_K(n)$  the supremum over all rigid adelic spaces  $E$  over  $K$  of the real numbers

$$\inf \{ H_E(x)^n H(E)^{-1}; x \in E \setminus \{0\} \}$$

( $H_E(x)$  and  $H(E)$  are the heights of  $x$  and  $E$  with respect to the metrics on  $E$ ). According to [6], the field  $K$  is called a *Siegel field* if  $c_K(n) < +\infty$  for all  $n \geq 1$ . We have  $c_{\mathbb{Q}}(n) = \gamma_n^{n/2}$ ,

$$c_K(n) \leq \left( n |\Delta_{K/\mathbb{Q}}|^{1/[K:\mathbb{Q}]} \right)^{n/2}$$

if  $K$  is a number field of absolute discriminant  $\Delta_{K/\mathbb{Q}}$  and

$$c_{\overline{\mathbb{Q}}}(n) = \exp \left\{ \frac{n}{2} \left( \frac{1}{2} + \dots + \frac{1}{n} \right) \right\}$$

(see [6]).

An *adelic quadratic space*  $(E, q)$  over  $K$  is a rigid adelic space  $E/K$  endowed with a quadratic form  $q: E \rightarrow K$ . In this framework, several problems can be raised (here, small = of small height):

- 1) Existence of a small isotropic vector,
- 2) Existence of a small maximal totally isotropic subspace,
- 3) Existence of a basis of  $E$  composed of small isotropic vectors.

There exist between 25 and 30 articles in the literature dealing with these questions (essentially when  $K$  a number field or  $\overline{\mathbb{Q}}$ ). A common divisor to these works is the notion of Siegel's lemma. We shall provide solutions to these three problems, which are optimal with respect to the height of  $q$ .

**0.2.** The following statement gives an answer to the problem 2.

**Theorem 1.** *Assume  $q$  is isotropic. Then, for all  $\varepsilon > 0$ , there exists a maximal totally isotropic subspace  $F$  of  $E$  of dimension  $d \geq 1$  and height*

$$H(F) \leq (1 + \varepsilon)c_K(n - d) (2H(q))^{(n-d)/2} H(E).$$

Here  $H(q)$  is the height of  $q$  built from local operators norms (see [7]). For instance, in the context of Cassels and Davenport Theorem, one can prove that  $H(q) \leq (\sum_{i,j} a_{i,j}^2)^{1/2}$ . Theorem 1 generalizes and improves theorems by Schlickewei (1985,  $K = \mathbb{Q}$ , [9]), Vaaler (1987,  $K$  number field, [10]) and Fukshansky (2008,  $K = \overline{\mathbb{Q}}$ , [5]). Using a Siegel's lemma in such a subspace  $F$ , we obtain an answer to Problem 1:

**Quadratic Siegel's lemma.** *If  $q$  is isotropic then, for all  $\varepsilon > 0$ , there exists  $x \in E \setminus \{0\}$  such that  $q(x) = 0$  and*

$$H_E(x) \leq (1 + \varepsilon) \left( c_K(n) (2H(q))^{(n-d)/2} H(E) \right)^{1/d}.$$

The proof of Theorem 1 follows from an estimate of the height of a suitable  $q$ -orthogonal symmetric of an almost minimal height subspace  $F$  (chosen among maximal totally isotropic subspaces of  $E$ ) and from a Siegel's lemma used with the quotient  $E/F$ . To be interesting, Theorem 1 must be applied in a Siegel field ( $c_K(n - d) < \infty$ ). But the converse is true: it can be also proved that to be a Siegel field is a necessary condition when a quadratic Siegel's lemma exists (take  $q(x) = \ell(x)^2$  with  $\ell: E \rightarrow K$  a linear form and use [6, § 4.8]).

**0.3.** Now, let us tackle the problem of a small isotropic basis of an adelic quadratic space  $(E, q)$  over a Siegel field  $K$ . Assume that there exists a nondegenerate isotropic vector in  $E$ . It is well known then that there exists a basis  $(e_1, \dots, e_n)$  of  $E$  such that  $q(e_i) = 0$  for all  $1 \leq i \leq n$ . Our goal is to have also the heights of  $e_i$ 's *small*. An obvious approach rests on an induction process, choosing  $e_i \in E \setminus K.e_1 \oplus \dots \oplus K.e_{i-1}$  with small height and  $q(e_i) = 0$ . That leads us to the following variant of the quadratic Siegel's lemma:

- 1a) Let  $I$  be an ideal of the ring of polynomials of  $E$  and denote by  $Z(I)$  the set of zeros  $\{x \in E; \forall P \in I, P(x) = 0\}$ . How to bound

$$\inf \{H_E(x); q(x) = 0 \text{ and } x \notin Z(I)\} ?$$

(Quadratic Siegel's lemma avoiding an algebraic set.)

To simplify, we state our result only for the standard adelic space  $E = K^n$ .

**Theorem 2.** *Let  $q: K^n \rightarrow K$  be a quadratic form and let  $I$  be an ideal of  $K[X_1, \dots, X_n]$  generated by polynomials of (total) degree  $\leq M$ . Assume (i)  $q \neq 0$  and (ii)  $\exists x \notin Z(I); q(x) = 0$ . Then there exists a constant  $c(n, K) \geq 1$ , which depends only on  $n$  and  $K$ , such that the vector  $x$  in condition (ii) can also be chosen with height*

$$H_{K^n}(x) \leq c(n, K) M^3 H(q)^{(n-d+1)/2}$$

where  $d$  is the dimension of maximal totally isotropic subspaces of  $(K^n, q)$ .

The constant  $c(n, K)$  can be made fully explicit (see [7, § 7]). This statement generalizes and improves previous results by Masser (1998,  $K = \mathbb{Q}$ ,  $Z(I)$  hyperplane, [8]), Fukshansky (2004,  $K$  number field,  $Z(I)$  union of hyperplanes, [4]) and Chan, Fukshansky & Henshaw (2014, [2]). Moreover the exponent  $(n - d + 1)/2$  of  $H(q)$  is best possible: take  $E = \mathbb{Q}^n$ ,  $a, d \geq 1$  integers,  $Z(I) = \{x_d = 0\}$  and

$$q(x) = 2x_{d+1}x_d - a^2x_d^2 - (x_{d+2} - ax_{d+1})^2 - \cdots - (x_n - ax_{n-1})^2.$$

We have  $H(q) = O_{a \rightarrow +\infty}(a^2)$  and if  $x$  is isotropic then  $|x_n| \geq a^{n-d+1}|x_d|/4$ . The proof of Theorem 2 relies on an avoiding Siegel's lemma and a geometric lemma. From Theorem 2 can easily be deduced a small-height isotropic basis of  $E$ .

Complete proofs and further results are given in [7].

#### REFERENCES

- [1] J. Cassels, *Bounds for the least solutions of homogeneous quadratic equations*, Proc. Cambridge Philos. Soc. **51** (1955), 262–264.
- [2] W.K. Chan, L. Fukshansky and G. Henshaw, *Small zeros of quadratic forms outside a union of varieties*, Trans. Amer. Math. Soc. **366** (2014), 5587–5612.
- [3] H. Davenport, *Note on a theorem of Cassels*, Proc. Cambridge Philos. Soc. **53** (1957), 539–540.
- [4] L. Fukshansky, *Small zeros of quadratic forms with linear conditions*, J. Number Theory **108** (2004), 29–43.
- [5] L. Fukshansky, *Small zeros of quadratic forms over  $\overline{\mathbb{Q}}$* , Int. J. Number Theory **4** (2008), 503–523.
- [6] É. Gaudron and G. Rémond, *Corps de Siegel*, J. reine angew. Math. to appear, 61 pages. DOI: 10.1515/crelle-2014-0096
- [7] É. Gaudron and G. Rémond, *Espaces adéliques quadratiques*, 34 pages, preprint available on <http://math.univ-bpclermont.fr/~gaudron>
- [8] D. Masser, *How to solve a quadratic equation in rationals*, Bull. London Math. Soc. **30** (1998), 24–28.
- [9] H.P. Schlickewei, *Kleine Nullstellen homogener quadratischer Gleichungen*, Monatsh. Math. **100** (1985), 35–45.
- [10] J. Vaaler, *Small zeros of quadratic forms over number fields*, Trans. Amer. Math. Soc. **302** (1987), 281–296.